

# Polynomial Time Recognition of Uniform Cocircuit Graphs

Ricardo Gómez Aiza, Juan José Montellano Ballesteros,  
Ricardo Strausz

*Instituto de Matemáticas  
Universidad Nacional Autónoma de México  
México D.F., Mexico*

Kolja Knauer<sup>1,2</sup>

*Institut für Mathematik  
Technische Universität Berlin  
Berlin, Germany*

---

## Abstract

We present an algorithm which takes a graph as input and decides in polynomial time if the graph is the cocircuit graph of a uniform oriented matroid. In the affirmative case the algorithm returns the set of signed cocircuits of the oriented matroid.

*Keywords:* Oriented matroid, cocircuit graph, recognition algorithm, polynomial algorithm.

---

<sup>1</sup> Supported by the Research Training Group *Methods for Discrete Structures* (DFG-GRK 1408). Part of this work was done during visits to the Departments of Mathematics of UNAM and CINVESTAV, Mexico City.

<sup>2</sup> Email: [knauer@math.tu-berlin.de](mailto:knauer@math.tu-berlin.de)

# 1 Introduction

The cocircuit graph is a natural combinatorial construction associated with oriented matroids. In the case of pseudoline-arrangements, i.e., uniform rank 3 oriented matroids, its vertices are the intersection points of the lines and two points share an edge if they are connected by a line segment which does not intersect other lines. More generally, the Topological Representation Theorem of Folkman and Lawrence [4] says that every oriented matroid can be represented as an arrangement of pseudospheres. The cocircuit graph is the 1-skeleton of this arrangement.

For the uniform case, Montellano-Ballesteros and Strausz [6] provide a graph theoretical characterization of cocircuit graphs. Cordovil, Fukuda and Guedes de Oliveira [2] show that a uniform oriented matroid is basically determined by its cocircuit graph (up to isomorphism and reorientation).

After introducing basic notions of oriented matroids we provide an algorithm inspired by [2] which given a graph  $G$  decides in polynomial time if  $G$  is the cocircuit graph of a uniform oriented matroid. In the affirmative case the algorithm returns the set of signed cocircuits of the oriented matroid.

Given a finite ground set  $E$  we define a *signed set*  $X \subseteq E$  as an *underlying set*  $\underline{X}$  with a bipartition  $(X^+, X^-)$  into a *positive* and a *negative* part. The parts may be empty. By  $X^0$  we refer to the *zero-support*  $E \setminus \underline{X}$ . *Reorienting*  $X$  on  $A \subseteq E$  yields the signed set  ${}_A X := ((X^+ \setminus A) \cup (X^- \cap A), (X^- \setminus A) \cup (X^+ \cap A))$ . By  $-X$  we refer to  ${}_E X$ . Given signed sets  $X, Y$  we denote by  $S(X, Y) := (X^+ \cap Y^-) \cup (X^- \cap Y^+)$  their *separator*.

We define an *oriented matroid* as a pair  $\mathcal{M} = (E, \mathcal{C}^*)$  of a ground set  $E$  and a multiset  $\mathcal{C}^*$  of signed sets called *cocircuits* satisfying the following axioms:

- (C1)  $\emptyset \notin \mathcal{C}^*$
- (C2)  $X \in \mathcal{C}^* \Rightarrow -X \in \mathcal{C}^*$
- (C3)  $X, Y \in \mathcal{C}^*$  and  $\underline{X} = \underline{Y} \Rightarrow X = \pm Y$
- (C4)  $X, Y \in \mathcal{C}^*$  and  $\underline{X} \neq \underline{Y}$  and  $e \in S(X, Y) \Rightarrow$  there is  $Z \in \mathcal{C}^*$  with  $Z^+ \subseteq X^+ \cup Y^+$  and  $Z^- \subseteq X^- \cup Y^-$  and  $e \in Z^0$ .

The *cocircuit graph*  $G_{\mathcal{M}}$  of an oriented matroid  $\mathcal{M} = (E, \mathcal{C}^*)$  has as vertex set  $\mathcal{C}^*$  and  $X, Y \in \mathcal{C}^*$  share an edge if and only if  $S(X, Y) = \emptyset$  and the symmetric difference  $|X^0 \Delta Y^0| = 2$ .

A *uniform oriented matroid*  $\mathcal{M} = (E, \mathcal{C}^*)$  of *order*  $n$  and *rank*  $r$  is an oriented matroid with  $|E| = n$  and  $\mathcal{C}^*$  having exactly the subsets of size  $n - r + 1$  as underlying sets. For more about oriented matroids, see [1].

## 2 The Algorithm

We are given a (simple, connected) graph  $G = (V, A)$  and want to test in polynomial time whether  $G$  is the cocircuit graph of a uniform oriented matroid. In the affirmative case we assign a signed set  $X(v)$  to every vertex  $v$  such that  $X(V)$  is the set of cocircuits of  $\mathcal{M}$  with  $G = G_{\mathcal{M}}$ . The algorithm has the following structure:

- (A) Determine the parameters  $n$  and  $r$
- (B) Construct the set of great cycles  $\mathcal{G}$  (defined below)
- (C) Assign zero-supports  $X^0$  to vertices and great cycles
- (D) Assign signed sets  $X(v) = (X^+(v), X^-(v))$  to vertices
- (E) Check the cocircuit axioms

Each of these steps works if  $G$  is the cocircuit graph of a uniform oriented matroid and outputs that  $G$  is not, otherwise. The main idea of step (C) is contained in [2].

**(A) Determine the parameters.** Cocircuit graphs are regular and antipodal. These properties are checked here.

- Check if  $G$  is regular. If so, let  $\delta$  denote the degree and set  $r := \frac{\delta}{2} + 1$ .
- Check if for every  $v \in V$  there exists a unique  $v^- \in V$  with  $\text{dist}(v, v^-) = \text{diam}(G)$ . In the affirmative case set  $n := \text{diam}(G) + r - 1$ .
- Check if  $|V| = 2 \binom{n}{r-1}$ .
- If any of the above tests results negative,  $G$  is not a cocircuit graph.

All the checked properties are necessary conditions for  $G$  to be a cocircuit graph of some uniform  $\mathcal{M}$ . Moreover  $n$  and  $r$  must be the order and the rank of the oriented matroid of  $\mathcal{M}$ , respectively. The runtime here is bounded from above by calculating a shortest path matrix for every starting vertex in the second step. This can be solved in  $\mathcal{O}(V^3)$  by applying Dijkstra's Algorithm [3]  $|V|$  times.

**(B) Construct the set of great cycles** A set  $C \subseteq \mathcal{C}^*$  is called a *great cycle* with zero-support  $X^0(C)$  if it consists of the cocircuits whose zero-support contains the  $(r-1)$ -set  $X^0(C)$ . The set of great cycles is denoted by  $\mathcal{G}$ . The following part of the algorithm computes  $\mathcal{G}$ .

- For every  $v \in V(G)$  and every neighbor  $w \in N(v)$ . if the edge  $\{v, w\}$  is not contained in any  $C \in \mathcal{G}$  then

- compute a shortest path  $P$  from  $w$  to  $v^-$ ,
- set  $C' := (\{v, w\}, P, \{v^-, w^-\}, P^-)$ ,
- add  $C'$  to  $\mathcal{G}$ .
- If  $\mathcal{G}$  is no partition of the edge set then  $G$  is not a cocircuit graph.

For the correctness of this step we need:

**Lemma 2.1** *The set of great cycles  $\mathcal{G}$  of a cocircuit graph  $G$  partitions the edge set. Moreover the edge  $\{v, w\}$  is contained in the cycle  $C'$  constructed in the algorithm.*

The lemma tells us in particular that  $|\mathcal{G}||C| = |A| = \frac{\delta}{2}|V| = 4(r-1) \binom{n}{r-1}$  and that  $|C| = 2\text{diam}(G) = 2(n-r+1)$ . Hence the runtime  $|V|\delta|C||V|^2$  is bounded by  $4(r-1)(n-r+1)|V|^3 \in \mathcal{O}(|V|^4)$ .

**(C) Assign zero-supports to vertices and great cycles.** In this part we assign zero supports to great cycles and vertices. We use the incidence structure on  $\mathcal{G}$ , i.e., let  $J$  be the graph with vertex set  $\mathcal{G}$  where  $C$  and  $C'$  share an edge if they have a vertex in common in  $G$ . Note that  $J$  is isomorphic to the *Johnson Graph*  $J(n, n-r+2)$ , see [5] for a definition. During the algorithm we denote by  $\mathcal{P}$  the set of cycles with already assigned zero-support. Cycles with all their vertices having a zero-support already are in  $\mathcal{T}$ . We assign supports starting from a vertex  $v$ , then all cycles containing  $v$ . These cycles form a clique in  $J$ . By  $\mathcal{G}_i, \mathcal{P}_i, \mathcal{T}_i$  we denote the respective sets at distance  $i$  from this clique. We write  $[k]$  for  $\{1, \dots, k\}$ .

- Take any  $v \in V$  and set  $X^0(v) := [r-1]$ .
- For every  $C_i$  in the set  $\{C_1, \dots, C_{r-1}\}$  of great cycles containing  $v$ ,
  - set  $X^0(C_i) := [r-1] \setminus \{i\}$ ,
  - add  $C_i$  to  $\mathcal{P}_0$ .
- Take any  $\tilde{C} = (v_1, \dots, v_{2(n-r+1)})$  from  $\mathcal{P}_0$ .
- Set  $X^0(v_{(n-r+1)+i}) := X^0(v_i) := X^0(C) \cup \{i\}$  for every  $i \in [n-r+1]$ .
- Add  $\tilde{C}$  to  $\mathcal{T}_0$  and  $\mathcal{T}_{-1}$ .
- Set  $i := 0$  and until all  $V$  is labelled repeat:
  - T-loop: For every  $C \in \mathcal{P}_i$  take  $\tilde{C} \in \mathcal{T}_{i-1}$  such that  $C \cap \tilde{C} \neq \emptyset$ .  
For every  $v \in C$  find a  $C' \in \mathcal{G}$  with  $v \in C'$  and  $C' \cap \tilde{C}$  containing a  $w$ :  
set  $X^0(v) := X^0(C) \cup (X^0(w) \setminus X^0(\tilde{C}))$ ,  
add  $C$  to  $\mathcal{T}_i$ .
  - P-loop: For every  $\tilde{C} \in \mathcal{T}_i$  and  $v \in \tilde{C}$ :  
If there exists an unlabeled great cycle  $C$  containing  $v$  look for another

$w \in C$  with known  $X^0(w)$ ,  
 set  $X^0(C) := X^0(v) \cap X^0(w)$ ,  
 add  $C$  to  $\mathcal{P}_{i+1}$ .

- If any vertex receives several different labels  $G$  is not a cocircuit graph.
- Increase  $i$  by one.

The runtime is bounded by the T-loop  $|\mathcal{G}|(|C|\delta + |C|\delta|C||C|) \in \mathcal{O}(V^3)$ .  
 For the existence of  $C'$  in the T-loop we need:

**Lemma 2.2** *Let  $C \in \mathcal{G}_i$  and  $C' \in \mathcal{G}_{i-1}$  be intersecting. Then for every  $v \in C$  there is a  $C'' \in \mathcal{G}_i$  that contains  $v$  and intersects with  $C'$ .*

For the existence of  $w$  in the P-loop we need:

**Lemma 2.3** *For every  $C \in \mathcal{G}_i$  there are at least two  $C', C'' \in \mathcal{G}_{i-1}$  which intersect with  $C$ .*

**(D) Assign signed sets to vertices.** We will label the vertices with signed sets. As in (C) we start with cycles  $\mathcal{S}_0$  which form a clique in  $J$ . Then we label cycles at increasing distance of  $\mathcal{S}_0$ . Since we calculated zero-supports in (C) we set  $\underline{X}(v) := E \setminus X^0(v)$  for every  $v \in V$ .

- Take any  $v \in V$  and set  $X(v) := (\underline{X}(v), \emptyset)$ .
- For all great cycles  $C \ni v$  and  $w, w' \in N(v) \cap C$  set  $X(w) := (X^+(v) \cap \underline{X}(w), \underline{X}(w) \setminus X^+(v))$  and  $X(w') := (\underline{X}(w), \emptyset)$ .
- Add all  $C \in \mathcal{G}$  containing  $v$  to  $\mathcal{S}_0$ .
- Set  $i := 0$  and repeat until all  $V$  is signed:
  - For every great cycle  $C \in \mathcal{S}_i$ 
    - get two signed non-antipodal vertices  $v, w \in C$  and take  $u \in N(v) \cap C$  such that  $0 < d(u, w) < d(u, w^-)$ , i.e.,  $u, v, w$  lie on the same half of  $C$ , sign  $u$  identically to  $v$  on  $\underline{X}(v) \cap \underline{X}(u)$  and take the signing of  $X(w)$  on  $e = \underline{X}(u) \setminus \underline{X}(v)$ .
    - This way we sign all vertices in  $C$ .
    - If a vertex receives two different labels  $G$  is not a cocircuit graph.
    - Add all unsigned cycles  $C' \cap C \neq \emptyset$  to  $\mathcal{S}_{i+1}$ .
  - Increase  $i$  by one.

The runtime is  $|\mathcal{G}||C|^2 \in \mathcal{O}(|V|^4)$ . We need Lemma 2.3 for the existence of  $v, w \in C$ . For the correctness of the signing we have:

**Lemma 2.4** *Take a great cycle  $C$  of a uniform cocircuit graph,  $X, Y \in C$ ,  $\underline{X} \neq \underline{Y}$  and let  $(X = X_0, \dots, X_k = Y)$  be the shortest  $(X, Y)$ -path in  $C$  and  $e \in \underline{X} \cap \underline{Y}$ . Then  $e \in S(X, Y)$  if and only if there is  $X_i^0 \ni e$ .*

**(E) Check the cocircuit axioms.** Here we check if the signed sets we assigned to the vertices satisfy (C4) of the cocircuit axioms. The other axioms are satisfied by construction.

- For every two non-antipodal vertices  $u, v \in V$  and every element in the separator of their signed labels  $e \in S(X(u), X(v))$  check if there exists a  $w \in V$  with  $X(w)^+ \subseteq X(u)^+ \cup X(v)^+$  and  $X(w)^- \subseteq X(u)^- \cup X(v)^-$  and  $e \in X(w)^0$ .

Here the runtime is  $|V|^3(n - r + 1) \in \mathcal{O}(|V|^4)$ . The correctness is obvious. The last step of the algorithm could also be done by checking whether the constructed  $\mathcal{C}^*$  is a certain metrical and antipodal embedding into the  $(r - 1)$ -dual of the  $n$ -cube, see [6].

Thus we have the following result:

**Theorem 2.5** *The preceding algorithm checks if a given graph  $G = (V, A)$  is the cocircuit graph of a uniform oriented matroid  $\mathcal{M}$  and constructs such  $\mathcal{M}$  in the affirmative case in time  $\mathcal{O}(|V|^4)$ .*

## References

- [1] Björner, A., M. Las Vergnas, B. Sturmfels, N. White and G. M. Ziegler, “Oriented matroids,” *Encyclopedia of Mathematics and its Applications* **46**, Cambridge University Press, Cambridge, 1993.
- [2] Cordovil, R., K. Fukuda and A. Guedes de Oliveira, *On the cocircuit graph of an oriented matroid*, *Discrete Comput. Geom.* **24** (2000), pp. 257–265, the Branko Grünbaum birthday issue.
- [3] Dijkstra, E. W., *A note on two problems in connexion with graphs*, *Numer. Math.* **1** (1959), pp. 269–271.
- [4] Folkman, J. and J. Lawrence, *Oriented matroids*, *J. Combin. Theory Ser. B* **25** (1978), pp. 199–236.
- [5] Godsil, C. and G. Royle, “Algebraic graph theory,” *Graduate Texts in Mathematics* **207**, Springer-Verlag, New York, 2001.
- [6] Montellano-Ballesteros, J. J. and R. Strausz, *A characterization of cocircuit graphs of uniform oriented matroids*, *J. Combin. Theory Ser. B* **96** (2006), pp. 445–454.