Simple Treewidth

Kolja Knauer¹

Technical University Berlin knauer@math.tu-berlin.de Joint work with Torsten Ueckerdt².

1 Introduction

A *k*-tree is a graph that can be constructed starting with a (k+1)-clique and in every step attaching a new vertex to a *k*-clique of the already constructed graph. The *treewidth* tw(*G*) of a graph *G* is the minimum *k* such that *G* is a *partial k*-tree, i.e., *G* is a subgraph of some *k*-tree [7].

We consider a variation of treewidth, called *simple treewidth*. A simple k-tree is a k-tree with the extra requirement that there is a construction sequence in which no two vertices are attached to the same k-clique. (Equivalently, a k-tree is simple if it has a tree representation of width k in which every (k - 1)-set of subtrees intersects at most 2 tree-vertices.) Now, the simple treewidth $\operatorname{stw}(G)$ of G is the minimum k such that G is a partial simple k-tree, i.e., G is a subgraph of some simple k-tree.

We have encountered simple treewidth as a natural parameter in questions concerning geometric representations of graphs, i.e., representing graphs as intersection graphs of geometrical objects where the quality of the representation is measured by the complexity of the objects. E.g., we have shown that the maximal *interval-number*([3]) of the class of treewidth k graphs is k + 1, whereas for the class of simple treewidth k graphs it is k, see [6]. Another example is the *bend-number*([4]), which for treewidth 3 graphs is 4 and for simple treewidth 3 graphs is 3, see [5] and a corresponding statement for higher values is conjectured in [4].

The aim of this note is to compare these two parameters and to motivate simple treewidth by indicating that it endows treewidth with a topological flavor. We pose several interesting open problems.

 $^{^1\}mathrm{The}$ research was supported by DFG grant FE-340/8-1 as part of ESF project GraDR EUROGIGA

²Karlsruhe Institute of Technology. ueckerdt@gmail.com.

2 Properties of simple treewidth

Let us first observe, that both parameters cannot differ too much.

Observation 2.1. For every G we have $tw(G) \le stw(G) \le tw(G) + 1$.

Proof. The first inequality is clear. For the second inequality we show that every k-tree G is a subgraph of a simple (k + 1)-tree H. Whenever in the construction sequence of G several vertices $\{v_1, \ldots, v_n\}$ are attached to the same k-clique C we consider $C \cup \{v_1\}$ as a (k+1)-clique in the construction sequence for H. Attaching v_i to C can be interpreted as attaching v_i to $C \cup \{v_{i-1}\}$ and omitting the edge $\{v_{i-1}, v_i\}$. This way we avoid that several vertices are attached to the same k-clique by considering (k+1)-cliques. \Box

Simple treewidth endows the notion of treewidth with a more topological flavor, as indicated for small k in the table below:

	≤ 1	≤ 2	≤ 3
stw	paths	outerplanar	planar & tw ≤ 3 , [2]
tw	forest	series-parallel	$tw \le 3$

A linkless embedding of G is an embedding into \mathbb{R}^3 with the property that no two cycles of G form a link, see [9].

Observation 2.2. If $stw(G) \leq 4$ then G has a linkless embedding.

Proof. It suffices to show that simple 4-trees have linkless embeddings, since edge-deletion does not destroy the linkless embedding. Therefore consider K_5 embedded in \mathbb{R}^3 as a tetrahedron plus a vertex in its interior. In each step of the construction sequence every available 4-clique is represented by a tetrahedron with empty interior, where we insert the new vertex. It is easy to see that the resulting embedding is linkless.

Non-simple 4-trees do not have linkless embeddings, which is easy to see using the forbidden-minor chracterization of linkless embeddable graphs [8].

Problem 2.3. $\operatorname{stw}(G) \leq 4 \Leftrightarrow G$ is linkless embeddable and $\operatorname{tw}(G) \leq 4$.

Simple treewidth also has connections to discrete geometry. In [1] a stacked polytope is defined to be a polytope admitting a triangulation whose dual graph is a tree. In that paper it is proved that a full-dimensional polytope $P \subset \mathbb{R}^d$ is stacked if and only if $\operatorname{tw}(G(P)) \leq d$, were G(P) denotes the 1-skeleton of P. Indeed, we strongly suspect:

Problem 2.4. A graph G is the 1-skeleton of a stacked d-polytope if and only if stw(G) = d and G is d-connected.

One can show that the class of simple treewidth at most k graphs is minor-closed. A proof of the following statement would imply that for planar graphs with treewidth at least 3 treewidth and simple treewidth coincide.

Problem 2.5. If G has no $K_{3,k}$ -minor and tw(G) = k then stw(G) = k.

References

- Alexander Below, Jesús A. De Loera, and Jürgen Richter-Gebert. The complexity of finding small triangulations of convex 3-polytopes. J. Algorithms, 50(2):134–167, 2004. SODA 2000 special issue.
- [2] Ehab S. El-Mallah and Charles J. Colbourn. On two dual classes of planar graphs. *Discrete Math.*, 80(1):21–40, 1990.
- [3] Frank Harary and William T. Trotter, Jr. On double and multiple interval graphs. J. Graph Theory, 3(3):205–211, 1979.
- [4] Daniel Heldt, Kolja Knauer, and Torsten Ueckerdt. Edge-intersection graphs of grid paths: the bend-number. *ArXiv e-prints*, September 2010.
- [5] Daniel Heldt, Kolja Knauer, and Torsten Ueckerdt. On the bend-number of planar and outerplanar graphs. In *LATIN*, pages 458–469, 2012.
- [6] Kolja Knauer and Torsten Ueckerdt. Three ways to cover a graph. ArXiv e-prints, May 2012.
- [7] Neil Robertson and Paul D. Seymour. Graph minors—a survey. In Surveys in combinatorics 1985 (Glasgow, 1985), volume 103 of London Math. Soc. Lecture Note Ser., pages 153–171. Cambridge Univ. Press, Cambridge, 1985.
- [8] Neil Robertson, Paul D. Seymour, and Robin Thomas. Sachs' linkless embedding conjecture. J. Combin. Theory Ser. B, 64(2):185–227, 1995.
- [9] Horst Sachs. On a spatial analogue of Kuratowski's theorem on planar graphs—an open problem. In *Graph theory (Lagów, 1981)*, volume 1018 of *Lecture Notes in Math.*, pages 230–241. Springer, Berlin, 1983.