

# Graphs of Some CAT(0) Complexes

Victor Chepoi<sup>1</sup>

*SFB343 Diskrete Strukturen in der Mathematik, Universität Bielefeld,  
D-33615 Bielefeld, Germany*

Received February, 1998; accepted October 29, 1999

In this note, we characterize the graphs (1-skeletons) of some piecewise Euclidean simplicial and cubical complexes having nonpositive curvature in the sense of Gromov's CAT(0) inequality. Each such cell complex  $K$  is simply connected and obeys a certain flag condition. It turns out that if, in addition, all maximal cells are either regular Euclidean cubes or right Euclidean triangles glued in a special way, then the underlying graph  $G(K)$  is either a median graph or a hereditary modular graph without two forbidden induced subgraphs. We also characterize the simplicial complexes arising from bridged graphs, a class of graphs whose metric enjoys one of the basic properties of CAT(0) spaces. Additionally, we show that the graphs of all these complexes and some more general classes of graphs have geodesic combings and bicomings verifying the 1- or 2-fellow traveler property.

© 2000 Academic Press

## 1. INTRODUCTION

The idea of formulating nonpositive curvature for general geodesic metric spaces goes back to work of Alexandrov and Busemann [1, 2, 18]. Gromov [31] introduced the term CAT(0) for simply connected geodesic metric spaces in which the geodesic triangles are thinner than in the Euclidean plane. CAT(0) spaces have many nice equivalent definitions, reflecting different geometric features [6, 16, 17, 47]. A particularly important class of such spaces is formed by CAT(0) simplicial and cubical complexes [16]. Gromov [31] has shown that CAT(0) cubical complexes can be characterized in a purely combinatorial way; see [16, 17, 20, 21, 47] for related results and applications. Further properties of CAT(0) cubical complexes

<sup>1</sup> Current address: Laboratoire d'Informatique de Marseille, Université d'Aix Marseille II, Faculté des Sciences de Luminy, F-13288 Marseille Cedex 8, France.



have been presented in the paper of Sageev [51], who called them cubings. Niblo and Reeves [44] further investigate the properties of cubings and show how to find a bicombing of Caley graphs of their fundamental groups. On the other hand, Gersten and Short [29, 30] use the CAT(0) singular disk diagrams to provide an automatic structure for fundamental groups of some two-dimensional piecewise Euclidean complexes. Other examples of CAT(0) polygonal complexes were given in [16, 17, 31, 47].

In this article we show that certain CAT(0) simplicial or cubical cell complexes can be characterized by their underlying graphs. In particular, we show that the graphs of cubings are exactly the median graphs. They and the associated cubical complexes have many nice geometric, convexity, and algebraic properties already known before [11, 35, 42, 43, 55]. This allows us to simplify some results from [51]. Also we characterize the graphs of CAT(0) two-dimensional complexes whose cells are right triangles as graphs in which all isometric cycles have length four and do not contain two forbidden subgraphs. Next we present a characterization of simplicial complexes arising from graphs in which all isometric cycles have length three (so-called bridged graphs). It is known that bridged graphs share one of the main properties of CAT(0) spaces, namely the convexity of the neighborhoods of convex sets. We present examples of simplicial complexes with regular cells arising from bridged graphs which are not CAT(0). Nevertheless, in few important cases we obtain CAT(0) complexes. An important feature of bridged graphs is that their graph-metric admits a bicombing satisfying the  $k$ -fellow traveler condition for  $k \leq 2$ . This property (for different  $k \leq 3$ ) is shared by all classes of graphs arising in this note, in particular, by median graphs and Helly graphs (a class of graphs containing the Caley graphs of fundamental groups of cubings).

The paper is organized as follows. In Section 2 we establish the notation and definitions used in the rest of the paper. In Section 3 we recall the basic characterizations of CAT(0) complexes; in Section 4 we introduce the classes of graphs we deal with and recall their main properties. In Section 5 we briefly discuss the minimal disks diagrams which are used in the subsequent proofs. In Section 6 we establish the correspondence between the cubings and median complexes. Using this result we derive some properties of cubings. In Section 7 we focus our attention on CAT(0) 2-complexes whose cells are right triangles (we call them folder complexes). We characterize folder complexes and present their basic properties which can be derived from the specificity of their skeletons. In Section 8 we establish that the flag complexes whose graphs are bridged are exactly the simply connected simplicial complexes in which the links of vertices do not contain induced 4-cycles and 5-cycles. Finally, in Section 9 we present the results about combings and bcombings of the underlying graphs of complexes under consideration.

## 2. PRELIMINARIES

This section summarizes some preliminaries from topology and metric spaces. Basic notions and results from topology can be found in any textbook on combinatorial topology (e.g., [52]).

### 2.1. Simplicial and Cubical Complexes

Let  $K$  be an abstract *simplicial complex*, i.e., a collection of sets (called *simplexes*) such that  $\sigma \in K$  and  $\sigma' \subseteq \sigma$  implies  $\sigma' \in K$ . A *cubical complex* is a set  $K$  of cubes of any dimensions which is closed under taking subcubes and nonempty intersections. We denote by  $V(K)$  and  $E(K)$  the *vertex set* and the *edge set* of  $K$ , namely, the set of all 0-simplexes and 1-simplexes of  $K$ . The pair  $(V(K), E(K))$  is called the (*underlying*) *graph* or the *1-skeleton* of  $K$  and is denoted by  $G(K)$  or  $K^1$  (the underlying graph of a cubical complex is defined similarly). More generally, for each  $r \geq 0$  the *r-skeleton* of  $K$  is the simplicial complex  $K^r = \{\sigma \in K : \#\sigma \leq r + 1\}$ . A *geometric realization*  $|K|$  of  $K$  is a polyhedron obtained by the following construction. Let  $\phi: V(K) \rightarrow \mathbb{E}^k$  be an injection such that for any two simplexes  $\sigma', \sigma''$ ,  $\text{conv } \phi(\sigma') \cap \text{conv } \phi(\sigma'') = \text{conv } \phi(\sigma' \cap \sigma'')$ . Then  $|K| = \cup\{\text{conv } \phi(\sigma) : \sigma \in K\}$ .

The *link* of a simplex  $\sigma$  in  $K$ , denoted  $\text{link}(\sigma, K)$  is the abstract simplicial complex consisting of all simplexes  $\sigma'$  such that  $\sigma \cap \sigma' = \emptyset$  and  $\sigma \cup \sigma' \in K$ . The *star*  $\text{star}(\sigma, K)$  of  $\sigma$  is the set of all simplexes containing  $\sigma$ . (The star of a cube in a cubical complex is defined analogously.) A simplicial complex  $K$  is a *flag complex* if any set of vertices is included in a member of  $K$  whenever each pair of its vertices is contained in a member of  $K$ . In the theory of hypergraphs this condition is called conformality. The link of any simplex in a flag complex is again a flag complex. A flag complex can be recovered by its underlying graph  $G(K)$ : the complete subgraphs of  $G(K)$  are exactly the simplexes of  $K$ .

A topological space  $X$  is *1-connected* if every continuous mapping  $f$  of the one-dimensional sphere  $S^1$  into  $X$  can be extended to a continuous mapping of the disk  $D^2$  with boundary  $S^1$  into  $X$ . A simplicial complex is *simply connected* if its underlying space  $|K|$  is 1-connected. It is well known that  $K$  is simply connected if and only if  $|K^2|$  is 1-connected.

A *cycle*  $C$  is a sequence  $(e_1, \dots, e_m)$  of edges of the graph  $G(K)$  such that the edges  $e_i$  and  $e_{i+1(\text{mod } m)}$  are incident and  $e_{i-1} \cap e_i \neq e_i \cap e_{i+1(\text{mod } m)}$  for  $i = 1, \dots, m$ . If  $e_i$  and  $e_j$  are incident if and only if  $j = i + 1(\text{mod } m)$  or  $i = j + 1(\text{mod } m)$  then  $C$  is called a *simple cycle* of the 1-skeleton of  $K$ . Equivalently, one can define a cycle as a sequence  $(v_1, \dots, v_m, v_1)$  of vertices such that the consecutive vertices are adjacent. If  $K$  is simply connected, then every cycle  $C$  of  $G(K)$  is null-homologous; i.e., it can be writ-

ten as a modulo 2 sum of 2-simplexes. Moreover,  $C$  can be transferred to a point by means of combinatorial deformations; for definitions see, for example, [52, pp. 163–167].

## 2.2. Piecewise Euclidean Cell Complexes

A *Euclidean cell* is a convex polytope in some Euclidean space. By a *piecewise Euclidean (PE) cell complex* we shall mean a space  $\mathcal{X}$  formed by gluing together Euclidean cells via isometries of their faces, together with the subdivision of  $\mathcal{X}$  into cells; for a precise definition, see, e.g., [17, 20, 47]. Additionally we assume that the intersection of two cells either is empty or a single face of each of cells. If all cells of  $\mathcal{X}$  are Euclidean simplexes (respectively, cubes), we say that  $\mathcal{X}$  is a *simplicial* (respectively, *cubical*) *cell complex*. With each simplicial or cubical complex  $\mathcal{X}$  one can associate in a canonical way an abstract simplicial or cubical complex  $K = K(\mathcal{X})$ . If all cells of  $\mathcal{X}$  are regular Euclidean simplexes or cubes (see Sections 7 and 8), then  $\mathcal{X} = |K|$ . A *piecewise spherical (PS) cell complex* is defined similarly using spherical cells. Such complexes arise naturally as links of cells and points (see [17, 20, 47] for more details).

Suppose that  $P$  is an Euclidean cell of  $\mathbb{E}^n$  and  $x \in P$ . Then  $\text{Link}(x, P)$  is the subset of the unit sphere  $S^{n-1}$  centered at  $x$  consisting of all unit vectors which originate in  $x$  and pointing into  $P$ . If  $P$  is a regular  $n$ -cube or a regular  $n$ -simplex and  $x$  is a vertex of  $P$ , then  $\text{Link}(x, P)$  is a regular spherical simplex on  $n$  vertices. The distance between any two vertices of  $\text{Link}(x, P)$  is  $\pi/2$  in the first case and  $\pi/3$  in the second case (recall that the spherical distance between two points  $u$  and  $v$  is the angle at  $x$  between the line segments joining  $x$  to  $u$  and  $v$ , respectively). Now suppose that  $P$  is a cell in  $\mathcal{X}$ ,  $x \in P$ , and  $f: P \rightarrow \mathbb{E}^n$  is a homeomorphism onto a Euclidean cell of  $\mathbb{E}^n$ . Put  $\text{Link}(x, P) = \text{Link}(f(x), f(P))$ , and  $\text{Link}(x, \mathcal{X}) = \cup \text{Link}(x, P)$ , where the union is taken over all cells  $P$  containing  $x$ . Then  $\text{Link}(x, \mathcal{X})$  is a piecewise spherical cell complex, called the *link* of  $x$  in  $\mathcal{X}$ . If  $\mathcal{X}$  is a simplicial PE complex and  $x$  is a vertex of  $K$ , then the abstract simplicial complex associated with  $\text{Link}(x, \mathcal{X})$  coincides with  $\text{link}(x, K(\mathcal{X}))$ .

If  $P$  is a Euclidean cell and  $F$  is a  $k$ -face of  $P$ , then the set of unit vectors normal to the  $k$ -plane supported by  $F$  and pointing into  $P$  form  $\text{Link}(F, P)$ . If  $P$  is a cell of  $\mathcal{X}$  and  $F$  is a face of  $P$ , then set  $\text{Link}(F, P) = \text{Link}(f(F), f(P))$  and  $\text{Link}(F, \mathcal{X}) = \cup \text{Link}(F, P)$ , where the union is taken over all cells containing  $F$  (if  $F \subseteq Q \subseteq P$ , then we identify  $\text{Link}(F, Q)$  with the corresponding face of  $\text{Link}(F, P)$ ). The PS complex  $\text{Link}(F, \mathcal{X})$  is called the *link of  $F$  in  $\mathcal{X}$* .

## 2.3. Metric Spaces

Let  $(X, d)$  be a metric space. The closed *ball* of center  $x$  and radius  $r \geq 0$  is denoted by  $B_r(x)$ . More generally, the closed  *$r$ -neighborhood* of a set  $S$

is the set

$$B_r(S) = \{x \in V : d(x, S) \leq r\},$$

where  $d(x, S) = \inf\{d(x, y) : y \in S\}$ . A *path* is a map from a segment  $[a, b]$  to  $X$ . A *geodesic* joining two points  $x, y \in X$  is a map  $g$  from  $[a, b]$  to  $X$  such that  $g(a) = x, g(b) = y$  and  $d(g(t), g(t')) = |t - t'|$  for all  $t, t' \in [a, b]$  (in particular  $d(x, y) = |a - b|$ ). A *geodesic segment* in  $X$  is the subset that is the image of a geodesic. A *geodesic metric space* is a metric space in which every pair of points can be joined by a geodesic segment. Geodesic metric spaces sometimes are called *convex*: for any points  $x, y \in X$  and nonnegative real numbers  $r(x)$  and  $r(y)$  satisfying  $d(x, y) \leq r(x) + r(y)$ , there is a point  $z$  such that  $d(x, z) \leq r(x)$  and  $d(y, z) \leq r(y)$ . More generally, a metric space  $X$  is called *hyperconvex* if for any collection of closed balls in  $X, B_{r_i}(x_i), i \in I,$  satisfying the condition that  $d(x_i, x_j) \leq r_i + r_j$  for all  $i, j \in I,$  the intersection  $\bigcap_{i \in I} B_{r_i}(x_i)$  is nonempty; i.e., the family of balls of  $X$  has the Helly property. The notion of hyperconvex spaces has been introduced by Aronszajn and Panitchpakdi [4] who proved that hyperconvex spaces are exactly the absolute retracts (alias injective spaces). To be more precise, here are the basic notions: a metric space  $(X, d)$  is *isometrically embedded* into a metric space  $(Y, d')$  if there is a map  $h: X \rightarrow Y$  such that  $d'(h(x), h(y)) = d(x, y)$  for all  $x, y \in X$ . In this case we say that  $X$  is a *subspace* of  $Y$  and that  $Y$  is an *extension* of  $X$ . Now, a *retraction*  $h: Y \rightarrow X$  from a metric space  $(Y, d')$  to a subspace  $X$  is an idempotent ( $h(x) = x$  for any  $x \in X$ ) nonexpansive ( $d'(h(x), h(y)) \leq d'(x, y)$  for any  $x, y \in Y$ ) map; its image  $X$  is called a *retract* of  $Y$ . A metric space  $(X, d)$  is an *absolute retract* if  $X$  is a retract of every metric space in which  $X$  embeds isometrically.

The *interval*  $I(x, y)$  between two points  $x$  and  $y$  is the set

$$I(x, y) = \{z \in X : d(x, y) = d(x, z) + d(z, y)\}.$$

A subset  $Y \subseteq X$  is *convex* if  $Y$  includes every interval  $I(x, y)$  with  $x, y$  in  $Y$ . A convex set  $Y$  whose complement  $X - Y$  is convex is called a *halfspace*. A subset  $Y$  of a metric space is called *gated* [26] if for any point  $x \notin Y$  there exists a (unique) point  $x_Y \in Y$  (the *gate* for  $x$  in  $Y$ ) such that  $d(x, z) = d(x, x_Y) + d(x_Y, z)$  for all  $z \in Y$ . Evidently, every gated set is convex. One can easily show by induction that the family of gated sets has Helly property: if  $S_1, \dots, S_n$  is a collection of pairwise intersecting gated sets then  $\bigcap_{i=1}^n S_i \neq \emptyset$ . Pick a common vertex  $x_j$  of  $S_1, \dots, S_{j-1}, S_{j+1}, \dots, S_n,$  and let  $g$  be the gate of  $x_j$  in  $S_j$ . Since  $g \in I(x_j, x_i)$  for all  $x_i \neq x_j,$  we conclude that  $g \in \bigcap_{i=1}^n S_i$ .

We conclude this section by recalling the following important notion. Fix a point  $b$  and consider a partial order  $\leq_b$  on  $X$  defined as follows:

$$x \leq_b y \quad \text{iff } x \in I(y, b).$$

The relation  $\leq_b$  is called the *base point order* at  $b$  [55].

### 3. CAT(0) CELL COMPLEXES

We will briefly review some definitions and some major results about the geometry of non-positively curved cell complexes. This theory originates from classical papers of Alexandrov, Busemann, Bruhat, Cartan, Hadamard, Tits, Toponogov, and others; for earlier developments see [1, 2, 18, 17]. In the most generality it has been outlined in the seminal paper of Gromov [31]. The detailed proofs of the results are from [31]; further extensions and applications have been presented in a number of papers, among which we cite [6, 16, 20, 47]. For a survey in more depth and background, the reader should refer to the forthcoming book of Bridson and Haefliger [17], whose terminology we follow.

A *geodesic triangle*  $\Delta(x_1, x_2, x_3)$  in a geodesic metric space  $(X, d)$  consists of three points in  $X$  (the vertices of  $\Delta$ ) and a geodesic segment between each pair of vertices (the edges of  $\Delta$ ). A *comparison triangle* for  $\Delta(x_1, x_2, x_3)$  is a triangle  $\Delta(x'_1, x'_2, x'_3)$  in  $\mathbb{E}^2$  such that  $d_{\mathbb{E}^2}(x'_i, x'_j) = d(x_i, x_j)$  for  $i, j \in \{1, 2, 3\}$ . Gromov [31] defines a geodesic metric space  $X$  to be a *CAT(0) space* if all geodesic triangles of  $X$  satisfy the comparison axiom of Cartan–Alexandrov–Toponogov:

*CAT(0)*: Let  $\Delta(x_1, x_2, x_3)$  be a geodesic triangle in  $X$ , and let  $y$  be a point on the side of  $\Delta(x_1, x_2, x_3)$  with vertices  $x_1$  and  $x_2$ . Let  $y'$  denote a unique point on the line segment  $[x'_1, x'_2]$  of the comparison triangle  $\Delta(x'_1, x'_2, x'_3)$ , such that  $d_{\mathbb{E}^2}(x'_i, y') = d(x_i, y)$  for  $i = 1, 2$ . Then  $d(x_3, y) \leq d_{\mathbb{E}^2}(x'_3, y)$ .

The *intrinsic pseudometric* of a PE complex  $\mathcal{X}$  is defined by assuming the distance between two points equal to the greatest lower bound on the length of piecewise linear paths joining them; see [16, 17]. The next theorem is a generalization of some results from [1, 18] for locally finite complexes.

**THEOREM 3.1** [16]. *If  $\mathcal{X}$  is a PE complex with only finitely many isometry types of cells then  $\mathcal{X}$  endowed with the intrinsic pseudometric is a complete geodesic metric space. If  $\mathcal{X}$  is a CAT(0) space then every local geodesic is a geodesic.*

A cell complex  $\mathcal{X}$  satisfies the *link condition* if for every  $x \in \mathcal{X}$  any two points  $y, z$  in the PS complex  $\text{Link}(x, \mathcal{X})$  at distance less than  $\pi$  can be

joined by a *unique geodesic*. The *systole* of  $\mathcal{X}$  is defined to be the infimum of the length of an isometrically embedded circle. The following result collects some basic characterizations of CAT(0) cell complexes (the full list of criteria with detailed proofs is given in [16, 17]; see also [6, 47]). The most part of the following conditions characterize CAT(0) spaces as well.

**THEOREM 3.2** [16, 31]. *Let  $\mathcal{X}$  be a simply connected PE complex with only finitely many isometry types of cells and endowed with the intrinsic metric. Then the following conditions are equivalent:*

- (i)  $\mathcal{X}$  satisfies CAT(0);
- (ii) any two points of  $\mathcal{X}$  can be joined by a unique geodesic segment;
- (iii) if  $\alpha$  and  $\beta$  are geodesic segments in  $\mathcal{X}$  then the function  $f: [0, 1] \rightarrow \mathcal{X}$  given by  $f(t) = d(\alpha(t), \beta(t))$  is convex;
- (iv)  $\mathcal{X}$  satisfies the link condition;
- (v) for each cell  $C$  of  $\mathcal{X}$  the systole of  $\text{Link}(C, \mathcal{X})$  is at least  $2\pi$ .

If the dimensions of all cells of  $\mathcal{X}$  are at most 2, then (i) is equivalent to

$(i^+)$  for any vertex  $x \in \mathcal{X}$  every simple cycle in  $\text{Link}(x, \mathcal{X})$  has length at least  $2\pi$ .

From (iii) one can immediately conclude that in a CAT(0) space the  $r$ -neighborhoods  $B_r(S)$  of convex sets  $S$  are convex. It also implies that a CAT(0) space does not contain isometrically embedded cycles; furthermore, in [17, 20] CAT(0) PE complexes were characterized as spaces of nonpositive curvature without isometrically embedded cycles. The convexity of balls yields that CAT(0) spaces are simply connected, even more, that they are contractible; see, for example, [17, Chap. II].

In the case of cube complexes, Gromov [31] presented a combinatorial characterization of CAT(0) condition (for a proof see [17, 47]).

**PROPOSITION 3.3** [31]. *A cube complex  $\mathcal{X}$  is CAT(0) if and only if  $\mathcal{X}$  is simply connected and for any cube  $C$  the abstract simplicial complex of  $\text{Link}(C, \mathcal{X})$  is a flag complex.*

This condition can be restated as follows: if three  $(k + 2)$ -cubes of  $\mathcal{X}$  share a common  $k$ -cube, and pairwise share common  $(k + 1)$ -cubes, then they are contained in a  $(k + 3)$ -cube of  $\mathcal{X}$ . In [44, 51] the CAT(0) cube complexes were called *cubings*.

A rich class of CAT(0) 2-complexes is formed by PE complexes arising in small cancellation theory [40]. An  $(m, n)$ -complex is a 2-complex  $\mathcal{X}$  in which each face (2-cell) has at least  $m$  sides and for any vertex  $x$  every simple cycle in  $\text{Link}(x, \mathcal{X})$  has at least  $n$  edges. If  $\mathcal{X}$  is a simply connected  $(m, n)$ -complex with  $mn \geq 2(m + n)$  and each face of  $\mathcal{X}$  is a regular Euclidean

polygon, then from Theorem 3.2( $i^+$ ) it follows that  $\mathcal{X}$  is a CAT(0) space. In particular, (6,3), (4,4), and (3,6)-complexes are CAT(0).

Gersten and Short [29, 30] consider four types  $(A_1 \times A_1, A_2, B_2, G_2)$ , of 2-complexes of nonpositive curvature and show that the fundamental group of every such complex has an automatic structure. A complex of type  $A_1 \times A_1$  is a piecewise Euclidean 2-complex in which each 2-cell is isometric to a square of side 1 in the Euclidean plane. A complex of type  $A_2$  is a 2-complex in which the 2-cells are isometric to an equilateral triangle of side 1; i.e., they are the (3,6)-complexes defined above. The complexes of types  $B_2$  and  $G_2$  are defined similarly, but each 2-cell in a  $B_2$  complex is isometric to an isosceles right triangle with short side of length one, and each 2-cell in a  $G_2$  complex is isometric to a triangle with angles  $\pi/6, \pi/3, \pi/2$  and with short side of length  $1/2$ . The CAT(0) complexes of types  $A_1 \times A_1$  and  $A_2$  are two-dimensional cases of median and bridged complexes defined below.

Now, suppose that  $D$  is a PE disk, i.e., a PE complex, homeomorphic to a 2-disk. For a vertex  $v$  of  $D$ , let  $\alpha(v)$  denote the sum of the corner angles of the 2-cells of  $D$  with vertex  $v$ . If  $v$  is an interior vertex of  $D$ , define the curvature at  $v$  to be  $\kappa(v) = 2\pi - \alpha(v)$ . When  $v$  is a vertex in the boundary  $\partial D$ , define the turning angle at  $v$  to be  $\tau(v) = \pi - \alpha(v)$ . A vertex  $v \in \partial D$  with  $\tau(v) > 0$  is called a *corner* of  $D$ . The following Lyndon's curvature theorem [40] is a PE version of the Gauss–Bonnet theorem:

$$\sum_{v \in D - \partial D} \kappa(v) + \sum_{v \in \partial D} \tau(v) = 2\pi.$$

If a PE disk  $D$  is CAT(0), then  $\kappa(v) \leq 0$  for any interior vertex  $v$ . Consequently,  $D$  has at least two corners.

#### 4. WEAKLY MODULAR GRAPHS

In this section we briefly introduce some classes of graphs and recall their basic properties. All graphs occurring here are connected, and without loops or multiple edges. The *distance*  $d_G(u, v)$  between two vertices  $u$  and  $v$  of a graph  $G = (V, E)$  is the length of a shortest path between  $u$  and  $v$ . The set of all vertices  $w$  on shortest paths between  $u$  and  $v$  is the *interval*  $I(u, v)$  between  $u$  and  $v$ . An induced subgraph  $H$  of a graph  $G$  is *isometric* if the distance between any pair of vertices in  $H$  is the same as that in  $G$ . By  $G(S)$  we denote the subgraph induced by a set  $S \subseteq V$ . An  $n$ -*cycle* is a simple cycle with  $n$  edges.

A graph  $G$  is *weakly modular* [10] if its shortest-path metric  $d_G$  satisfies the following two conditions:

*triangle condition:* for any three vertices  $u, v, w$  with

$$1 = d_G(v, w) < d_G(u, v) = d_G(u, w)$$

there exists a common neighbor  $x$  of  $v$  and  $w$  such that  $d_G(u, x) = d_G(u, v) - 1$ ;

*quadrangle condition:* for any four vertices  $u, v, w, z$  with

$$d_G(v, z) = d_G(w, z) = 1 \quad \text{and} \quad d_G(u, v) = d_G(u, w) = d_G(u, z) - 1,$$

there exists a common neighbor  $x$  of  $v$  and  $w$  such that  $d_G(u, x) = d_G(u, v) - 1$ .

We can define weakly modular graphs using the concept of metric triangle. Recall that three vertices  $u, v$ , and  $w$  form a *metric triangle*  $uvw$  if the intervals  $I(u, v)$ ,  $I(v, w)$ , and  $I(w, u)$  pairwise intersect only in the common end vertices. According to [22],  $G$  is weakly modular if and only if for every metric triangle  $uvw$  all vertices of the interval  $I(v, w)$  are at the same distance  $k = d_G(u, v)$  from  $u$ . The number  $k$  is called the *size* of the metric triangle  $uvw$ . A metric triangle  $uvw$  is a *pseudo-median* of the triplet  $x, y, z$  if the following metric equalities are satisfied:

$$\begin{aligned} d_G(x, y) &= d_G(x, u) + d_G(u, v) + d_G(v, y), \\ d_G(y, z) &= d_G(y, v) + d_G(v, w) + d_G(w, z), \\ d_G(z, x) &= d_G(z, w) + d_G(w, u) + d_G(u, x). \end{aligned}$$

A graph in which every metric triangle is degenerate, that is, has size 0, is called *modular* [9]. In other words, a graph is modular if  $m(x, y, z) = I(x, y) \cap I(y, z) \cap I(z, x)$  is nonempty for every triplet  $x, y, z$ . In this case, each vertex from  $m(x, y, z)$  is called a *median* of  $x, y, z$ . If the median is unique for all triplets, then such a graph is called *median*.

LEMMA 4.1 [9, 42]. *A graph  $G$  is modular if and only if it is triangle-free and satisfies the quadrangle condition. A graph  $G$  is median if and only if it is modular and does not contain  $K_{2,3}$  as a subgraph.*

Median graphs and the related median structures have many nice characterizations and properties, investigated by several authors; [5, 7, 11, 14, 15, 35, 42, 43, 54, 55, 56] is a sample of papers on this subject.

In order to check whether a given subgraph of a weakly modular graph  $G$  is convex or gated the following lemma is useful. It can be proved quite easily by induction.

LEMMA 4.2 [22]. *A connected subgraph  $H$  of a weakly modular graph  $G$  is convex if and only if it is 2-convex, i.e.,  $I(x, y) \subseteq H$  whenever  $x, y \in H$  and  $d_G(x, y) = 2$ . Every convex set of a modular graph  $G$  is gated.*

A graph is called *hereditary weakly modular* if every isometric subgraph is weakly modular. In a similar way we can define *hereditary modular* graphs.

**THEOREM 4.3** [9]. *A graph  $G$  is hereditary modular if and only if all isometric cycles of  $G$  have length four. Equivalently,  $G$  is hereditary modular if and only if it is modular and does not contain induced 6-cycles.*

This result has been extended to hereditary weakly modular graphs (by a *house* is meant the graph obtained by gluing a 3-cycle and a 4-cycle along a common edge):

**THEOREM 4.4** [22]. *A graph  $G$  is hereditary weakly modular if and only if it does not contain induced houses and isometric  $n$ -cycles with  $n \geq 5$ .*

From the characterization of weakly modular graphs via metric triangles we conclude that the balls of weakly modular graphs are isometric subgraphs. Therefore, the balls of hereditary weakly modular graphs induce hereditary weakly modular graphs. An important class of hereditary weakly modular graphs is formed by bridged graphs. A graph is called *bridged* if all isometric cycles of  $G$  have length three; that is, each cycle of length greater than 3 has a shortcut in  $G$ ; see [33, 53].

**THEOREM 4.5** [33, 53]. *A graph  $G$  is bridged if and only if for any convex subset  $S$  of  $G$  and any integer  $k \geq 0$  the  $k$ -neighborhood  $B_k(S)$  is convex.*

A cycle  $C$  of a graph  $G$  is *well bridged* [33] if, for each vertex  $x$  of  $C$ , either the two neighbors  $u, v$  of  $x$  in  $C$  are adjacent, or  $d_G(x, y) < d_C(x, y)$  for some antipode  $y$  of  $x$  in  $C$ . In a bridged graph  $G$  all cycles  $C$  are well bridged [33, 53]. Indeed, if the length of  $C$  is  $2k$ ,  $y$  is the antipode of  $x$  in  $C$ , and  $d_G(x, y) = k$ , then from the convexity of the ball  $B_{k-1}(y)$  we infer that  $u, v$  are adjacent. Similarly, if  $C$  has length  $2k + 1$ ,  $y, z$  are the antipodes of  $x$  in  $C$ , and  $d_G(x, y) = d_G(x, z) = k$ , then from the convexity of  $B_{k-1}(\{y, z\})$  we obtain that  $u$  and  $v$  are adjacent. We continue with a characterization of graphs with convex balls.

**THEOREM 4.6** [33, 53]. *In a graph  $G$  all balls  $B_k(v)$  are convex if and only if every cycle in  $G$  of length other than 5 is well bridged.*

The last class of weakly modular spaces is that of absolute retracts of graphs. A connected graph  $G$  is an *absolute retract* if  $G$  (regarded as a metric space) is a retract of any connected graph in which  $G$  is embedded as an isometric subgraph. A *Helly graph* is a graph in which every family of pairwise intersecting balls has a nonempty intersection. The following result is a discrete analogous of the characterization of absolute retracts due to Aronszajn and Panitchpakdi (for recursive characterizations of Helly graphs see [12]).

**THEOREM 4.7** [36, 46, 50]. *For a graph  $G$ , the following conditions are equivalent:*

- (i)  $G$  is an absolute retract;
- (ii)  $G$  is a Helly graph;
- (iii)  $G$  is a retract of a direct product of paths.

Recall, the direct product of paths is a subspace of an  $l_\infty$ -normed space: it has the integer points of some vector space as vertices and two vertices are adjacent if and only if their  $l_\infty$ -distance is one.

## 5. MINIMAL DISKS

Let  $D$  and  $K$  be two simplicial complexes. A map  $f: V(D) \rightarrow V(K)$  is called *simplicial* if  $f(\sigma) \in K$  for all  $\sigma \in D$ . Any simplicial map  $f: V(D) \rightarrow V(K)$  induces a continuous map  $f: |D| \rightarrow |K|$  by extending  $f$  affinely over the geometric simplexes  $\text{conv}(\sigma)$ ,  $\sigma \in D$ . If  $D$  is a planar triangulation (i.e., the 1-skeleton of  $D$  is an embedded planar graph whose all interior 2-faces are triangles) and  $C = f(\partial D)$ , then  $(D, f)$  is called a *singular disk diagram* (or Van Kampen diagram) for cycle  $C$ ; (for more details see [40, Chap. V]). According to Van Kampen's lemma ([40, pp. 150–151]), for every cycle of a simply connected simplicial complex one can construct a singular disk diagram. A singular disk diagram with no cut vertices (i.e., its 1-skeleton is 2-connected) is called a *disk diagram*. A *minimal (singular) disk* of  $C$  is a (singular) disk diagram  $D$  of  $C$  with a minimum number of 2-faces. This number is called the (*combinatorial*) *area* of  $C$  and is denoted  $\text{Area}(C)$ .

The following result makes precise the structure of minimal disks of simple cycles in 1-skeletons of simply connected simplicial complexes.

**LEMMA 5.1.** (1) *Let  $C = (e_1, \dots, e_m)$  be a simple cycle in the 1-skeleton of a simply connected simplicial complex  $K$ . There exists a disk diagram  $(D, f)$  such that  $f$  bijectively maps  $\partial D$  to  $C$ .*

(2) *If  $(D, f)$  is a minimal disk for a cycle  $C$ , then (a) the image of a 2-simplex of  $D$  under  $f$  is a 2-simplex, and (b) adjacent 2-simplexes of  $D$  have distinct images under  $f$ .*

*Proof.* The first assertion can be easily proved by induction on the number of 2-simplexes involved in a (combinatorial) homotopy of  $C$  to a point. To prove the second assertion, let  $(D, f)$  be a minimal disk of  $C$ . Suppose that  $f(x) = f(y)$  for two adjacent vertices  $x$  and  $y$  of  $D$ . First notice that each 3-cycle  $(x, y, z, x)$  bounds a face of  $D$ . Indeed, otherwise deleting the interior vertices, edges, and simplexes in the region bounded by this cycle, we will obtain a new triangulation  $D'$  with  $\partial D' = \partial D$ . The restriction of  $f$

to  $D'$  is a simplicial map, because it maps the new simplex  $xyz$  to a vertex or to an edge of  $K$ , contrary to the choice of  $(D, f)$ . Therefore  $x$  and  $y$  have only two common neighbors  $p$  and  $q$ , and  $\tau = pxy$  and  $\sigma = qxy$  are simplexes of  $D$ . Consider the simplicial complex  $D'$  obtained by identifying  $zx$  and  $zy$  for  $z \in \{p, q\}$ , by creating a vertex  $xy$ , and by ignoring the simplexes  $\tau$  and  $\sigma$ . One can check that  $D'$  is a disk-triangulation and that the map  $f': V(D') \rightarrow V(K)$  defined by  $f'(u) = f(u)$  if  $u \notin \{x, y\}$  and  $f'(xy) = f(x) = f(y)$  obeys the conditions, contradicting the minimality of  $(D, f)$ .

Suppose that  $f(\sigma) = f(\tau)$  where  $\sigma = xyp$ ,  $\tau = xyq$ . Then  $f(p) = f(q)$  by (a). Consider the triangulation  $D'$  obtained from  $D$  by deleting the edge  $xy$  and the triangles  $\sigma$ ,  $\tau$ , and by adding the edge  $pq$  and the 2-simplexes  $pxq$  and  $pyq$ . The map  $f: V(D') \rightarrow V(K)$  is simplicial, because  $V(D') = V(D)$  and  $f$  maps  $pxq$  and  $pyq$  to the edges  $f(x)f(p)$  and  $f(y)f(q)$  of  $K$ . Therefore,  $(D', f)$  is a minimal disk, contrary to (a). ■

*Remark 5.2.* Let  $(D, f)$  be a minimal disk of  $C$ , and let  $P$  be a path of  $D$  which intersects  $\partial D$  only in the end-vertices. These two vertices split  $\partial D$  into two paths  $P_1, P_2$ , while  $P$  divides  $D$  into two disk-triangulations  $D_1$  and  $D_2$  with  $\partial D_1 = P_1 \cup P$ ,  $\partial D_2 = P_2 \cup P$ . Let  $C_1 = f|_{\partial D_1}$ ,  $C_2 = f|_{\partial D_2}$  (these cycles are not necessarily simple). Then  $(D_1, f)$ ,  $(D_2, f)$  are minimal disks of  $C_1$  and  $C_2$ , respectively.

## 6. MEDIAN COMPLEXES

In this section we show that the cubings (alias CAT(0) cube complexes) coincide with the cubical cell complexes arising from median graphs. We will apply this to derive some results from [51].

### 6.1. Cubings as Median Complexes

Every median graph  $G$  gives rise to an abstract cubical complex  $K(G)$  consisting of all cubes of  $G$ , i.e., subgraphs of  $G$  isomorphic to cubes of any dimensions. The geometric realization  $|K(G)|$  is called a *median complex* (in [27] they are called Buneman complexes). Trivially,  $G$  is recovered from its complex  $|K(G)|$  as the underlying graph. We shall prove the following result.

**THEOREM 6.1.** *Cubings and median complexes are the same.*

*Proof.* Let  $G$  be a median graph. The proof that  $K(G)$  is simply connected is straightforward (see also the proof of (iii) $\Rightarrow$ (i) in Theorem 7.1 below). To prove that  $|K(G)|$  is a cubing we have to show that if three  $(n+2)$ -cubes  $q_1, q_2, q_3$  of  $K(G)$  share a common  $n$ -cube  $q$ , and pairwise

share common  $(n + 1)$ -cubes  $q_{ij}$ , then they are contained in a  $(n + 3)$ -cube of  $K(G)$ . Choose an arbitrary vertex  $x \in q$  and let  $x_{ij}$  denote its neighbor in the facet of  $q_{ij}$  disjoint from  $q$ . Further, denote by  $x_i$  the second common neighbor (it belongs to  $q_i$ ) of the vertices  $x_{ij}$  and  $x_{ik}$ . The median of the triplet  $x_i, x_j, x_k$  is a vertex  $m_x$  outside  $q_1 \cup q_2 \cup q_3$ ; otherwise we get an induced  $K_{2,3}$ , contrary to Lemma 4.1. For the same reason,  $m_x \neq m_y$  if  $x$  and  $y$  are distinct vertices of  $q$ . Moreover, one can easily verify that two vertices  $x$  and  $y$  of  $q$  are adjacent if and only if the vertices  $m_x$  and  $m_y$  are adjacent. Hence the set  $\{m_x : x \in q\}$  induces an  $n$ -cube  $q'$  which together with  $q_1, q_2$ , and  $q_3$  yields the desired  $(n + 3)$ -cube.

To prove the converse, we actually use a weaker flag condition. By a *3-wheel* in a cubical complex we will mean three 2-cubes which share a common vertex, and pairwise share common edges. Let  $K$  be a simply connected abstract cubical complex in which each 3-wheel is contained in a 3-cube of  $K$ . We claim that the 1-skeleton of  $K$  is a median graph.

First, triangulate  $|K^2|$  by adding one diagonal in each 2-cube of  $K$ , thus obtaining a simply connected simplicial 2-complex. Let  $(D, f)$  be a minimal disk triangulation of a cycle  $C$  of  $G(K)$  (we do not require that  $C$  is simple, although from Lemma 5.1 we know that for simple cycles such disks exist). Removing the edges of  $D$  which are mapped to diagonals of 2-cells, we obtain a disk-quadrangulation  $Q$  and a (dimension-preserving) cellular map  $f: Q \rightarrow K^2$  such that  $f|_{\partial Q} = C$  (we also call  $(Q, f)$  a minimal disk of  $C$ ). Applying Lemma 5.1(b) to  $(D, f)$  we deduce that  $f$  maps incident faces of  $Q$  to distinct incident 2-cubes of  $K$ . In particular, every 3-wheel of  $Q$  is mapped to a 3-wheel of  $K$ . Finally, two faces of  $Q$  either are disjoint, or intersect in a single vertex or a single edge (in particular, the graph of  $Q$  does not contain  $K_{2,3}$  as an induced subgraph).

*Claim.* If a cycle  $C$  of  $G(K)$  admits a disk diagram, then there exists a minimal disk  $(Q, f)$  of  $C$  with at least three corners.

We argue by induction on the area of a minimal disk  $(Q, f)$  of  $C$ . Let  $\partial Q = (e_1, e_2, \dots, e_n)$ , where  $e_i = x_i x_{i+1(\text{mod } n)}$ . First suppose that there exists an edge  $e = x_i x_j$  (with  $1 < i < j < n$  for simplicity) which splits  $\partial Q$  into two cycles  $C_1 = (e, e_{j+1}, \dots, e_{i-1})$  and  $C_2 = (e_i, e_{i+1}, \dots, e_j, e)$ . Assume, in addition, that among minimal disks of  $C$  containing such diagonals the disk  $Q$  and the edge  $e$  of  $Q$  are selected to minimize  $n - |i - j|$ . By the induction assumption we can find two minimal disks  $(Q_1, f_1)$  and  $(Q_2, f_2)$  each containing at least three corners and such that  $f_1|_{\partial Q_1} = C_1, f_2|_{\partial Q_2} = C_2$ . Piece them together along the preimages of  $e$  in  $\partial Q_1$  and  $\partial Q_2$ , thus obtaining a minimal disk  $(Q', f')$  for  $C$ . The corners of  $Q_1$  and  $Q_2$  are corners of  $Q'$ , unless they coincide with  $x_i$  or  $x_j$ . If both  $x_i$  and  $x_j$  are corners of  $Q_1$ , then their neighbors  $x_{i-1}$  and  $x_{j+1}$  in  $\partial Q_1$  are adjacent. From the choice of  $e$  we

deduce that  $x_{i-1}x_{j+1} \in \partial Q_1$ . Then  $x_{i-1}, x_{j+1}$  are corners, showing that  $Q'$  has the desired property.

Further we may assume that  $x_i, x_j$  are adjacent in  $Q$  if and only if they are consecutive vertices of  $\partial Q$ . First we show that  $Q$  can be chosen to have at least one corner. Indeed, suppose that  $e_1$  belongs to the face  $q = x_1y_1y_2x_2$  of  $Q$ . If  $y_1$  or  $y_2$  belongs to  $\partial Q$ , then  $x_1$  or  $x_2$  is a corner. Otherwise, replace in  $\partial Q$  the edge  $e_1$  by the path  $(x_1, y_1, y_2, x_2)$ . By the induction assumption, the obtained cycle has a minimal disk  $Q'_0$  with at least three corners. Let  $Q'$  be obtained by gluing  $q$  and  $Q'_0$ . If  $y_1$  or  $y_2$  is a corner of  $Q'_0$ , then we obtain two faces of  $Q'$  which intersect more than in one edge, which is impossible because  $Q'$  is a minimal disk. Thus at least one corner of  $Q'_0$  is a corner of  $Q'$ .

So, let  $Q$  contain at least one corner, say  $x_1$ . Denote by  $q_1 = x_1x_2y_1x_n$  the unique face of  $Q$  containing  $x_1$  and set  $Q'_1 = q_1$ . Let  $\Gamma_1$  be a cycle obtained from  $\partial Q$  by replacing  $x_1$  by  $y_1$ . By the induction assumption there exists a minimal disk  $Q_1$  with  $\partial Q_1 = \Gamma_1$  possessing at least three corners. If  $x_2$  and  $x_n$  are not corners in  $Q_1$ , then gluing  $Q_1$  and  $q_1$  we obtain a minimal disk of  $C$  with the desired property. So, assume without loss of generality that  $x_2$  is a corner of  $Q_1$ . Again, denote by  $q_2 = x_2x_3y_2y_1$  the unique face of  $Q_1$  containing  $x_2$ , set  $Q'_2 = Q'_1 \cup \{q_2\}$ , and let  $\Gamma_2$  be the cycle obtained from  $\Gamma_1$  by replacing  $x_2$  by  $y_2$ . Continuing this way, assume that we have defined the 2-chain  $Q'_i$  being the union of the faces  $q_1, q_2, \dots, q_i$ , and the cycle  $\Gamma_i$  obtained from  $\partial Q$  by replacing the path  $P_i = (x_n, x_1, x_2, \dots, x_{i+1})$  by the path  $P'_i = (x_n, y_1, y_2, \dots, y_i, x_{i+1})$ ; see Fig. 1 for an illustration. By the induction hypothesis,  $\Gamma_i$  has a minimal disk  $Q_i$  with at least three corners. Clearly  $Q' = Q_i \cup Q'_i$  is a minimal disk of  $C$ . The vertex  $y_i$  is not a corner of  $Q_i$ ; otherwise  $Q'$  would contain an induced  $K_{2,3}$ . If  $y_i = x_{i+2}$ , then  $x_{i+1}$  is a corner of  $Q'$  and we can find a required minimal disk applying the transformation described below. So let  $y_i \neq x_{i+2}$ . If  $x_{i+1}$  is a corner of  $Q_i$  and  $x_{i+1} \neq x_n$ , then define new  $y_{i+1}, q_{i+1}, \Gamma_{i+1}, Q'_{i+1}$ , and  $Q_{i+1}$ . Finally, assume that  $x_{i+1}$  is not a corner of  $Q_i$ . Since any corner of  $Q_i$  located on the path  $\partial Q - P'_i$  is a corner of  $Q'$ , it suffices to consider the case when all corners of  $Q_i$  except possibly one belong to  $P'_i$ . Pick a corner  $y_j \in P'_i$ . Let  $q = y_{j-1}y_jy_{j+1}z$  be the unique face of  $Q_i$  containing  $y_j$ . The faces  $q, q_j, q_{j+1}$  form a 3-wheel  $W$  in  $Q'$  which will be mapped to a 3-wheel of  $K$ . By the flag condition, the image of  $W$  is contained in a 3-cube  $Z$  of  $K$ . Perform the following operation with  $Q'$ : remove the faces  $q, q_j, q_{j+1}$  and add a new vertex  $z'$  and new faces  $q' = x_jx_{j+1}x_{j+2}z', q'_j = x_{j+2}z'zy_{j+1}, q'_{j+1} = y_{j-1}x_i z'z$ . We obtain a new minimal disk of  $C$  in which the vertex  $x_{j+1}$  becomes a corner. Indeed, one can map  $q', q'_j, q'_{j+1}$  to the 2-faces of the cube  $Z$  opposite to  $q, q'_j, q'_{j+1}$ , respectively. If  $z \in P'_i$ , say  $z = y_{j-2}$ , then necessarily  $z' = x_{j-1}$ . Replacing the faces  $q, q_{j-1}, q_j, q_{j+1}$  by  $q', q'_j$ , we obtain a disk of  $C$  with smaller combinatorial area, which is




FIG. 1.

impossible. Thus  $z \notin P'_i$ ; in particular,  $P'_i$  does not contain adjacent corners of  $Q_i$ . Therefore we can perform the transformation described above step by step with all corners from  $P'_i$ , yielding a new minimal disk  $Q''$  of  $C$  in which the corners of  $Q_i \cap P'_i$  are replaced by corners of  $Q''$ . Clearly,  $Q''$  obeys the desired property. The same reasoning applies when  $x_{i+1} = x_n$ ; three arising configurations in this case are illustrated in Fig. 1. This concludes the proof of the claim.

From this claim we conclude that the graph  $G(K)$  is triangle-free and its every 4-cycle spans a 2-face of  $K$ . This is so because a simple cycle  $C$  with  $\text{Area}(C) > 1$  must contain in its every disk diagram at least two non-corner vertices. Therefore from Proposition 3.3 we conclude that  $G(K)$  does not contain induced  $K_{2,3}$ . To establish that  $G(K)$  is median, in view of Lemma 4.1 it suffices to show that if  $u, v, w, z$  are distinct vertices and

$v, w \in I(u, z)$  are neighbors of  $z$ , then there exists a common neighbor  $x$  of  $v$  and  $w$  such that  $d_{G(K)}(u, x) = d_{G(K)}(u, v) - 1$ . We proceed by induction on  $k = d_{G(K)}(u, v)$ , departing from the trivial case  $k = 1$ . We may assume that  $I(u, v) \cap I(u, w) = \{u\}$ ; otherwise we replace  $u$  by a closest to  $v$  vertex from this intersection and apply the induction. Let  $P'$  and  $P''$  be the shortest paths connecting the pairs  $u, v$  and  $u, w$ , respectively. Let  $C$  be the (simple) cycle composed of  $P', P''$  and the edges  $vz, zw$ . Suppose that among all shortest paths between  $u, v$  and  $u, w$  the paths  $P', P''$  are selected to form a cycle  $C$  with  $\text{Area}(C) > 1$  minimum. Suppose, additionally, that the quadrangle condition holds for all quadruples of vertices defining a cycle  $C'$  with  $\text{Area}(C') < \text{Area}(C)$ . Choose a minimal disk  $Q$  of  $C$  with at least three corners. For simplicity, we will use the same labels for the vertices of  $Q$  and their images in  $K$ . If at least one corner of  $Q$  is distinct from  $v, w$ , then necessarily we find a corner  $p$  distinct from  $u, v, w, z$ . Let  $p \in P'$ , and by  $a, b$  denote the neighbors of  $p$  in  $P'$ . Consider the face  $pacb$  of  $Q$  containing  $p$ . If we replace in  $P'$  the vertex  $p$  by  $c$ , we obtain a shortest path between  $u$  and  $w$ . The area of a minimal disk of the cycle formed by this path,  $P''$ , and the edges  $wz, zv$  is smaller than  $\text{Area}(C)$ , contrary to the choice of  $P'$ . Thus  $v, w$  are corners. If  $z$  is also a corner, then either  $w$  is adjacent to the neighbor of  $v$  in  $P'$  or  $v$  is adjacent to the neighbor of  $w$  in  $P''$ . In both cases this common neighbor of  $v, w$  is a median of  $u, v, w$ .

Now, assume that  $Q$  has exactly three corners  $u, v, w$ . Let  $y$  be the neighbor of  $v$  in  $P'$  and let  $vyv'z$  be the unique face of  $Q$  containing  $v$ . Clearly  $y \in I(v', u)$ . Consider a cycle  $C'$  formed by the path  $P''$ , the path  $(w, z, v'y)$ , and the part of  $P'$  comprised between  $y$  and  $u$ . Since  $\text{Area}(C') < \text{Area}(C)$  and  $v', w \in I(z, u)$ , by the choice of  $u, v, w, z$ , we can find a vertex  $y'$  adjacent to  $v', w$  at distance  $k - 1$  to  $u$ . Since  $y, y' \in I(v', u)$ , by the induction assumption there is a common neighbor  $u'$  of  $y, y'$  at distance  $k - 2$  to  $u$ . The 2-cells  $zvyv', zv'y'w$ , and  $v'yu'y'$  form a 3-wheel  $W$  of  $K$ . By the flag condition,  $W$  is contained in a 3-cube  $Z$  of  $K$ . Let  $x$  be the unique vertex in  $Z - W$ . Then  $x$  is adjacent to  $u', v$ , and  $w$ ; therefore  $d_{G(K)}(x, u) = k - 1$ . From this we infer that  $x$  is a median of the triplet  $u, v, w$ , whence  $G(K)$  is a median graph. Since all 4-cycles of  $G$  span 2-faces of  $K$ , an easy induction on the dimension together with the flag condition imply that all graphic cubes of  $G(K)$  induce cells of  $K$ . This concludes the proof. ■

In view of Theorem 6.1, the following result is obvious.

**COROLLARY 6.2.**  *$\mathcal{X}$  is a  $\text{CAT}(0)$  complex of type  $A_1 \times A_1$  if and only if its 1-skeleton  $G$  is a cube-free median graph and  $\mathcal{X} = |K(G)|$ .*

From the claim in the proof of Theorem 6.1 we obtain that if we fix in the 1-skeleton of a cubing two shortest paths  $P', P''$  with common end-vertices  $u, v$ , then there exists a sequence of shortest paths  $P_i$  from  $u$  to

$v$  with  $P_1 = P', P_n = P''$ , such that the symmetric difference  $P_i \Delta P_{i+1}$  is a boundary of some 2-cube. This result has been established in [51] and is a rather pleasant way to deform one shortest path to another by means of special elementary homotopies. Notice that the same holds true for shortest paths in the leg graphs of  $F$ -complexes investigated in the next section. Actually, it holds for all graphs satisfying the quadrangle condition.

### 6.2. Properties of Median Complexes

Recently, Sageev [51] and Niblo and Reeves [44] investigated different aspects of groups acting on cubings. For these purposes, the authors of both papers thoroughly explore various geometrical and structural properties of cubings. Theorem 6.1 essentially facilitates this. Moreover, most of such properties have been already known for median graphs; [11, 14, 15, 27, 42, 43, 54, 55] are just a few sources. On the other hand, the CAT(0) property sheds a new light on median complexes. Below, we present a brief account of properties of median graphs and median complexes, attempting a unifying approach in view of Theorem 6.1.

Let  $G$  be the underlying graph of a cubing  $\mathcal{X}$ . For an edge  $uv$  of  $G$  let

$$G(u, v) = \{x \in V(G) : d_G(u, x) < d_G(v, x)\},$$

$$G(v, u) = \{x \in V(G) : d_G(v, x) < d_G(u, x)\}.$$

For brevity, we use this notation also for the subgraphs induced by these sets.

LEMMA 6.3.  $G(u, v), G(v, u)$  are complementary halfspaces of  $G$ .

*Proof.* Since  $G$  is bipartite,  $G(v, u) \cup G(u, v) = V(G)$ . If  $x \in G(u, v)$ , then  $I(x, u) \subseteq G(u, v)$ , whence  $G(u, v)$  is connected and we can apply Lemma 4.2. Pick two vertices  $x, y \in G(u, v)$  at distance two and a vertex  $z$  adjacent to  $x$  and  $y$ . Suppose by way of contradiction that  $z \in G(v, u)$ . Then  $d_G(x, u) = d_G(z, v) = d_G(y, u)$ . Let  $m = m(x, y, u)$ . Since  $m \in I(x, u) \subseteq G(u, v)$ , we conclude that  $m \neq z$ . Since  $m$  and  $z$  are adjacent to  $x$  and  $y$  and are equidistant to  $v$ , we deduce that  $z$  is also a median of the triplet  $x, y, v$ , which is impossible. ■

Following Djoković [25], define for edges  $uv$  and  $u'v'$  of  $G$ ,

$$uv \Theta u'v' \iff u' \in G(u, v) \text{ and } v' \in G(v, u).$$

Lemma 6.3 implies that  $uv \Theta u'v'$  if and only if  $G(u, v) = G(u', v')$  and  $G(v, u) = G(v', u')$ ; i.e., the relation  $\Theta$  is transitive (and is an equivalence relation on  $E(G)$ ). We may compare  $\Theta$  to the following relation  $\Psi$ . For two edges  $e$  and  $e'$  define  $e\Psi e'$  if they either are equal or there exists edges  $e_0, e_1, \dots, e_n$  such that  $e = e_0, e' = e_n$  and  $e_i, e_{i+1} (i = 0, \dots, n - 1)$

constitute opposite edges on some 4-cycle of  $G$ . If  $uv$  and  $u'v'$  are opposite edges of a 2-cube, then  $u' \in G(u, v)$ ,  $v' \in G(v, u)$ ; hence  $uv \Theta u'v'$ . A trivial induction yields  $\Psi \subseteq \Theta$ . To show that  $\Psi$  includes  $\Theta$ , proceed by induction on the distance  $k$  between two distinct edges  $uv$  and  $u'v'$  in the relation  $\Theta$ . Select an arbitrary neighbor  $y$  of  $u'$  on a path joining the pair  $v, v'$ . Let  $x$  be the median of the triplet  $u, u', y$ . Since  $d_G(u, x) = d_G(v, y) < k$ ,  $x$  is distinct from  $v'$ . Convexity of  $G(u, v)$  yields  $x \in G(u, v)$ ; thus  $uv \Theta xy$ . By the induction assumption,  $uv \Psi xy$ . Since  $\Psi$  is an equivalence relation and  $xy \Psi u'v'$ , we infer that  $uv \Psi u'v'$ . This yields  $\Theta \subseteq \Psi$  and in conjunction with  $\Psi \subseteq \Theta$  establishes the following result.

LEMMA 6.4.  $\Theta = \Psi$ .

According to the theorem of Djoković [25] transitivity of  $\Theta$  ensures isometric embeddability into a hypercube (a *hypercube*  $H(\Lambda)$  has all finite subsets of some set  $\Lambda$  as vertices, and cardinality of the symmetric difference  $A\Delta B$  as the distance between  $A, B$ ). Let  $E_i, i \in \Lambda$  be the equivalence classes of  $\Theta = \Psi$  and let  $G'_i, G''_i$  be the complementary halfspaces defined by  $E_i$ . Define the following embedding  $\phi$  of  $G$  into the power set of  $\Lambda$ . Pick a base point  $b$  and set  $\phi(b) = \emptyset$ . For a vertex  $x$ ,  $\phi(x)$  consists of all  $i \in \Lambda$  such that  $b$  and  $x$  lie in different halfspaces defined by  $E_i$ . Clearly, each  $\phi(x)$  is finite. For arbitrary vertices  $x, y$  we have  $d_G(x, y) = \#\phi(x)\Delta\phi(y)$ . Indeed, the convexity of halfspaces implies that if  $x \in G'_i$  and  $y \in G''_i$  then every shortest path connecting  $x$  and  $y$  contains exactly one edge of  $E_i$ . Conversely, if an edge on a shortest path between  $x$  and  $y$  belongs to  $E_i$ , then  $x$  and  $y$  are located in different halfspaces defined by  $E_i$ . Then apply this to shortest paths between  $x, y$  and  $b$  which pass via  $m(x, y, b)$ .

*Remark 6.5.* Proposition 3.3 and the previous result remain valid if one replaces the cubical cells by boxes. Then all edges from one equivalence class  $E_i$  of  $\Theta$  have the same length  $\mu_i$ . We will also call such box complexes cubings.

For a class  $E_i$  of equivalent edges assume that  $b \in G'_i$ . Let  $F'_i$  be the subgraph induced by all vertices of  $G'_i$  which have a neighbor in  $G''_i$ . Analogously define  $F''_i$ . Set  $F_i = F'_i \cup F''_i$ . Let  $H'_i, H''_i$  be the cubical cell complexes induced by  $F'_i, F''_i$ . Obviously,  $H'_i$  and  $H''_i$  are isomorphic. Moreover, given  $0 < \mu < 1$ , one can define a cubical complex  $H_i(\mu)$  isomorphic to  $H'_i$  as follows: on each edge  $x'x'' \in E_i$  pick the point  $x_\mu$  verifying  $d(x_\mu, x') = \mu, d(x_\mu, x'') = 1 - \mu$ . Define a graph  $F_i(\mu)$  with this set of points as vertices: two vertices  $x_\mu \in x'x''$  and  $y_\mu \in y'y''$  are adjacent if and only if  $x'x''$  and  $y'y''$  are opposite edges of some 2-cube of  $G$ . Then  $H_i(\mu)$  is the cubical complex of  $F_i(\mu)$ . Following Sageev [51], we call  $H_i(\mu)$  *geometric hyperplanes*. An important result in [51] is to show that the geometric

hyperplanes of a cubing partition the complex into two connected components. The second important moment is to establish that a hyperplane in a cubing is a cubing and that any finite collection of pairwise intersecting hyperplanes has a point in common. The first assertion is immediate from previous properties, because  $\mathcal{X}$  is the union of  $|G'_i| \cup (|H'_i| \times [0, \mu])$  and  $([\mu, 1] \times |H''_i|) \cup |G''_i|$  glued together along  $H_i(\mu)$ . In particular,  $H_i(\mu)$  separates any pair of vertices from different halfspaces  $G'_i$  and  $G''_i$ .

PROPOSITION 6.6 [51]. *Every geometric hyperplane in a cubing is a cubing.*

*Proof.* First we prove that  $F'_i, F''_i$ , and  $F_i$  are convex subgraphs of  $G$ . From Lemma 6.4 we deduce that  $F'_i$  is connected, therefore it suffices to show 2-convexity of  $F'_i$ . Pick  $x', y' \in F'_i$  at distance 2, their neighbors  $x'', y''$  in  $F''_i$ , and a common neighbor  $z$  of  $x', y'$ . Since  $G''_i$  is convex and  $G$  is bipartite, we infer that  $d_G(x'', y'') = 2$ . Hence the median  $v$  of  $x'', y'', z$  is a vertex of  $G''_i$  adjacent to each vertex from the triplet, whence  $z \in F'_i$ . Thus  $F'_i, F''_i$ , and their union are convex. Since convex subgraphs of a median graph are median, by Theorem 6.1 and the isomorphism of  $H'_i$  and  $H_i(\mu)$  we conclude that  $H_i(\mu)$  is a cubing. ■

As to the Helly property, let  $H_{i_j}(\mu_j)$  ( $j = 1, \dots, m$ ) be a collection of pairwise intersecting geometric hyperplanes. First notice that  $H_i(\mu)$  and  $H_k(\lambda)$  intersect if and only if  $F_i$  and  $F_k$  intersect in  $G$ . Therefore  $F_{i_1}, \dots, F_{i_m}$  is a collection of pairwise intersecting convex sets of  $G$ . Being gated, by the Helly property for gated sets, they share a common vertex  $x'$ . Clearly, if  $x' \in F'_{i_j}$  then its neighbor  $x'' \in F''_{i_j}$  is also a vertex from the intersection. Thus in the intersection we can find a box  $|C|$  sharing a facet  $C_{i_j}$  with every  $F'_{i_j}$ . Take a point  $p \in |C|$  whose distance to  $C_{i_j}$  is  $\mu_j$ . Evidently,  $p \in \bigcap_{j=1}^m H_{i_j}(\mu_j)$ , thus establishing the Helly property.

We continue with a useful property established by some authors asserting that finite median graphs have thin halfspaces.

LEMMA 6.7. *If  $G'_i$  is a minimal proper halfspace of a finite median graph  $G$ , then  $G'_i = F'_i$ ; i.e. any vertex of  $G'_i$  is adjacent to a vertex of  $G''_i$ .*

*Proof.* Suppose not; then we can find a vertex  $v \in G'_i - F'_i$  adjacent to a vertex  $u' \in F'_i$ . Let  $u''$  be the neighbor of  $u'$  in  $F''_i$ . Suppose that the edge  $vu'$  belongs to the equivalence class  $E_j$ . Let  $G'_j$  be the halfspace containing  $v$ . By the choice of  $G'_i$ , there is a vertex  $x \in G'_j \cap G''_i$ . From the convexity of these halfspaces we deduce that the median of  $x, u'', v$  is a common neighbor of  $u'', v$  and belongs to  $G''_i$ . This shows that  $v$  is a vertex of  $F'_i$ , contrary to our assumption. ■

Given a geometric hyperplane  $H_i(\mu)$ , one can refine the cellular structure of  $\mathcal{X}$  by subdividing the cells  $|C|$  intersected by  $H_i(\mu)$  and adding the

intersections  $|C| \cap H_i(\mu)$  as new cells. The underlying graph of the new box complex is obtained from  $G$  by taking on each edge of  $E_i$  the corresponding vertex of  $F_i(\mu)$  and adding the edges of the graph  $F_i(\mu)$ . Since the new cubical complex has  $\mathcal{X}$  as the geometric realization, from Theorem 6.1 (see Remark 6.5) we conclude that the new graph is again median. More generally, we can refine  $\mathcal{X}$  by consequently cutting it with a finite number of geometric hyperplanes, arriving at a new CAT(0) box complex whose 1-skeleton is a median graph.

Every point  $p \in \mathcal{X}$  belongs to the relative interior of a unique cell  $|C|$ . The geometric hyperplanes  $H_{i_1}(\mu_{i_1}), \dots, H_{i_n}(\mu_{i_n})$  passing through  $p$  are all parallel to the facets of  $|C|$ . Define  $\mathbf{p} = (p_1, p_2, \dots)$ , where  $p_i = \mu_{i_j}$  if  $i = i_j$  and  $p_i = 0$  otherwise. If  $p$  is a vertex, then clearly  $\mathbf{p}$  is the characteristic vector of the set  $\phi(p)$ . If we endow each cell with the  $l_1$ -distance and extend it to the intrinsic distance  $\rho_1$  on  $\mathcal{X}$ , then one can show that for any points  $x, y \in \mathcal{X}$  we have

$$\rho_1(x, y) = \|\mathbf{x} - \mathbf{y}\|_1 = \sum_{i \in \Lambda} |x_i - y_i|.$$

(The sum is well-defined, because each of the vectors  $\mathbf{x}, \mathbf{y}$  has only a finite number of nonzero coordinates.)  $(\mathcal{X}, \rho_1)$  is a median space, i.e., for any three points  $x, y, z \in \mathcal{X}$  there exists a unique point  $m = m(x, y, z)$  such that  $m$  belongs to all intervals between  $x, y, z$ ; for details about this and previous result see [55]. In view of previous results, one can sketch the following proof of this fact: refine  $\mathcal{X}$  by cutting it with all geometric hyperplanes passing through at least one of the points  $x, y, z$ . We obtain a cubing whose 1-skeleton is a (weighted) median graph isometrically embedded in  $(\mathcal{X}, \rho)$ . Since  $x, y, z$  are vertices of this graph, their median will be the median in  $(\mathcal{X}, \rho)$  (its unicity easily follows).

**THEOREM 6.8** [55]. *If  $\mathcal{X}$  is a cubing, then  $(\mathcal{X}, \rho_1)$  is a median space, which embeds isometrically into a space of  $L_1$ -type.*

Median spaces are particular instances of a remarkable algebraic structure. A *median operator* on a set  $X$  is a function  $m: X^3 \rightarrow X$  satisfying the following conditions:

(absorption law)  $m(a, a, b) = a$ ;

(symmetry law) if  $\sigma$  is any permutation of  $a, b, c$ , then

$$m(\sigma(a), \sigma(b), \sigma(c)) = m(a, b, c);$$

(transitive law)  $m(m(a, b, c), d, c) = m(a, m(b, c, d), c)$ .

The resulting pair  $(X, m)$  is called a *median algebra*. Median algebras and related convexity and metric median structures have a rich theory. Their

study goes back to Birkhoff, Kiss and Sholander; for an extensive survey, see [11, 55]. A more general notion of median algebra is given in [35]. We only mention that there is a bijection between finite (discrete) median algebras and median graphs; with a discrete median algebra  $(X, m)$  one can associate a median graph by taking  $X$  as the vertex-set and all pairs  $xy$  such that  $m(x, y, z) = x$  or  $y$  for all  $z \in X$ , as edges.

Mai and Tang [41] established that any finite collapsible simplicial complex  $X$  is injectively metrizable. To prove this, they showed that  $X$  can be subdivided to a collapsible cubical complex  $K$ , which is an injective space with respect to the intrinsic  $l_\infty$ -metric. Recall, this metric  $\rho_\infty$  is defined in the following way: given two points  $x, y \in |K|$  so that if  $x$  and  $y$  are in a common cell, say in a  $k$ -cell, then  $\rho_\infty(x, y) = \max_i |x_i - y_i|$ , where  $x = (x_1, \dots, x_k), y = (y_1, \dots, y_k)$ ; otherwise the distance is the length of a shortest path joining them. Recently, van de Vel [56] observed that the cubical complexes occurring in [41] are actually the median complexes, leading to the following result.

**THEOREM 6.9** [41, 55]. *If  $\mathcal{X}$  is a locally finite cubing, then  $(\mathcal{X}, \rho_\infty)$  is an injective space.*

A discrete version of the result of Mai and Tang is given by Bandelt and van de Vel [15]. It turns out that there is a parallel between their result and the approach of Niblo and Reeves [44] to bicombing the fundamental group of a cubing. Given the median graph  $G$  of a cubing  $\mathcal{X}$ , let  $G^\Delta$  be the graph having the same vertex set as  $G$ , where two vertices are adjacent if and only if they belong to a common cube of  $G$ . Halfspaces  $G_{i_1}, G_{i_2}, \dots, G_{i_q}$  of  $G$  form a chain if

$$G_{i_1} \subset G_{i_2} \subset \dots \subset G_{i_q}.$$

Equivalently, for any  $0 < \mu < 1$ , the geometric hyperplanes  $H_{i_1}(\mu), \dots, H_{i_q}(\mu)$  are pairwise disjoint.

**PROPOSITION 6.10** [15]. *If  $G$  is a median graph, then  $G^\Delta$  is a Helly graph. The distance  $d_{G^\Delta}(x, y)$  is the largest possible length of a chain of nontrivial halfspaces separating  $x$  and  $y$ .*

Each path  $P_{(v_0, v_n)} = (v_0, v_1, \dots, v_{n-1}, v_n)$  of  $G^\Delta$  corresponds to a chain of cubes [15]: this is a sequence of cubes  $C_0, C_1, \dots, C_n$ , such that  $C_i$  is the cube of minimal dimension containing  $v_i$  and  $v_{i+1}$ . Call a chain of cubes satisfying  $C_i \cap C_{i+1} = \{v_{i+1}\}$  for all  $i < n$  a *cube-path*. A cube-path is called a *normal cube-path* [44] if  $C_{i+1}$  intersects  $\text{star}(C_i, \mathcal{X})$  in  $v_{i+1}$  (recall, that  $\text{star}(C_i, \mathcal{X})$  consists of all cubes containing  $C_i$ ). The following result is a nice observation of Niblo and Reeves.

PROPOSITION 6.11 [44]. *Given two vertices  $v, w$  of a median graph  $G$ , there is a unique normal cube-path from  $v$  to  $w$  and a unique normal cube-path from  $w$  to  $v$ . Both paths are shortest paths in the graph  $G^\Delta$ .*

Using Proposition 6.11 and Lemma 4.2 one can readily check that  $G^\Delta$  is a Helly graph. Indeed, as in the proof of Mai and Tang or that of Bandelt and van de Vel, it suffices to show that every ball  $B_r(b)$  of  $G^\Delta$  is a convex set of  $G$  (recall that the convex sets of a median graph are gated). By Lemma 4.2, it is enough to show that if  $x, y \in B_r(b)$ ,  $d_G(x, y) = 2$ , then any of their common neighbor  $z$  in  $G$  belongs to  $B_r(b)$ . Consider the last cubes  $C'$  and  $C''$  in the normal cube-paths  $P_{(b,x)}$  and  $P_{(b,y)}$  connecting  $b$  with  $x$  and  $y$ , respectively. The star of at least one cube, say of  $C'$ , must contain the vertex  $z$ ; otherwise, adding to  $P_{(b,x)}$  and  $P_{(b,y)}$  the edges  $xz$  and  $yz$ , respectively, we get two distinct normal-cube paths between  $b$  and  $z$ . Therefore  $d_{G^\Delta}(b, z) \leq d_{G^\Delta}(b, x)$ , whence  $z \in B_r(b)$ .

In some sense, Proposition 6.11 expresses the fact that given a base-point  $b$  of a median graph  $G$ , the pair  $(V(G), \leq_b)$  is a median semilattice; e.g. [55], each order interval  $[b, u]$  is distributive, and a nonempty finite set of vertices has an upper bound provided that each pair of its vertices has an upper bound. To construct the normal cube-path  $P_{(v,w)}$  from  $v$  to  $w$ , consider the set  $F$  of all neighbors of  $v$  in the interval  $I(v, w)$ . Take the upper bound  $u$  of  $F$ . Then  $I(v, u)$  induces a cube  $C$ . Define  $P_{(v,w)}$  as the union of  $C$  and  $P_{(u,w)}$ . From the choice of  $u$  we conclude that any cube containing  $C$  intersects the first cube of the path  $P_{(u,w)}$  in  $u$ , and by the induction assumption  $P_{(v,w)}$  is indeed a normal cube-path.

*Remark 6.12.* In fact, it is possible to deduce Theorem 6.9 from Proposition 6.10 and Proposition 6.11 from Lemma 6.7, but we will not present them here.

We conclude the list of properties of median structures with the following important characterization of median graphs.

THEOREM 6.13 [7]. *Median graphs are precisely the retracts of hypercubes.*

Bandelt and van de Vel [13] have proven that every edge-preserving map  $f: G \rightarrow G$  of a finite median graph has an invariant cube, i.e., a cube of  $G$  which is mapped isomorphically onto itself by  $f$ . On the other hand, Kirk [39], generalizing a result of Penot [48], proved that every nonexpansive map of a convex bounded weakly countably compact metric space in which all balls are convex has a fixed point. Van de Vel [55, p. 504] noticed that “the similarities between ...” two results “... are too detailed to be pure coincidence.” It seems that the CAT(0) property of median complexes gives a partial explanation to this fact. Namely, assume additionally that  $f: G \rightarrow G$  is a cell-to-cell map. Then  $f$  induces a continuous map  $f: |K(G)| \rightarrow$

$|K(G)|$  by extending  $f$  affinely over the geometric cubes. As a result, we obtain a nonexpansive map of a CAT(0) space  $|K(G)|$ . From Theorem 3.2(iii) and the result of Kirk,  $f$  has a fixed point  $p$ . Consider the smallest cube  $C$  of  $G$  such that  $p$  belongs to the relative interior of  $|C|$ . Clearly,  $C$  is an invariant cube of the initial map  $f$ . This remark can be extended to obtain fixed simplexes of edge-preserving maps of underlying graphs of CAT(0) simplicial complexes, but does not extend to arbitrary edge-preserving maps of CAT(0) polysimplicial complexes (cell complexes whose cells are products of simplexes).

## 7. SIMPLICIAL COMPLEXES WITH RIGHT TRIANGULAR CELLS

In this Section we assume that  $\mathcal{X}$  is a PE 2-complex with only finitely many isometry types of cells whose 2-cells are right triangles and the attaching map verifies the following conditions:

- (a) if the 2-cells  $\sigma', \sigma''$  share a common edge  $e$ , then  $e$  is either the hypotenuse of both  $\sigma', \sigma''$  or a leg of both  $\sigma', \sigma''$ ;
- (b) if the 2-cells  $\sigma', \sigma''$  are glued along the hypotenuse, then  $\sigma' \cup \sigma''$  is isometric to a Euclidean rectangle;
- (c) each hypotenuse belongs to at least two triangles.

Simple arguments show that in a cell complex obeying (a) and (b) all maximal cells are triangles; thus the assumption that  $\mathcal{X}$  is two-dimensional does not restrict the generality. Further, a 2-complex satisfying (a) and (b) can be complemented by adding “pendant triangles” to fulfill (c) as well. Every  $B_2$  complex (for a definition see Section 3) satisfies (a) and (b). A particular class of  $B_2$  complexes arises from orientable hereditary modular graphs (alias frames), and has been introduced and investigated by Karzanov [37]. His construction actually allows us to derive a  $B_2$  complex from an arbitrary graph.

Let  $K = K(\mathcal{X})$  denote the abstract simplicial complex associated with  $\mathcal{X}$ . A *leg graph*  $LG(K)$  is obtained by deleting the hypotenuse-edges from the 1-skeleton  $G(K)$  of  $K$ . Clearly, if a hypotenuse  $h$  is shared by  $n$  triangles, then the subgraph of  $LG(K)$  induced by  $\text{star}(h, K)$  is the complete bipartite graph  $K_{2,n}$ . Following [37], we call the union of all triangles of  $\mathcal{X}$  sharing the hypotenuse  $h$  a *folder* and denote it by  $F_h$ . Conversely, an arbitrary graph  $G$  gives rise to a cell complex  $\mathcal{X}(G)$  which fulfills the requirements (a), (b), and (c). For this, in each maximal complete bipartite subgraph  $K_{2,n}$  ( $n \geq 2$ ) of  $G$  (further called a *biclique* of  $G$ ) pick two vertices  $x, y$  which are adjacent to all other vertices  $z \in K_{2,n}$  ( $x, y$  are uniquely defined unless  $n = 2$ ). Define the right triangle  $xyz$  with the right angle at  $z$ , and




FIG. 2.

set  $\mathcal{X}(G)$  be the union of all such triangles. The obtained complex  $\mathcal{X}(G)$  verifies (a) and (c). If we require that the opposite edges in every 4-cycle of  $G$  have equal lengths, then  $\mathcal{X}(G)$  obeys the condition (b) as well. Trivially,  $G$  is recovered from its complex  $\mathcal{X}(G)$  as the leg graph. In general, the structure of  $\mathcal{X}(G)$  can be quite complicated: consider, for example, the complexes generated by the graphs  $K_{3,3}$  and  $K_{3,3}^-$  from Fig. 2.

7.1. Folder-Complexes

We say that a complex  $\mathcal{X}$  obeying (a) and (b) is a *folder complex* (*F-complex*) if the intersection of any two folders does not contain incident legs and  $\mathcal{X}$  does not contain any three folders  $F_{h_1}, F_{h_2}, F_{h_3}$  and three distinct legs  $e_1, e_2, e_3$  sharing a common vertex such that  $e_i$  belongs to  $F_{h_j}$  exactly when  $i \neq j$ . (This is a “folder” version of the flag condition for cubical complexes.)

**THEOREM 7.1.** *For a simply connected cell 2-complex  $\mathcal{X}$  satisfying the conditions (a), (b) and (c) the following conditions are equivalent:*

- (i)  $\mathcal{X}$  satisfies  $CAT(0)$ ;
- (ii)  $\mathcal{X}$  is an *F-complex*;
- (iii)  $\mathcal{X} = \mathcal{X}(G)$  for a hereditary modular graph  $G$  without  $K_{3,3}$  and  $K_{3,3}^-$  as induced subgraphs.

If  $\mathcal{X}$  is simply connected and satisfies (a) and (b) then (i)  $\Leftrightarrow$  (ii)  $\Rightarrow$  (iii).

*Proof.* (i)  $\Rightarrow$  (ii): Suppose by way of contradiction that  $\mathcal{X}$  contains three folders  $F_{h_1}, F_{h_2}, F_{h_3}$  and three distinct legs  $xy_1, xy_2, xy_3$  such that  $xy_i$  belongs to  $F_{h_j}$  exactly when  $i \neq j$ . The hypotenuse  $h_j$  is either  $y_i y_k$  ( $i, k \neq j$ ) or  $xx_j$ . In the first case,  $y_i y_k$  belongs to the PS complex  $\text{Link}(x, \mathcal{X})$  and its spherical length is  $\pi/2$ . In the second case,  $y_i x_j$  and  $y_k x_j$  belong to

$\text{Link}(x, \mathcal{X})$ , and by condition (b) the spherical length of the path  $y_i x_j y_k$  again is  $\pi/2$ . Concatenating such hypotenuses and paths, we obtain a simple cycle of  $\text{Link}(x, \mathcal{X})$  of length  $2\pi/3$ , contrary to Theorem 3.2.

Now, suppose that the folders  $F_{h'}$  and  $F_{h''}$  share two incident legs  $xy$  and  $xz$ . At least one of  $h'$  or  $h''$ , say  $h'$ , is distinct from  $yz$ . Then  $h' = xu$ , the union of the 2-cells  $xuy$  and  $xuz$  being a rectangle. If  $h'' = yz$ , then  $\text{Link}(x, \mathcal{X})$  contains a simple cycle  $(z, y, u, z)$  of length  $\pi$ . Otherwise, if  $h'' = xv$ , the simple cycle  $(z, v, y, u, z)$  of  $\text{Link}(x, \mathcal{X})$  has spherical length  $\pi$ , in contradiction to Theorem 3.2.

(ii)  $\Rightarrow$  (iii): Let  $C$  be a simple cycle in the graph  $LG(\mathcal{X})$ . Consider a minimal disk  $(D, f)$  of  $C$ . Delete from  $D$  all edges whose images under  $f$  are hypotenuses of  $\mathcal{X}$ . Since the map  $f$  is dimension-preserving, the resulting planar graph  $Q$  is a quadrangulation; i.e., all (interior) faces of  $Q$  are quadrangles.

*Claim.* The degree of every interior vertex  $x$  of  $Q$  is at least 4.

Suppose not. First, assume that  $x$  has exactly two neighbors  $y$  and  $z$ , and let  $yxzu$  and  $yxzv$  be the faces of  $Q$  incident to  $x$ . Necessarily,  $x$  is adjacent in  $D$  to at least one of the vertices  $u, v$ . We may thus assume that  $xu$  is an edge of  $D$ . If  $x$  is adjacent to  $v$  too, by Lemma 5.1 the triangles  $xuy, xuz, xvy, xvz$  will be mapped to four distinct triangles from the folders  $F_{f(xu)}$  and  $F_{f(xv)}$ . Since the legs  $f(xy), f(xz)$  are incident and belong to both folders, we obtain a contradiction with the assumption that  $\mathcal{X}$  is a folder complex. The same contradiction arises in the case when  $y$  and  $z$  are adjacent in  $D$  and  $f(v) \neq f(u)$ . So, let  $f(v) = f(u)$ . Then  $F_{f(yz)}$  and  $F_{f(xu)}$  share the 4-cycle  $(f(y), f(x), f(z), f(u) = f(v), f(y))$ , a contradiction.

Now, suppose that  $x$  is adjacent in  $Q$  to three vertices  $y, z, w$ . Let  $q_1 = x_1 y x z, q_2 = x_2 z x w, q_3 = x_3 w x y$  be the faces of  $Q$  incident to  $x$ , each of them being a union of two triangles of  $D$ ; see Fig. 3 in which the edges mapped to hypotenuses are heavy. Let  $e_i$  be the common edge of two triangles constituting  $q_i$ . If  $e_1, e_2, e_3$  have distinct images in  $K$ , we obtain three folders  $F_{f(e_1)}, F_{f(e_2)}, F_{f(e_3)}$  and three edges  $f(xy), f(xz), f(xw)$ , which contradicts the hypothesis that  $\mathcal{X}$  is a folder complex. From this and Lemma 5.1 we immediately conclude that  $x$  cannot be adjacent to all three vertices  $x_1, x_2, x_3$ , and that the vertices  $y, z, w$  are not pairwise adjacent, thus settling the first two cases from Fig. 3. An easy analysis of the remaining cases show that every simplicial map  $f$  with  $f(e_i) = f(e_j)$  violates one of the conditions of Lemma 5.1. This proves the claim.

From this claim and the Gauss–Bonnet formula one can easily show that  $\partial Q$  contains at least four corners (in this case they are vertices of degree two of  $Q$ ). Indeed,  $Q$  is a CAT(0) (4,4)-complex; thus  $\kappa(v) \leq 0$  for all interior vertices  $v$ . Since the turning angles are nonpositive or  $\pi/2$ , necessarily  $Q$  contains at least four corners. From this we immediately obtain that  $LG(K)$




FIG. 3.

is triangle-free. Now, suppose that  $C$  is a 4-cycle of  $LG(K)$ . Then  $Q$  consists of a single quadrangle; i.e.,  $C$  is the boundary of a union of two triangles of  $\mathcal{X}$  sharing a common hypotenuse. Since  $\mathcal{X}$  is simply connected, from this we deduce that  $LG(K)$  is bipartite. As  $\mathcal{X}$  is a folder complex, a simple analysis in conjunction with this implies that  $LG(K)$  does not contain  $K_{3,3}$  and  $K_{3,3}^-$  as induced subgraphs. To prove that  $LG(K)$  is hereditary modular, in view of Theorem 4.3 we must show that  $LG(K)$  does not contain isometric cycles of length at least 6. Let  $C$  be a simple cycle with at least six edges, and let  $(Q, f)$  be a minimal disk quadrangulation of  $C$ . From Theorems 6.1 and 4.3 we have that  $Q$  is a cube-free median graph, whence  $Q$  is a hereditary modular graph. Hence  $\partial Q$  is not an isometric cycle of  $Q$ . Since  $f: Q \rightarrow LG(K)$  is nonexpansive, the image of  $\partial Q$  is a non-isometric cycle of  $LG(K)$ , yielding that  $LG(K)$  is hereditary modular.

(iii)  $\Rightarrow$  (i): Let  $G$  be a hereditary modular graph which does not contain  $K_{3,3}$  and  $K_{3,3}^-$  as induced subgraphs and denote  $\mathcal{X} = \mathcal{X}(G)$ . Each 4-cycle is the boundary of two triangles of  $\mathcal{X}$ . To show that  $\mathcal{X}$  is simply connected, it suffices to prove that every simple cycle  $C$  of  $G$  can be (combinatorially) deformed to an arbitrary vertex  $x \in C$ . We apply the induction on  $\sum_{w \in C} d_G(x, w)$ . Walking around  $C$  starting from  $x$ , we arrive at a vertex  $y \in C$  such that its neighbors  $u, v$  in  $C$  obey  $d_G(x, u) = d_G(x, v) < d_G(x, y)$ . By modularity condition we can find a common neighbor  $y'$  of  $u$  and  $v$  one step closer to  $x$ . Let  $C'$  be the cycle obtained from  $C$  by replacing  $y$  by  $y'$  (and eventually removing one of the vertices  $u$  or  $v$  if  $y' \in C$ ). Clearly, for  $C'$  we can apply the induction hypothesis; thus  $C'$  can be deformed to  $x$ . Since  $C$  can be transferred to  $C'$  using two elementary deformations in  $\mathcal{X}$ , we are done.

To establish that  $\mathcal{X}$  is nonpositively curved, by Theorem 3.2 we must show that for a vertex  $x$  the length of every simple cycle in the PS complex  $\text{Link}(x, \mathcal{X})$  is at least  $2\pi$ . To prove this, first we present few simple properties of  $\mathcal{X}$ . One can readily check that the maximal complete bipartite subgraphs of  $G$  are bicliques. Therefore every 4-cycle of  $G$  belongs to a unique folder of  $\mathcal{X}$ . Two folders either are disjoint or intersect in a vertex or in an edge; otherwise we get a forbidden subgraph  $K_{3,3}$  or  $K_{3,3}^-$ . This implies that  $\text{Link}(x, \mathcal{X})$  does not contain 4-cycles of  $G$ . Indeed, if we could

find such a cycle  $(y, z, u, v, y)$ , then  $x$  must be adjacent in  $G$  to some opposite vertices of this cycle, say to  $y$  and  $u$ . Hence  $xz$  and  $xv$  are hypotenuses, yielding two folders  $F_{xz}$  and  $F_{xv}$  with a forbidden intersection.

Consider a simple cycle  $C$  in  $\text{Link}(x, \mathcal{X})$ . The legs of  $C$  combine in pairs  $l', l''$  to form paths whose spherical length is  $\pi/2$ . Indeed, pick a leg  $l' = yz$ , and suppose that  $xy$  is the hypotenuse of the triangle  $xyz$ . If  $u$  is the neighbor of  $y$  in  $C$ , then  $xuy$  is a 2-cell, and therefore we can set  $l'' = yu$ . On the other hand, any hypotenuse  $h = uv$  of  $C$  has length  $\pi/2$ . In addition, we can find a vertex  $x_h \notin C$  such that  $(x, u, x_h, v, x)$  is a 4-cycle of  $G$ . If we take another hypotenuse  $h'$  of  $C$ , then the vertices  $x_h$  and  $x_{h'}$  are distinct. Moreover,  $x_h$  cannot be adjacent in  $G$  to any end-vertex of  $h'$ ; otherwise  $G$  would contain a 3-cycle.

Now, if  $C$  contains at least four hypotenuses and/or  $\pi/2$ -pairs of legs, then the length of  $C$  is at least  $2\pi$  and we are done. The subsequent case analysis will show that the converse cannot happen. First suppose that  $C$  has only three edges, say  $C = (y, z, u, y)$ . Since  $G$  is triangle-free, at least one edge of  $C$ , say  $h = yu$ , is a hypotenuse. If  $yz$  and  $zu$  are legs, then  $xz$  or  $xu$ , say the first one, is a hypotenuse, providing two folders  $F_h$  and  $F_{xz}$  with a forbidden intersection. Otherwise, if  $h' = yz$  and  $h'' = zu$  are hypotenuses, then  $(u, x_h, y, x_{h'}, z, x_{h''}, u)$  is an induced 6-cycle of  $G$ , contrary to the fact that  $G$  is hereditary modular. A similar contradiction arises when  $C$  has six edges and is a cycle of  $G$ . Then  $C$  is not induced, and this would imply that the link of  $x$  contains a 4-cycle of  $G$ , which we have proven to be impossible. If  $C$  has two hypotenuses  $h' = yz$ ,  $h'' = yv$  and two legs  $zu, uv$ , then in  $G$  we obtain an induced 6-cycle  $(y, x_{h'}, z, u, v, x_{h''}, y)$ . Indeed, as we have shown before,  $x_{h'}v, x_{h''}z \notin E(G)$ . On the other hand,  $uy \in E(G)$  would imply that  $u \in F_{h'} \cap F_{h''}$ , which is impossible because these folders intersect already along the edge  $xy$ . Finally, assume  $C$  consists of a hypotenuse  $yz$  and two pairs of legs  $zu, uv$  and  $vw, wy$ . Consider the 6-cycle  $(y, x_h, z, u, v, w, y)$  of  $G$ , which cannot be induced. However, if  $G$  contains any one of the edges  $x_hv, yu$ , or  $zw$ , then any two of the folders  $F_{yz}, F_{xu}, F_{xw}$  intersect more than in one edge. This proves that every simple cycle  $C$  in the link of  $x$  has length at least  $2\pi$ , concluding the proof that  $\mathcal{X}$  is a CAT(0) space. ■

Motivated by Theorem 7.1, we call a hereditary modular graph which does not contain  $K_{3,3}$  and  $K_{3,3}^-$  as induced subgraphs an *F-graph*. These graphs lie in “between” frames (orientable hereditary modular graphs) and semiframes (hereditary modular graphs without induced  $K_{3,3}^-$ ) investigated in [37, 38], respectively.

As we noticed already, a  $B_2$  complex verifies the conditions (a) and (b); therefore, the CAT(0) complexes of this type can be characterized via

Theorem 7.1. Namely, given an F-graph  $G$ , one can define a unique  $B_2$  complex  $\mathcal{X}(G)$  by setting all edges of  $G$  to have length one.

To present the next example, let  $\mathcal{K}$  be a simply connected (4,4)-complex. Let  $\mathcal{X}$  be the subdivision of  $\mathcal{K}$  obtained by adding a new vertex at the barycenter of each 2-face, and then joining it in the face with the vertices on the boundary. Then  $\mathcal{X}$  is a simplicial complex. Model each simplex of  $\mathcal{X}$  as an isosceles triangle with angles  $\pi/2, \pi/4, \pi/4$ , the edge of  $\mathcal{K}$  of length 1, and two new edges of length  $1/\sqrt{2}$ . Clearly,  $\mathcal{X}$  is a  $B_2$  complex, and  $G(\mathcal{X})$  coincides with the graph  $HG(\mathcal{X}) = G(\mathcal{X}) - LG(\mathcal{X})$ . The converse is not true. One can construct examples of F-complexes  $\mathcal{X}$  in which the graph  $HG(\mathcal{X})$  has a prescribed number of connected components, whose interplay can be quite complicated.

## 7.2. Properties of F-Complexes and F-Graphs

Since every two folders intersect in an edge or a vertex, with some abuse of terminology, we can assume that  $\mathcal{X}$  is a cell complex, whose 2-cells are the folders and the 1-cells are the legs of triangles. Let  $G$  be an F-graph endowed with the standard graph-metric  $d_G$ . Due to the bijection between folders and bicliques, in some places below, by a folder we actually mean the biclique of  $G$  induced by a folder of  $\mathcal{X}(G)$ . The first two assertion can be easily derived from the definitions and Lemma 4.2; Lemmas 7.4 and 7.6 have been proven in [37] for frames; however, we give their direct proofs.

LEMMA 7.2. *Every folder is a gated subgraph of  $G$ .*

LEMMA 7.3. *Each ball  $B_k(x)$  induces an isometric subgraph of  $G$ .*

LEMMA 7.4. *If  $d_G(b, v) = k + 1$ , then either  $v$  has a unique neighbor  $v^b$  in  $B_k(b)$  or there is a unique vertex  $v^b$  at distance  $k - 1$  to  $b$  which is adjacent to all neighbors of  $v$  in  $B_k(b)$ .*

*Proof.* Pick two neighbors  $x, y \in B_k(b)$  of  $v$ , and let  $v' \in m(x, y, b)$ . Assume that  $v''$  is another median of this triplet. Both  $v'$  and  $v''$  are common neighbors of  $x, y$  at distance  $k - 1$  to  $b$ . Let  $z \in m(v', v'', b)$ . Then the vertices  $v, x, y, v', v'', z$  induce the forbidden  $K_{3,3}^-$ . Hence the median  $v'$  is unique. Let  $z \in B_k(b)$  be another neighbor of  $v$ . We assert that  $z$  is adjacent to  $v'$ . Indeed, suppose not, and let  $z', z''$  be the medians of the triplets  $x, z, b$  and  $y, z, b$ , respectively. Then we have constructed an induced 6-cycle  $(x, v', y, z'', z, z', x)$ , a contradiction. ■

This property implies that in  $B_{k+1}(b)$  the vertex  $v$  belongs either to a unique edge  $vv^b$  or to a unique folder  $F$ . From Lemma 7.2 we infer that  $v^b$  is the gate of  $b$  in  $F$ . Therefore, a total order  $\prec_b$  refining  $\leq_b$  has the property that in the subgraph induced by  $\{z : z \prec_b v\}$  the vertex  $v$  belongs

to a unique cell  $F_{vv^b}$  spanned by the pair  $vv^b$  (we will call this cell *pendant*, specifying its type, if necessary).

A *folder-path* between two vertices  $v', v$ , is a sequence of folders and/or edges  $F_0, \dots, F_n$ , such that  $v' \in F_1, v \in F_n$  and  $F_i \cap F_{i+1} = \{v_i\}$  for  $i = 0, \dots, n - 1$ . By induction on  $n = d_G(b, v)$  we define a special folder-path  $P_{(v,b)}$  between  $b$  and each vertex  $v$ . Namely, take the pendant cell  $F_{vv^b}$  spanned by  $v$  and  $v^b$  and set  $P_{(v,b)} = \{F_{vv^b}\} \cup P_{(v^b,b)}$ . One can readily check that if  $F_0, F_1, \dots, F_n$  are the cells of  $P_{(v,b)}$ , then any maximal cell containing  $F_i$  intersects  $F_{i+1}$  in exactly one vertex for all  $i < n$ . Due to the similarity with normal cube-paths, we call the folder-paths obeying this condition a *normal folder-path*. As in the case of cube-paths, every folder-path can be identified with a sequence  $(v_0, v_1, \dots, v_n)$  of vertices, such that each pair of consecutive vertices spans an edge or a folder. We can extend such a path to a (geometric) path in the complex  $\mathcal{X}(G)$ . If we take the union of all paths  $P_{(v,b)}$  pointing at  $b$ , we will obtain a tree  $T_b$  (a dendron in  $\mathcal{X}(G)$ ). One can construct examples of F-complexes in which certain normal folder-paths are not paths of the underlying graph. However, as in the case of cubings, folder-paths correspond to paths in an appropriate graph  $G^\Delta$  associated with an F-graph  $G$ . Namely, let  $G^\Delta$  be the graph having the same vertex set as  $G$ , where two vertices  $v, w$  are adjacent iff they belong to a common folder or edge of  $\mathcal{X}$ .

PROPOSITION 7.5. (1)  $P_{(v,b)}$  is the unique normal folder-path between  $v$  and  $b$ . All normal folder-paths are shortest paths in  $G^\Delta$ .

(2)  $G^\Delta$  is a Helly graph.

*Proof.* (1) By induction on  $d_{G^\Delta}(b, v)$  we prove that  $P_{(v,b)}$  is a shortest path in  $G^\Delta$ . In  $G^\Delta$  pick a neighbor  $x$  on a shortest path between  $v$  and  $b$ . By the induction assumption,  $P_{(x,b)}$  is a shortest path. If  $d_G(x, b) > d_G(v, b)$ , Lemma 7.4 would imply that  $x, v$ , and  $x^b$  lie in a common folder; thus  $v$  and  $x^b$  are adjacent in  $G^\Delta$ , contrary to the choice of  $x$ . Hence  $d_G(x, b) \leq d_G(v, b)$ . If  $x = v^b$ , we are done. If  $x$  and  $v$  are adjacent in  $G$ , then  $v^b$  is adjacent to both  $x$  and  $x^b$ . Consequently,  $v^b$  is adjacent to both  $v$  and  $x^b$  in  $G^\Delta$ , whence  $d_{G^\Delta}(v^b, b) < d_{G^\Delta}(v, b)$ . Finally, assume that  $d_G(x, b) = d_G(v, b)$ . The median  $m$  of the triplet  $v, x, b$  is adjacent to  $v^b$  and  $x^b$ ; therefore  $m, v^b, x^b$  belong to a common folder. We obtain  $d_{G^\Delta}(v^b, b) < d_{G^\Delta}(v, b)$ , again establishing that  $P_{(v,b)}$  is a shortest path. As a consequence one obtains that  $u \leq_b v$  in  $G$  implies  $d_{G^\Delta}(b, u) \leq d_{G^\Delta}(b, v)$ .

Suppose now that there exists another normal folder-path  $P'_{(v,b)}$ , and let  $x$  be the neighbor of  $v$  in it. We may assume that  $P_{(x,b)}$  is the unique normal folder-path connecting  $x$  and  $b$ . This implies  $d_G(x, b) \leq d_G(v, b)$ . If  $v$  and  $x$  are adjacent in  $G$ , then the folders spanned by  $vv^b$  and  $xx^b$  intersect in an edge  $xv^b$ , contrary to the fact that  $F_{vv^b} \in \text{star}(vx, \mathcal{X})$ . Finally, if

$d_G(v, b) = d_G(x, b)$ , we obtain the same contradiction: the cell spanned by  $vx$  intersects  $F_{xx^b}$  in the edge  $xm$ , where  $m$  denotes the median of  $v, x, b$ .

Let  $B_r^\Delta(b)$  be a ball in  $G^\Delta$  with center  $x$  and radius  $r$ . Using Lemma 4.2 and the induction on  $r$  we will show that  $B_r^\Delta(b)$  is a convex set in  $G$ . Pick  $y, z \in B_r^\Delta(b)$  with  $d_G(y, z) = 2$  and let  $x$  be their common neighbor in  $G$ . If  $d_G(b, x) < \max\{d_G(b, y), d_G(b, z)\}$  then  $d_{G^\Delta}(b, x) \leq \max\{d_{G^\Delta}(b, y), d_{G^\Delta}(b, z)\} \leq r$ , and we are done. So assume that  $k = d_G(b, y) = d_G(b, z) < d_G(b, x)$ . If  $x^b$  is the unique neighbor in  $B_{k-1}(b)$  of  $y$  or of  $z$ , we infer that  $x \in B_r^\Delta(b)$ . Hence in  $B_{k-1}(b)$  we can find the neighbors  $u$  and  $w$  of  $y$  and  $z$ , respectively. If  $y^b = z^b$  then  $(x, y, u, y^b = z^b, z, x)$  is an induced 6-cycle of  $G$ . So,  $y^b \neq z^b$ . By the induction hypothesis  $d_{G^\Delta}(b, x_b) \leq \max\{d_{G^\Delta}(b, y^b), d_{G^\Delta}(b, z^b)\} \leq r - 1$ , leading us to the conclusion that  $x \in B_r^\Delta(b)$ . Therefore  $B_r^\Delta(b)$  is a convex set of  $G$ , and by Lemma 4.2 it is gated in  $G$ . From the Helly property for gated sets we infer that  $G^\Delta$  is a Helly graph. ■

**LEMMA 7.6.** *Let  $F$  be a folder of  $G$  and let  $x, y \notin F$  be two vertices of  $G$  whose gates  $g_x, g_y$  in  $F$  are distinct and nonadjacent. Then  $d_G(x, y) = d_G(x, g_x) + d_G(g_y, y) + 2$ ; i.e.,  $g_x$  and  $g_y$  lie on a common shortest path between  $x$  and  $y$ .*

*Proof.* We argue by induction on  $k = d_G(x, g_x) + d_G(y, g_y)$ . Suppose by way of contradiction that  $d_G(x, y) < k + 2$ . First, let  $k = 2$ ; i.e.,  $x, g_x$  are adjacent and  $y, g_y$  are adjacent. Then  $d_G(x, y) = 2$ . Take a common neighbor  $z$  of  $x$  and  $y$  and two common neighbors  $u, v \in F$  of  $g_x$  and  $g_y$ . To avoid an induced 6-cycle,  $z$  must be adjacent to both  $u$  and  $v$ . But then the subgraph induced by  $x, z, y, u, v, g_x, g_y$  contains the forbidden  $K_{3,3}^-$ . So, let  $k \geq 3$ , say  $d_G(x, g_x) \geq 2$ . In  $I(x, g_x)$  pick a neighbor  $w$  of  $x$  and a neighbor  $p \neq x$  of  $w$ . Clearly  $g_x$  is also the gate of  $w$  and  $p$ . By the induction hypothesis  $d_G(w, y) = k + 1$ . Since  $x, p \in I(w, y)$ , we will find a common neighbor  $w'$  of  $p$  and  $x$  one step closer to  $y$ . As  $w' \in I(x, g_x)$ , we arrive at a contradiction to the induction assumption. ■

We continue by presenting a special subdivision of  $\mathcal{X} = \mathcal{X}(G)$  ([38] presents a related “orbit splitting” operation which allows us to transform edge weighted semiframes into frames). Let  $E_i$   $i \in \Lambda$  be the classes of the equivalence relation  $\Psi$  defined in the previous section. Clearly, all edges from a folder  $K_{2,n}$ ,  $n \geq 3$ , belong to one equivalence class. If  $G$  is an F-graph then all edges in  $E_i$  have the same length  $\lambda_i$ . An *ij-rectangle* is a rectangle of  $\mathcal{X}$  with two opposite edges in  $E_i$  and two other opposite edges in  $E_j$ . In this case define  $\lambda_{ij} = \lambda_i/\lambda_j$ .

Let  $0 < \lambda < \lambda_i$ . Consider an *ij-rectangle*  $R = xuvy$  with  $xy, uv \in E_i$  and  $xu, vy \in E_j$ . Define the points  $p(x), p(y) \in [x, y]$ ,  $p(u), p(v) \in [u, v]$ ,

$q(x), q(u) \in [x, u], q(y), q(v) \in [y, v]$ , such that

$$|x - p(x)| = |y - p(y)| = |u - p(u)| = |v - p(v)| = \lambda$$

and

$$|x - q(x)| = |y - q(y)| = |u - q(u)| = |v - q(v)| = \lambda \cdot \lambda_{ji}.$$

Subdivide  $R$  into nine rectangles by means of the segments  $[p(x), p(u)], [p(y), p(v)], [q(x), q(y)],$  and  $[q(u), q(v)]$ . In  $\mathcal{X}(G)$   $R$  is a union of two right triangles with a common hypotenuse  $h$ . Each of these triangles will be subdivided into three rectangles and three right triangles whose hypotenuses constitute  $h$ . Refine this subdivision further by taking in each of three rectangles the diagonal parallel to  $h$ . If we provide this operation with all  $ij$ -rectangles, we obtain a 2-complex  $\mathcal{X}'$  whose 2-cells are right triangles. Clearly  $\mathcal{X}'$  obeys the conditions (a), (b), and (c). Since  $\mathcal{X}$  and  $\mathcal{X}'$  have the same underlying space,  $\mathcal{X}'$  is CAT(0). By Theorem 7.1, the leg graph of  $\mathcal{X}'$  is an F-graph. More generally, given a finite subset of equivalence classes  $\Lambda' \subseteq \Lambda$ , we can apply the same procedure step by step for all  $i \in \Lambda'$ , yielding an F-complex and an F-graph as the leg graph. We summarize this in the following property.

LEMMA 7.7. *Performing the subdivision operation with respect to a finite subset of equivalence classes of an F-graph we obtain again an F-graph.*

Recently, Karzanov [37] presented a characterization of minimizable graphs (alias frames); we refrain from giving definitions here, referring to [37, 38] for precise formulations and interesting details. These are hereditary modular graphs whose edges can be oriented so that in every  $ij$ -rectangle each pair of opposite edges has the same orientation. The main (and rather difficult) step in the proof of the result from [37] is to associate with each frame  $G$  the  $B_2$  complex  $\mathcal{X}(G)$  endowed with the intrinsic  $l_1$ -metric and to prove that every tight extension of the graph-metric  $d_G$  is a subspace of this metric space. Below we show that for an F-complex  $\mathcal{X}$  the intrinsic  $l_1$ -metric is hyperconvex. In view of the result of Aronszajn and Panitchpakdi [4] characterizing hyperconvex spaces as injective spaces, the result of [37] will follow. In [34] Isbell introduced collapsible cubical 2-complexes and proved that their rectilinear metric is injective. One can show that such complexes are exactly the CAT(0)  $A_1 \times A_1$  complexes (alias cube-free median complexes).

Let  $\mathcal{X}$  be a 2-complex obeying the requirements (a), (b), (c). Metrize  $\mathcal{X}$  so that each solid  $ij$ -rectangle is a copy of the rectangle  $[0, \lambda_i] \times [0, \lambda_j]$  in the  $l_1$ -plane; define the distance between two points not in a common rectangle as the length of a shortest piecewise-linear path joining them. The length of such a path  $P$  is computed in the following way: suppose

that there is a subdivision  $t_0 < \dots < t_k$  such that  $P(t_i, t_{i+1})$  is contained in some cell. Then sum up the lengths of  $P(t_i, t_{i+1})$ , the length inside a cell being measured with respect to the  $l_1$ -metric. We denote the obtained metric space by  $(\mathcal{X}, \rho_1)$ . One can see that the restriction of  $\rho_1$  on  $LG(K)$  coincides with the weighted shortest-path metric. If  $\mathcal{X}$  is a  $B_2$  complex, then  $LG(K)$  endowed with the standard graph-metric  $d_{LG(K)}$  is isometrically embedded in  $(\mathcal{X}, \rho)$ . Since the F-graphs are hereditary modular, from [8, 10] we deduce that in F-complexes  $\mathcal{X}$  every shortest path of  $(LG(K), \rho)$  is a shortest path of  $(LG(K), d_{LG(K)})$  and vice versa. Now we formulate the second characterization of F-complexes.

**THEOREM 7.8.** *A locally-finite 2-complex  $\mathcal{X}$  satisfying (a), (b), and (c) is an F-complex if and only if  $(\mathcal{X}, \rho_1)$  is hyperconvex.*

*Proof.* Assume that the metric  $\rho_1$  is injective (thus  $\mathcal{X}$  is simply connected, because as is shown in [4, p. 422, 34, Theorem 1.1] injective spaces are contractible); however,  $\mathcal{X}$  is not an F-complex. First, let  $\mathcal{X}$  contain three folders  $F_{h_1}, F_{h_2}, F_{h_3}$  and three distinct legs  $e_1, e_2, e_3$  sharing a common vertex such that  $e_i$  belongs to  $F_{h_j}$  exactly when  $i \neq j$ . Denote by  $\nu_1, \nu_2, \nu_3$  the lengths of  $e_1, e_2, e_3$ , respectively. In  $F_{h_i}$  pick a vertex  $v_i$  which does not belong to other two folders. Set  $r_i = \max\{\nu_j, \nu_k\}$ , where  $i \neq j, k$ . One can verify that the balls  $B_{r_1}(v_1), B_{r_2}(v_2), B_{r_3}(v_3)$  pairwise intersect, but their common intersection is empty. Second, suppose that two folders  $F_{h_1}$  and  $F_{h_2}$  have two legs  $xy$  and  $xz$  in common. Let  $v \in F_{h_1}$  and  $w \in F_{h_2}$  be some common neighbors of  $y$  and  $z$  in  $LG(K)$ . Let  $\mu_1, \mu_2$  be the lengths of  $xy, xz$ , where  $\mu_1 \leq \mu_2$ . If  $v \neq w$ , consider the collection of balls with  $x, y, z, v, w$  as centers and the following radii:  $r(x) = r(y) = r(z) = \mu_2$  and  $r(v) = r(w) = \mu_1$ . The balls pairwise intersect; however, their intersection is empty, because  $B_{\mu_2}(x), B_{\mu_1}(v), B_{\mu_1}(w)$  intersect in at most two points  $z, y$ , but  $z \notin B_{\mu_2}(y), y \notin B_{\mu_2}(z)$ . So assume that  $v = w$ ; i.e., we have two solid rectangles  $R_1, R_2$  with the common boundary  $xyvz$ . Take the balls  $B_{\mu_2}(x), B_{\mu_2}(y), B_{\mu_1}(z), B_{\mu_1}(v)$ . They intersect in two interior points  $p_1 \in R_1, p_2 \in R_2$ . Add the balls centered at  $p_1$  and  $p_2$ , each having radius  $\mu_1/2$ . We will get a collection of six pairwise intersecting balls with an empty intersection, a contradiction.

As to the converse, since the balls are compact, it suffices to verify the Helly property only for finite collections of balls. It will be enough to check this only for finite F-complexes  $\mathcal{X}$ . Indeed, any finite collection of balls is contained in the subcomplex spanned by a sufficiently large ball of the leg graph. Since  $\mathcal{X}$  is locally finite, this ball is finite and by Lemma 7.3 it spans an F-complex. Second, it suffices to verify the Helly property for pairwise intersecting balls whose centers are all located at vertices of  $\mathcal{X}$ . Indeed, we can subdivide sufficiently many times  $\mathcal{X}$  as described before, arriving at a

new complex with the same underlying space as  $\mathcal{X}$ , but in which all ball-centers are vertices. By Lemma 7.7 the resulting complex is an F-complex as well. Finally, we will allow balls with negative radii, by letting  $B_r(x) := \{x\}$  if  $r < 0$ .

So assume that  $\mathcal{X}$  is a finite F-complex, and let  $\mathcal{B} = \{B_{r_k}(x_k) : k = 1, \dots, n\}$  be a collection of pairwise intersecting balls whose centers are vertices of  $\mathcal{X}$ . Suppose that the result holds for any collection of less than  $n$  balls. The result is also true if  $\mathcal{X}$  is a folder  $F$ . In this case we will find a common point of balls on the hypotenuse  $h_F$  if all  $x_i$  are located at the ends of  $h_F$ , or in a triangle containing a ball-center with the smallest radius. So assume that  $\mathcal{X}$  contains at least two cells. To prove that  $\bigcap_{k=1}^n B_{r_k}(x_k) \neq \emptyset$  we proceed by induction on the number of vertices of  $\mathcal{X}$ . We will show how to replace in  $\mathcal{B}$  certain balls by balls which are their proper subsets, to obtain a new collection  $\mathcal{B}'$  of pairwise intersecting balls in a proper subcomplex of  $\mathcal{X}$ .

By Lemma 7.4 we can find a vertex  $v$  which belongs to a unique edge or to a pendant folder  $F$  (as  $v$  one can take any vertex at maximum distance from a base point). Let  $\mathcal{X}'$  be the subcomplex of  $\mathcal{X}$  obtained by removing  $v$  and all (open) cells containing  $v$ . Notice that  $\mathcal{X}'$  is an isometric subspace of  $\mathcal{X}$ . If  $v \notin X$ , then replace each ball of  $\mathcal{B}$  by its intersection with  $\mathcal{X}'$  and apply induction. So we may assume that  $v = x_1$ .

First suppose that  $F$  contains at least three triangles and that  $x_1$  is not incident to the hypotenuse  $h_F$  of  $F$ . From the choice of  $x_1$  we deduce that  $x_1$  and all non-neighbors of  $x_1$  in  $F$ , except possibly one vertex, have degree 2 in  $LG(\mathcal{X})$ . Delete from  $\mathcal{B}$  the balls around such vertices except one vertex with the smallest radius. The reduced family of pairwise intersecting disks has a point  $p$  in common. Clearly,  $p$  belongs to all deleted balls, and by the induction assumption we are done. So, further we may assume that  $h_F = [x_1, z]$ . If  $F$  is a rectangle, then its edges have two lengths  $\lambda$  and  $\mu$  (which may coincide), otherwise all edges of  $F$  have the same length  $\lambda = \mu$ .

For a ball-center  $x_k$  ( $k > 1$ ) by  $g_k$  denote its gate in  $F$ , and set  $r'_k := r_k - \rho_1(x_k, g_k)$ . Let  $r'_1 := r_1 - \lambda$ ,  $r''_1 := r_1 - \mu$ . First notice that

$$r_1 + r'_k \geq \lambda + \mu \tag{1}$$

for any vertex  $x_k$  with  $g_k = z$ . Second, if the gates  $g_i$  and  $g_j$  of  $x_i$  and  $x_j$  are distinct and both different from  $z$ , then

$$r'_i + r'_j \geq \lambda + \mu. \tag{2}$$

Indeed, by Lemma 7.6 and since the shortest paths in  $(LG(\mathcal{X}), \rho_1)$  and  $(LG(\mathcal{X}), d_{LG(\mathcal{X})})$  are the same, we conclude that  $r_i + r_j \geq \rho_1(x_i, x_j) = \rho_1(x_i, g_i) + \lambda + \mu + \rho_1(g_j, x_j)$ , yielding the required inequality.

Pick a vertex  $w$  of  $F$ ,  $w \neq x_1, z$ , and suppose that  $\rho_1(w, x_1) = \lambda$ . Replace  $B_{r_1}(x_1)$  by  $B_{r'_1}(w)$ . Note that  $r_k + r'_1 \geq \rho_1(x_k, w)$  for all  $x_k$  whose gates in

$F$  coincide with  $z$  or  $w$ . If this inequality is true when  $g_k$  and  $w$  are not adjacent, by the induction hypothesis the balls intersect. Since  $B_{r'_1}(w) \subset B_{r_1}(x_1)$ , we are done. Hence, further we may assume that there exists at least one  $x_j$  such that  $\rho_1(x_1, g_j) = \mu$  and  $r'_1 + r_j < \lambda + \mu + \rho_1(x_j, g_j)$ ; i.e.,

$$r_1 + r'_j < 2\lambda + \mu. \tag{3}$$

Similarly, we may assume that there exists a vertex  $x_i$  such that  $\rho_1(x_1, g_i) = \lambda$  and

$$r_1 + r'_i < 2\mu + \lambda. \tag{4}$$

Replace every ball  $B_{r'_k}(x_k)$  ( $k > 1$ ) by  $B_{r'_k}(g_k)$ . Together with  $B_{r_1}(x_1)$  these balls constitute  $\mathcal{B}'$ . Notice that  $r_1 + r'_k \geq \lambda + \mu$  for all  $x_k$  with  $g_k = z$ . On the other hand, by (2)  $r'_i + r'_j \geq \lambda + \mu$  for all  $x_i, x_j$  whose gates are distinct and both different from  $z$ . If  $r'_i + r'_k \geq \rho_1(g_i, g_k)$  for all  $g_i \neq g_k = z$ , then in  $F$  we find a common point of balls from  $\mathcal{B}'$ , which belongs to all balls from  $\mathcal{B}$  as well. Hence,  $r'_{i^*} + r'_{k^*} < \rho_1(g_{i^*}, g_{k^*})$  for some  $g_{i^*}, g_{k^*}$  with  $g_{i^*} \neq g_{k^*} = z$ . Suppose without loss of generality that  $\rho_1(x_1, g_{i^*}) = \lambda$ ; i.e., the previous inequality can be rewritten as

$$r'_{i^*} + r'_{k^*} < \mu. \tag{5}$$

Evidently  $x_{i^*}$  may be assumed to satisfy the inequality (4). From (3) and (5) we conclude that  $r_1 + r'_j + r'_{i^*} + r'_{k^*} < 2\lambda + 2\mu$ . On the other hand, from (1) and (2) we have  $2\lambda + 2\mu \leq r_1 + r'_{i^*} + r'_j + r'_{k^*}$ , thus yielding a final contradiction. ■

An alternative metrization of an F-complex  $\mathcal{X}$  is obtained by extending the  $l_\infty$ -metric from the cells. Each  $ij$ -rectangle is a copy of the rectangle  $[0, \lambda_i] \times [0, \lambda_j]$  in the  $l_\infty$ -plane. As in the previous section, denote the resulting intrinsic metric on  $\mathcal{X}$  by  $\rho_\infty$ . The (weighted) Helly graph  $G^\Delta$  associated to the 1-skeleton  $G$  of  $\mathcal{X}$  is isometrically embedded in  $(\mathcal{X}, \rho_\infty)$ . This suggests that  $(\mathcal{X}, \rho_\infty)$  is injective. We only sketch the proof of this property in the case when all cells are isosceles right triangles. It suffices to show that any ball  $B_r(x)$  of  $(\mathcal{X}, \rho_\infty)$  is a convex set of  $(\mathcal{X}, \rho_1)$  (the second space being modular, all its compact convex sets are gated). It is sufficient to show this for finite F-complexes and balls with rational radii. Since regular subdivision of all cells with some given mesh yields again an F-complex, it suffices to consider a ball  $B_r(x)$  whose center is a vertex and whose radius is an integer. Then one can easily see that  $B_r(x)$  is a union of cells of  $\mathcal{X}$ . Moreover, from the proof of Proposition 7.5 we infer that if two non-adjacent vertices of some folder  $F$  belong to  $B_r(x)$ , then the whole folder lies in  $B_r(x)$ . By Lemma 4.2 the subgraph of  $G$  induced by  $B_r(x)$  is convex in  $G$ . This implies that  $B_r(x)$  is convex in  $(\mathcal{X}, \rho_1)$ . Therefore, we have proven the following property of flag complexes.

PROPOSITION 7.9. *If  $\mathcal{X}$  is a locally finite  $F$ -complex, then  $(\mathcal{X}, \rho_\infty)$  is an injective space.*

### 8. BRIDGED COMPLEXES

Every bridged graph  $G$  gives rise to an abstract simplicial complex  $K(G)$  whose simplexes are the complete subgraphs of  $G$ . Trivially,  $K(G)$  is a flag complex. The geometric realization  $|K(G)|$  is called a *bridged complex*. In this section we present a characterization of flag complexes whose underlying graphs are bridged.

THEOREM 8.1. *For a simplicial flag complex  $K$  the following conditions are equivalent:*

- (i)  *$K$  is simply connected and the link of every vertex  $v$  does not contain induced 4-cycles and 5-cycles;*
- (ii)  *$G(K)$  is weakly modular and does not contain induced 4-cycles and 5-cycles;*
- (iii)  *$G(K)$  is bridged.*

*Proof.* (i)  $\Rightarrow$  (ii): We actually need a weaker version of the flag condition: if three 2-simplexes share a common vertex and pairwise share common edges then they are contained in a 3-simplex of  $K$ . Consider a simple cycle  $C$  in the graph  $G(K)$  and let  $(D, f)$  be a minimal disk of  $C$ .

*Claim 1.* The degree of every interior vertex  $x$  of  $D$  is at least 6.

Let  $x_1, \dots, x_n$  be the neighbors of  $x$ , where  $\sigma_i = xx_i x_{i+1(\text{mod } n)}$  ( $i = 1, \dots, n$ ) are the faces of  $D$  incident to  $x$ . Trivially,  $n \geq 3$ . Suppose by way of contradiction that  $n \leq 5$ . From Lemma 5.1 we conclude that  $f(\sigma_1), \dots, f(\sigma_n)$  are distinct 2-simplexes of  $K$ . If  $n = 3$  then  $f(\sigma_1), f(\sigma_2), f(\sigma_3)$  intersect in  $f(x)$  and pairwise share edges of  $K$ . Therefore they are contained in a 3-simplex of  $K$ . This implies that  $\delta = f(x_1)f(x_2)f(x_3)$  is a 2-face of  $K$ . Let  $D'$  be a disk triangulation obtained from  $D$  by deleting the vertex  $x$  and the triangles  $\sigma_1, \sigma_2, \sigma_3$ , and adding the 2-simplex  $x_1x_2x_3$ . The map  $f: V(D') \rightarrow V(K)$  is simplicial, because it maps  $x_1x_2x_3$  to  $\delta$ . Therefore  $(D', f)$  is a disk diagram for  $C$ , contrary to the choice of  $D$ . Now, let  $x$  have four neighbors. The cycle  $(x_1, x_2, x_3, x_4, x_1)$  is sent to a 4-cycle in  $\text{link}(f(x), K)$ , in which two opposite vertices, say  $f(x_1)$  and  $f(x_3)$ , are adjacent. Consequently,  $\delta' = f(x_1)f(x_3)f(x_2)$  and  $\delta'' = f(x_1)f(x_3)f(x_4)$  are 2-faces of  $K$ . Let  $D'$  be a disk triangulation obtained from  $D$  by deleting the vertex  $x$  and the triangles  $\sigma_i (i = 1, \dots, 4)$  and adding the 2-simplexes  $\sigma' = x_1x_3x_2$  and  $\sigma'' = x_1x_3x_4$ . The map  $f$  remains simplicial, since it sends  $\sigma', \sigma''$

to  $\delta'$ ,  $\delta''$ , respectively, contrary to the choice of  $D$ . Finally, suppose that  $n = 5$ . Consider the 5-cycle of  $\text{link}(f(x), K)$  induced by the vertices  $f(x_1), f(x_2), f(x_3), f(x_4), f(x_5)$ . By the initial hypothesis concerning  $K$ , we deduce that there is a vertex, say  $f(x_1)$ , which is adjacent to all other vertices of this cycle. This implies that  $f(x_1)f(x_3)f(x_2)$ ,  $f(x_1)f(x_4)f(x_3)$ , and  $f(x_1)f(x_5)f(x_4)$  are 2-faces of  $K$ . Let  $D'$  be obtained from  $D$  by deleting the vertex  $x$  and the faces  $\sigma_i (i = 1, \dots, 5)$  and adding three new faces  $x_1x_3x_2$ ,  $x_1x_4x_3$ , and  $x_1x_5x_4$ . As in preceding cases, we can conclude that  $(D', f)$  is a disk diagram for  $C$ , yielding a contradiction to the minimality of  $C$ .

From this claim we obtain that any minimal disk  $D$  is a (3,6)-complex and thus a CAT(0) space. The turning angle  $\tau(v)$  at any corner  $v$  of  $D$  is  $2\pi/3$  or  $\pi/3$ . The corners of first type are the boundary vertices of degree two. The corners of second type are boundary vertices of degree three. In the first case two neighbors of  $v$  are adjacent. In the second case  $v$  and its neighbors in  $\partial D$  are adjacent to the third neighbor of  $v$ . From the Gauss-Bonnet formula we infer that  $D$  contains at least three corners, and if  $D$  has exactly three corners then all are of first type. Together with the claim this ensures that all 3-cycles of  $G(K)$  bound 2-simplexes of  $K$  and that  $G(K)$  does not contain induced 4-cycles and 5-cycles.

Next we show that  $G(K)$  is weakly modular. To verify the triangle condition, pick three vertices  $u, v, w$  with  $1 = d_{G(K)}(v, w) < d_{G(K)}(u, v) = d_{G(K)}(u, w) = k$ . We claim that if  $I(u, v) \cap I(u, w) = \{u\}$ , then  $k = 1$ . Suppose not. Pick two shortest paths  $P'$  and  $P''$  joining the pairs  $u, v$  and  $u, w$ , respectively, such that the cycle  $C$  composed of  $P', P''$  and the edge  $vw$  has minimum area. Choose a minimal disk  $D$  of  $C$ , which necessarily has a corner  $x$  distinct from  $u, v, w$ . Let  $x \in P'$ . Notice that  $x$  is a corner of second type; otherwise its neighbors  $y, z$  in  $P'$  are adjacent, contrary to the assumption that  $P'$  is a shortest path. Let  $p$  be the vertex of  $D$  adjacent to  $x, y, z$ . If we replace in  $P'$  the vertex  $x$  by  $p$ , we obtain a new shortest path between  $u$  and  $v$ . Together with  $P''$  and the edge  $vw$  this path forms a cycle  $C'$  whose area is strictly smaller than  $\text{Area}(C)$ , contrary to the choice of  $C$ . As to the quadrangle condition, suppose by way of contradiction that we can find distinct vertices  $u, v, w, z$  such that  $v, w \in I(u, z)$  are neighbors of  $z$  and  $I(u, v) \cap I(u, w) = \{u\}$ ; however,  $u$  is not adjacent to  $v$  and  $w$ . Again, select two shortest paths  $P'$  and  $P''$  between  $u, v$  and  $u, w$ , respectively, so that the cycle  $C$  composed of  $P', P''$  and the edges  $vz$  and  $zw$  has minimum area. Choose a minimal disk  $D$  of  $C$ . From the hypothesis concerning the vertices  $u, v, w, z$  we deduce that  $D$  has at most one corner of first type located at  $u$ . Hence  $D$  contains at least four corners. If one corner  $x$  is distinct from  $u, v, w, z$ , then proceeding in the same way as before, we obtain a contradiction with the choice of  $P', P''$ . Therefore  $u, v, w, z$  are the only corners of  $D$ , where  $\tau(u) = 2\pi/3, \tau(v) = \tau(w) = \tau(z) = \pi/3$ .

Since the sum of turning angles at corners is smaller than  $2\pi$ , we arrive at a contradiction with the Gauss–Bonnet formula. The proof of (ii)  $\Rightarrow$  (iii)  $\Rightarrow$  (i) follows from the following result.

*Claim 2.* A graph  $G$  is bridged if and only if it is weakly modular and does not contain 4-cycles and 5-cycles as induced subgraphs.

If  $G$  is bridged, then trivially  $G$  does not contain induced 4-cycles and 5-cycles. Weak modularity of  $G$  easily follows from the fact that the cycles of  $G$  are well bridged. Conversely, let  $G$  be a weakly modular graph without induced 4-cycles and 5-cycles. To show that  $G$  is bridged, by Theorem 4.5 and since  $B_{k+r}(S) = B_k(B_r(S))$ , it suffices to show that if  $S$  is convex then the neighborhood  $B_1(S)$  of  $S$  is convex, too. By Lemma 4.2 it suffices to check 2-convexity only. Pick two vertices  $x, y \in B_1(S)$  at distance two. We assume without loss of generality that  $x, y \notin S$ ; otherwise we immediately obtain  $I(x, y) \subseteq B_1(S)$ . Let  $x'$  and  $y'$  be neighbors of  $x$  and  $y$  in  $S$  minimizing  $d_G(x', y')$ . Suppose by way of contradiction that there is a common neighbor  $z$  of  $x$  and  $y$  outside  $B_1(S)$ . Since  $S$  is convex,  $d_G(x', y') \leq 3$ . If  $x'$  and  $y'$  were adjacent, then the vertices  $x', x, z, y, y'$  would induce a forbidden cycle. If  $d_G(x', y') = 3$ , then necessarily  $d_G(x, y') = 3$ . By the triangle condition there exists a common neighbor  $u$  of  $x$  and  $x'$  at distance 2 to  $y'$ . Since  $S$  is convex,  $u \in S$ , contrary to the choice of  $x'$ . Finally, assume  $d_G(x', y') = 2$ , and let  $z'$  be adjacent to  $x'$  and  $y'$ . If  $d_G(z, z') = 3$ , by the quadrangle condition we find a common neighbor  $v$  of  $x', y', z$ . Since  $v \in I(x', y') \subseteq S$ , we arrive at a contradiction to the choice of  $z$ . Otherwise, if  $d_G(z', z) = 2$ , by the triangle condition we can find a common neighbor  $s$  of  $z, x', z'$  and a common neighbor  $t$  of  $z, z', y'$ . In order to avoid forbidden 4-cycles and 5-cycles, the vertices  $s, t$  must be adjacent to  $x, x', y, y'$ . This implies  $s, t \in I(x', y') \subseteq S$ , contrary to the choice of  $z$ . This concludes the proof. ■

**COROLLARY 8.2.**  *$\mathcal{X}$  is a CAT(0) complex of type  $A_2$  if and only if it is a bridged 2-complex. In particular, any triangle building is bridged.*

Another example of a CAT(0) complex whose 1-skeleton is bridged is provided by any tiling of the plane into acute angled triangles.

Although bridged graphs are characterized by a discrete variant of a property describing CAT(0) spaces (see the remarks after Theorem 3.2), not all bridged complexes are CAT(0). We present two such examples. First, take a 6-wheel consisting of a 6-cycle  $(x, u_1, u_2, y, u_3, u_4, x)$  and a central vertex  $z$ , and replace the vertices  $u_1, u_2, u_3, u_4$  by  $n$ -cliques  $C_1, C_2, C_3, C_4$ . Make  $x$  adjacent to all vertices of  $C_1$  and  $C_4$ ,  $y$  to all vertices of  $C_2$  and  $C_3$ , and  $z$  to the vertices of all four cliques. Finally, assume that  $C_1 \cup C_2$  and  $C_3 \cup C_4$  form  $(n + 1)$ -cliques. We obtain a new bridged graph  $G_n$ . One can imagine  $G_n$  as glued from two identical halves along the path

$(x, z, y)$ .  $G_1$  coincides with the initial 6-wheel. In this case  $(x, z, y)$  is the unique geodesic in  $|G_1|$  connecting the vertices  $x$  and  $y$ . This is no longer true in  $|G_n|$  for sufficiently large  $n$ . Indeed, let  $b_i$  be the barycenter of the simplex  $|C_i \cup \{z\}|$ ,  $i = 1, \dots, 4$ . Then  $d(x, b_1) = d(x, b_4) = d(y, b_2) = d(y, b_3) = \sqrt{\frac{n+1}{2n}}$  and  $d(b_1, b_2) = d(b_3, b_4) = \frac{\sqrt{n-1}}{n}$ . The paths  $(x, b_1, b_2, y)$  and  $(x, b_4, b_3, y)$  have length  $\sqrt{\frac{2(n+1)}{n}} + \frac{\sqrt{n-1}}{n}$ , which is smaller than 2. This shows that the path  $(x, z, y)$  is not a geodesic. Due to the symmetry between the halves, we conclude that  $x$  and  $y$  can be connected with more than one shortest path, showing that  $|G_n|$  is not CAT(0).

Even not all bridged 3-complexes are CAT(0). For this, let  $\Gamma$  be the bridged graph with vertices  $z, x_1, x_2, x_3, u_1, u_2, v_1, v_2, w_1, w_2$ , where  $z$  is adjacent to all vertices,  $u_1, u_2$  are adjacent to  $x_1, x_2$ ,  $v_1, v_2$  are adjacent to  $x_2, x_3$ , and  $w_1, w_2$  are adjacent to  $x_3, x_1$ . In the link of  $z$  consider the geodesic segments joining each pair of vertices  $x_1, x_2, x_3$ . For example, the geodesic segment between  $x_1$  and  $x_2$  consists of two edges sharing the midpoint of the segment  $u_1 u_2$ . Their union is an isometric cycle of  $\text{Link}(z, |\Gamma|)$  of length  $5\pi/2$ ; thus  $|\Gamma|$  is not CAT(0).

Bridson [16] established that all simplicial 3-complexes whose cells are regular and whose links of all vertices are PE (3,6)-disks are CAT(0). Clearly, such complexes are bridged. Following the lines of the proof from [16], one can show that all bridged 3-complexes in which the links of vertices are bridged are CAT(0). Bridged complexes  $K$  in which the links of all simplexes are bridged form a very special class of simplicial complexes. As we will show below their skeletons do not contain induced cycles of length larger than three (such graphs are the well known *chordal graphs*).

**COROLLARY 8.3.** *The links of all simplexes of a simply connected flag complex  $K$  are bridged if and only if the 1-skeleton  $G(K)$  is a chordal graph.*

*Proof.* Let  $\sigma$  be a maximal simplex whose link contains an induced cycle  $C$  of length larger than three. Since  $\text{link}(\sigma, K)$  is bridged, one can construct a minimal disk  $D$  for  $C$ . We may assume that all cycles of length  $\geq 4$  of  $\text{link}(\sigma, K)$  whose minimal disks have smaller area are not induced. From the proof of Theorem 8.1 we know that  $D$  is a (3,6)-disk; thus it has a corner  $x$ . Since  $C$  is induced, the turning angle at  $x$  is  $\pi/3$ ; i.e., it belongs to exactly two faces  $xyz_1$  and  $xyz_2$  of  $D$ . The cycle  $C'$  obtained from  $C$  by replacing  $x$  by  $y$  has smaller area; thus it is not induced. Therefore  $y$  is adjacent to some vertex  $v \neq x$  of  $C$ . The edge  $yv$  splits  $C'$  into two cycles of smaller area. Applying the above argument to each of these cycles, we deduce that  $y$  is adjacent to some their vertices distinct from  $z_1, v$ , and  $z_2$ . Continuing this way, we obtain that  $y$  is adjacent to all vertices of  $C$ . This means that  $C$  is a cycle in the link of the simplex  $\sigma \cup \{y\}$ , contrary to the

choice of  $\sigma$ . The converse is immediate, because the skeletons of links of all simplexes will be chordal graphs. ■

It is well known (cf. [32]) that every finite chordal graph  $G$  has a simplicial vertex, i.e., a vertex  $x$  such that its neighbors form a complete subgraph. If  $G$  is the graph of a complex  $K$ , this means that  $x$  belongs to a unique maximal simplex  $\sigma$ . Consequently,  $|K|$  can be obtained by gluing  $|\sigma|$  and  $|K'|$ , where  $K'$  is the subcomplex of  $K$  spanned by  $\sigma' = \sigma - \{x\}$  and the maximal simplexes distinct from  $\sigma$ . Since the gluing is performed along a convex set  $|\sigma'|$ , from the result of Rechetniak (cf. [17, Theorem 4.5]) we obtain that  $|K|$  is CAT(0) if and only if  $|\sigma|$  and  $|K'|$  are CAT(0). In particular, we have the following observation (to pass from finite to arbitrary complexes we use Theorem 3.2).

**COROLLARY 8.4.** *If the graph of a flag complex  $K$  is chordal, then  $|K|$  is CAT(0).*

We conclude this section with the following interesting characterization of bridged subgraphs of bridged graphs.

**COROLLARY 8.5.** *An induced subgraph  $G'$  of a bridged graph  $G$  is bridged if and only if the simplicial complex  $K(G')$  is connected and simply connected.*

## 9. COMBINGS AND BICOMBINGS OF SOME GRAPHS: CLIN D'OEIL

Both notions of combing and bicombing have been introduced in [28] for standard word metric on the Caley graph of a group in order to define an automatic or a biautomatic structure on the group. A (geodesic) combing of a group is a selection for each element  $g$  of one shortest path beginning at the identity and ending at  $g$ . A reasonable requirement to a combing is that the paths to neighboring points are uniformly close. A bicombing is a choice of shortest paths between each pair of points which, in some sense, imitates the convexity of the distance function between geodesics in a CAT(0) space. The study of infinite groups by means of the geometry of their Caley graphs has proven to be a powerful technique for understanding large classes of groups; [19, 28–30, 44, 49] are only few references.

In this section we investigate the concepts of combing and bicombing in the framework of graphs. We show that for all classes of graphs introduced in the previous sections there exist combings and bicombings having a tight fellow traveler property. Their particular instances are Caley graphs of groups acting on some CAT(0) complexes, for example,  $A_2$  and  $B_2$  complexes and cubings.

During this section we assume that the graphs are locally finite (i.e., every vertex has a finite number of neighbors). Let  $[0, n]^*$  denote the set of integer points from the segment  $[0, n]$ . Given a path  $p$  of length  $n$  of a graph  $G$ , we can parameterize it and denote  $p: [0, n]^* \rightarrow G$ . It is often convenient to extend  $p$  over a larger interval,  $p: [0, m]^* \rightarrow G$ , for some  $m > n$ . We do this by setting  $p(t) = p(n)$  for all  $t \geq n$ . (In particular, when comparing two such paths, we may assume that they are defined on the same interval.)

Suppose that  $p, q: [0, n]^* \rightarrow G$  are two paths in  $G$  and  $k$  a positive integer. We say that  $p$  and  $q$  are *k-fellow travelers* if  $d_G(p(t), q(t)) \leq k$  for all integers  $t$  with  $0 \leq t \leq n$ . Following [29], we say that two paths  $p, q: [0, n]^* \rightarrow G$  lie in a *k-Hausdorff neighborhood* of each other if every vertex of  $p$  is at distance at most  $k$  from  $q$  and vice versa; i.e., for every  $t \in [0, n]$  there exist  $s', s'' \in [0, n]$  such that  $d_G(p(t), q(s')) \leq k$  and  $d_G(p(s''), q(t)) \leq k$ .

Let  $b$  be a base point of  $G$ . A (*geodesic*) *k-combing* is a choice of a shortest path  $P_{(b,x)}$  between  $b$  and each vertex  $x$ , such that  $P_{(b,v)}$  and  $P_{(b,w)}$  are *k-fellow travelers* for any adjacent vertices  $v$  and  $w$ . One can imagine the union of combing paths as a spanning tree  $T_b$  of  $G$  rooted at  $b$  and preserving the distances from  $b$  to all vertices. The neighbor of  $v$  in the unique path of  $T_b$  connecting  $v$  with the root will be called the *father* of  $v$ .

A (*geodesic*) *k-bicombing* is a choice of a shortest path  $P_{(v,v')}$  between each pair of vertices  $v, v'$ , such that  $P_{(v,v')}$  and  $P_{(w,w')}$  are *k-fellow travelers* for any vertices  $v, v', w, w'$  such that  $d_G(v, w) \leq 1$  and  $d_G(v', w') \leq 1$ . A particularly strong form of bicombing is the case in which the paths  $P_{(v,v')}$  and  $P_{(v',v)}$  are reverse. In this case we say that the bicombing is *symmetric*.

### 9.1. Combing

Given a base point  $b$ , a natural way to comb a graph  $G$  is to find a total order  $<_b$  refining the base point order  $\leq_b$ . Let  $\alpha$  be a strictly isotone map from  $V(G)$  to  $\mathbb{N}$ ; i.e.,  $\alpha(v) < \alpha(w)$  if and only if  $v <_b w$ . Denote by  $G_n$  the subgraph of  $G$  induced by vertices  $v$  with  $\alpha(v) \leq n$ . The *father* of a vertex  $v \neq b$  is the neighbor  $v_b$  of  $v$  with the least value  $\alpha(v_b)$ . Clearly,  $d_G(b, v_b) < d_G(b, v)$ . Define the combing path  $P_{(b,v)}$  to be the union of the path  $P_{(b,v_b)}$  and the edge  $vv_b$ .

Perhaps the simplest algorithmic way to define  $<_b$  is provided by the well known *breadth-first search* (BFS) procedure. In the breadth-first search the vertices of  $G$  are numbered in increasing order. We number with 1 the vertex  $b$  and put it on an initially empty queue of vertices. We repeatedly remove the vertex  $v$  at the head of the queue and consequently number and place onto the queue all still unnumbered neighbors of  $v$ . BFS constructs a spanning tree  $T_b$  of  $G$  with the vertex  $b$  as a root. A vertex  $x$  is the father

of any of its neighbors in  $G$  included in the queue when  $x$  is removed (a detailed description of the BFS procedure is presented, for example, in the book of Golombic [32]). In [24] we have investigated the properties of BFS orderings of weakly modular graphs. In particular, we have shown that for all house-free weakly modular graphs this ordering is distance preserving; i.e., each  $G_n$  is an isometric subgraph of  $G$ . In [23] we established that any ordering of a bridged graph produced by BFS is a domination ordering. We continue this line of research here by showing that BFS can be used for combing hereditary weakly modular graphs and graphs with convex balls (for an illustration see the first graph from Fig. 4). A part of the next result is implicit in [24].

**PROPOSITION 9.1.** *Let  $G$  be a hereditary weakly modular graph and  $b$  a base point. Then the paths  $\{P_{(b,v)} : v \in V(G)\}$  constructed by BFS form a geodesic 2-combing of  $G$ . If  $G$  is bridged, then this combing satisfies the 1-fellow traveler property.*

*Proof.* Let  $v$  and  $w$  be two adjacent vertices of  $G$  with  $v <_b w$ , and let  $v_b$  and  $w_b$  be their fathers. First assume that  $d_G(v, b) = d_G(w, b) = k$ . To prove that the paths  $P_{(b,v)}$  and  $P_{(b,w)}$  are 2-fellow travelers it suffices to establish that  $v_b$  and  $w_b$  are adjacent or coincide and that  $w_b$  and  $v$  are adjacent. We proceed by induction on  $k$ . If  $k = 1$ , then  $v_b = b = w_b$  and we are done. So, let  $k > 1$ . Suppose, for a contradiction, that  $d_G(v_b, w_b) > 1$ . Since  $v <_b w$ , according to BFS  $v_b <_b w_b$ . Thus  $v_b$  and  $w$  must be non-adjacent.

First let  $d(v_b, w_b) = 3$ . Since  $d_G(v_b, b) = d_G(w_b, b)$ , each pseudo-median of  $b, v_b, w_b$  has size 3 or size 1. In the first case it consists of  $v_b, w_b$  and a vertex  $x$  at distance 3 from  $v_b$  and  $w_b$ . From the properties of weakly modular graphs presented in Section 4 we deduce that  $d_G(v, x) = d_G(w, x) = 3$ . Since  $d_G(x, b) = k - 4$ , we get a contradiction with  $d_G(b, v) = d_G(b, w) = k$ . So, assume that the pseudo-median  $y, z, x$  of  $v_b, w_b, b$  has size 1. Then  $d_G(y, b) = d_G(z, b) = k - 2$  and




FIG. 4. BFS-combing and a partial bicombing of bridged graphs.

$d_G(x, b) = k - 3$ . Since  $v, y \in I(v_b, w_b)$ , by the quadrangle condition there is a common neighbor  $s$  of  $v, y$ , and  $w_b$ . As  $G$  is house-free,  $s$  and  $z$  must be adjacent. But then the vertices  $v_b, v, y, s, z$  induce a house, because  $d_G(v_b, w_b) = 3$  and  $d_G(y, b) = d_G(z, b) = k - 3$ .

Now, assume  $d_G(v_b, w_b) = 2$ . By the triangle condition applied to the vertices  $v_b, w_b, w$ , we can find their common neighbor  $z$ . In order to avoid an induced house, the vertices  $z$  and  $v$  must be adjacent. According to BFS  $\alpha(z) > \alpha(w_b)$ . We assert that  $d_G(z, b) = k - 1$ . Indeed, otherwise  $d_G(z, b) = k$  and  $v_b, w_b \in I(z, b)$ . By the quadrangle condition there is a common neighbor  $r$  of  $v_b$  and  $w_b$  at distance  $k - 2$  from  $b$ . Then we get two houses, induced by the vertices  $v, w, v_b, w_b, z, r$ . So, let  $d_G(z, b) = k - 1$ . Consider the fathers  $x, z_b, y$  of the vertices  $v_b, z, w_b$ , respectively. By the induction assumption  $d_G(x, z_b) \leq 1$  and  $d_G(z_b, y) \leq 1$ . In addition,  $z_b$  must be adjacent to both  $v_b$  and  $w_b$ , because  $\alpha(v_b) < \alpha(z) > \alpha(w_b)$ . Then, however, the vertices  $z_b, v_b, v, w, w_b$  induce a 5-cycle, which is impossible. So,  $d_G(v_b, w_b) \leq 1$ .

It remains to prove that if  $v_b \neq w_b$ , then  $w_b$  and  $v$  are adjacent. Suppose the contrary. Since  $v_b$  and  $w_b$  are at distance  $k - 1$  to  $b$ , by the triangle condition there is a common neighbor  $b'$  of  $v_b$  and  $w_b$  at distance  $k - 2$  to  $b$ . As a result we get a house, induced by  $v, w, v_b, w_b, b'$ .

To complete the proof, consider the second case  $k = d_G(b, v) < d_G(b, w)$ . By induction on  $k$  we show that  $v_b$  and  $w_b$  are adjacent. We may suppose that  $w_b$  is not adjacent to  $v$ ; otherwise from the preceding case we are done. Let  $y$  be the father of  $w_b$ . Necessarily  $y$  is not adjacent to  $v$ ; otherwise we are in contradiction with BFS. By the quadrangle condition we can find a common neighbor  $z$  of  $v$  and  $w_b$  at distance  $k - 1$  to  $b$ . Since  $\alpha(v_b) < \alpha(z) > \alpha(w_b)$ , by the induction assumption,  $v_b$  and  $y$  must be adjacent to the father  $z_b$  of  $z$ . The 6-cycle  $(v, w, w_b, y, z_b, v_b, v)$  must contain at least one diagonal; otherwise by Lemma 10 of [24] there is a vertex adjacent to all vertices of this cycle, contrary to the fact that  $d_G(v, z_b) = 3$ . But  $z_b$  is not adjacent to any of  $v, w_b, w$ , because  $d_G(z_b, b) = k - 2$ . Before we noticed that  $v$  is not adjacent to  $w_b, y$ . Therefore  $v_b$  and  $y$  are adjacent. To avoid an induced 5-cycle, the vertices  $v_b$  and  $w_b$  must be adjacent, as desired.

If  $G$  is bridged and  $k = d_G(b, v) < d_G(b, w)$ , then necessarily  $w_b$  is adjacent to  $v$ , reducing the second case to the first one, thus establishing the 1-fellow traveler property. ■

The  $t$ th level of an interval  $I(b, v)$  consists of all vertices of  $I(b, v)$  at distance  $t$  to  $b$ . For simplicity, let  $v_t = P_{(b,v)}(t)$ ,  $w_t = P_{(b,v)}(t)$ . The following simple property of BFS-orderings is essential.

*Remark 9.2.* (1)  $v_t$  is the vertex of the  $t$ th level of  $I(b, v)$  with the minimum value of  $\alpha$ ;

(2) If  $d_G(b, v) = d_G(b, w) = n$  and  $t \in [0, n]^*$ , then  $v \prec_b w$  if and only if  $v_t \prec_b w_t$ .

**PROPOSITION 9.3.** *Let  $G$  be a graph with convex balls and  $b$  a base point. Then the paths  $\{P_{(b,v)} : v \in V(G)\}$  constructed by BFS form a geodesic 2-combing of  $G$ .*

*Proof.* Pick two adjacent vertices  $v$  and  $w$  of  $G$  such that  $v \prec_b w$ ,  $d_G(b, v) = n$  and  $v_b \neq w_b$ . If  $d_G(b, v) < d_G(b, w)$ , from the convexity of  $B_n(b)$  we infer that  $v$  and  $w_b$  are adjacent. Thus assume that  $d_G(b, w) = n$ . Let  $b'$  be the lowest common ancestor of  $v$  and  $w$  in the tree  $T_b$ , i.e., the furthest from the  $b$  common vertex of the combing paths  $P_{(b,v)}$  and  $P_{(b,w)}$ . Consider the cycle  $C$  formed by the edge  $vw$  and the subpaths of  $P_{(b,v)}$  and  $P_{(b,w)}$  between  $b'$  and  $v$  and  $w$ , respectively. If  $C$  is a 3- or a 5-cycle, then clearly  $d_G(v_t, w_t) \leq 2$  for any nonnegative integer  $t \leq n$ . Otherwise, since  $C$  is well bridged either  $v$  and  $v_b$  are adjacent to  $w_b$ , or there is a bridge from  $w$  to a vertex  $x \neq b'$  on the shortest path of  $C$  between  $v$  and  $b'$ . Since  $d_G(v, b) = d_G(w, b)$ , we obtain that  $d_G(v, x) = d_G(w, x)$ , whence  $x \in I(b, w)$ . Let  $y$  be the vertex on the shortest path of  $C$  between  $w$  and  $b'$  with  $d_G(y, b') = d_G(x, b')$ . Since  $x$  and  $y$  have the same distance to  $b$ , Remark 9.2(1) implies  $y \prec_b x$ . But then from Remark 9.2(2) we conclude that  $w \prec_b v$ , which is impossible. ■

To find a combing of a Helly graph  $G$  we define the rooted tree  $T_b$  as follows. Given a vertex  $v$  at distance  $n$  to  $b$ , there exists a neighbor  $v_b$  of  $v$  at distance  $n - 1$  to  $b$  which is adjacent to all neighbors  $x$  of  $v$  with  $d_G(b, x) \leq n$  (in fact, this property characterizes the Helly graphs [12]). Indeed, take the collection of pairwise intersecting balls consisting of  $B_{n-1}(b)$ ,  $B_1(v)$ , and  $B_1(x)$  for all neighbors  $x$  of  $v$  with  $d_G(b, x) \leq n$ . Then we can select any point from their intersection as the father  $v_b$  of  $v$ . Pick two adjacent vertices  $v, w$ . If  $d_G(b, v) = d_G(b, w)$ , then either their fathers coincide or  $v, w, v_b, w_b$  are pairwise adjacent. Otherwise, if  $d_G(b, w) > d_G(b, v)$  and  $w_b \neq v$ , necessarily  $w_b$  is adjacent to  $v$  and  $v_b$ , thus establishing the following observation.

**PROPOSITION 9.4.** *Every Helly graph  $G$  has a geodesic 1-combing.*

We continue with combings of hypercubes and median graphs. Notice that one cannot depart from an arbitrary BFS ordering, even if we want to comb the 3-cube. Let  $G$  be a median graph with a base point  $b$  isometrically embedded into the hypercube  $H(\Lambda)$  as was presented in Section 6. To comb  $H(\Lambda)$ , we define  $\prec_b$  as the lexicographic order on the finite subsets of the totally ordered set  $\Lambda$  ( $\prec_b$  can be obtained by a stronger version of BFS, namely, by the lexicographic breadth-first search). The father of a vertex  $A$  in the combing tree  $T_b$  is the vertex  $A - \{\lambda\}$ , where  $\lambda$  is the largest

element in  $A$ . If  $B$  is a neighbor of  $A$ , say  $B = A - \{\mu\}$ , then  $A - \{\lambda, \mu\}$  is the father of  $B$ , showing that  $T_b$  provides a geodesic 2-combing of  $H(\Lambda)$ .

Since  $G$  is a retract of  $H(\Lambda)$ ,  $G$  has a geodesic 2-combing as well. In fact, we can provide an explicit 2-combing of  $G$ . For this, let  $\Lambda_n = \{i \in \Lambda : i \leq n\}$  and let  $G(n)$  be the subgraph induced by the vertices of  $G$  contained in  $\Lambda_n$ . Clearly  $G(1) \subseteq G(2) \subseteq \dots$  and  $G = \bigcup_{n \in \Lambda} G(n)$ . Every  $G(n)$ , being the intersection of  $G$  with the hypercube  $H(\Lambda_n)$ , is a convex subgraph of  $G$ . The equivalence class  $E_i$  splits  $G$  into halfspaces  $G'_{n+1}$  and  $G''_{n+1}$ , where  $G(n)$  is contained in one of them, say  $G(n) \subseteq G'_{n+1}$ . One can easily show that the complement  $H''$  of  $G(n)$  in  $G(n+1)$  consists of all vertices  $v''$  of  $H''_{n+1}$  whose neighbors  $v'$  in  $H'_{n+1}$  belong to  $G(n)$ . (In other words,  $H''$  is a thin halfspace of  $G(n+1)$ .) Let  $H'$  denote the subgraph of  $G(n)$  induced by the vertices  $v'$ . Notice that both  $H', H''$  are convex subgraphs of the graphs  $G(n+1)$  and  $G(n)$  (see Section 6.2). The vertex  $v'$  will be called the father of  $v''$ .

Assume that we have defined the combing tree  $T_b^+(n)$  in the graph  $G(n)$  and we wish to extend it to the tree  $T_b^+(n+1)$  of  $G(n+1)$ . For this, add to  $T_b^+(n)$  all edges of the type  $v'v''$ , thus  $v'$  becomes the unique neighbor of  $v''$  in  $T_b^+(n+1)$ . If  $v'', w'' \in H''$  are adjacent, then by the convexity of  $G(n)$  we infer that their fathers are adjacent as well. This shows that the resulting tree  $T_b^+ = \bigcup_{n \in \Lambda} T_b^+(n)$  provides a geodesic 2-combing for  $G$ .

There is yet another way to grow a combing tree  $T_b^-$  of  $G$ . Namely, given the current tree  $T_b^-(n)$ , we intersect it with  $H'$  and consider the connected components of this intersection. In each such a component  $C$  we pick the vertex  $x_C$  closest to the root  $b$ . (In fact, one can view  $C$  as a tree rooted at  $x_C$ .) In  $H''$  take the copy of each  $C$ , add it to  $T_b^-(n+1)$ , previously connecting it with  $C$  by the edge between  $x_C$  and its neighbor in  $H''$ . Set  $T_b^- = \bigcap_{n \geq 1} T_b^-(n)$ . Before establishing the 2-fellow property we pause to emphasize two properties of the algorithm. First, notice that there are no 4-cycles of  $G$  in which exactly two incident edges belong to  $T_b^-$ . Second, if a 4-cycle  $C$  shares three edges with  $T_b^-$ , then the fourth edge belongs to an equivalence class with a smaller number than that of the equivalence class which contains the two opposite edges from  $C \cap T_b^-$ .

Let  $v, w$  be adjacent vertices of  $G(n+1)$ . Consider the paths  $P_{(b,v)}$  and  $P_{(b,w)}$  of  $T_b^-$  from  $b$  to  $v$  and  $w$ . To check that  $P_{(b,v)}$  and  $P_{(b,w)}$  are 2-fellow travelers, by induction on  $n$  we prove the following stronger condition, which we call the “1.5-fellow traveler property”:

each vertex of  $P_{(b,v)}$  is either adjacent to a vertex of  $P_{(b,w)}$  or its neighbors in  $P_{(b,v)}$  are adjacent to a common vertex in  $P_{(b,w)}$  and vice versa.

The result easily follows if  $v \in H'$  and  $w \in H''$ . So suppose that  $v, w \in H''$ , say  $v = v', w = w'$ . If  $v'$  and  $w'$  belong to a common connected component of  $T_b^-(n) \cap H'$ , or they belong to different components  $C', C''$ , but

neither  $v'$  nor  $w'$  are the roots, the result easily follows by the induction hypothesis and the algorithm. Finally, suppose that  $v' \in C'$  is the root of  $C'$ , while the root of the connected component  $C''$  containing  $w'$  is a vertex  $x' \neq w'$ . According to the procedure used,  $P_{(b,v)}$  consists of  $P_{(b,v')}$  and the edge  $v'v$ . The path  $P_{(b,w')}$  consists of  $P_{(b,x')}$ , the edge  $x'x''$ , and the image in  $H''$  of the path  $P_{(x',w')}$ . Notice that  $P_{(b,w')}$  consists of  $P_{(b,x')}$  and  $P_{(x',w')}$ . Let  $z'$  be the neighbor of  $w'$  in  $P_{(x',w')}$  and let  $y, u$  be the last two vertices in  $P_{(b,v')}$ . By the induction assumption, either  $u$  is adjacent to  $z'$  or  $w'$  and  $y$  are adjacent. In the first case  $u \in I(v', z') \subseteq H'$ , because  $H'$  is convex. Then  $u \in C'$ , contrary to the assumption that  $v'$  is the closest to  $b$  vertex of  $C'$ . Otherwise, if  $w'$  is adjacent to  $y$ , then we get a 4-cycle  $(y, u, v', w', y)$  which contradicts the properties established before.

For an illustration of both methods, see Fig. 5. In this figure, the median graph is isometrically embedded in  $\mathbb{E}^3$  endowed with the  $l_1$ -metric. The equivalence classes are performed in the order indicated by the labels of edges in the tree.

Slightly modifying the first procedure, one can obtain a geodesic 1-combing of the associated to  $G$  Helly graph  $G^\Delta$ . For this, notice that a cube  $C$  of  $G$  intersects every  $G(n)$  in a subcube. Therefore, if  $C$  intersects both  $G(n)$  and  $H'' = G(n+1) - G(n)$ , then it shares with each of them a facet. Let  $T_b^\Delta = \bigcup_{n \in \Lambda} T_b^\Delta(n)$ ; we extend the combing tree  $T_b^\Delta(n)$  in the following way. For a new vertex  $v''$  we consider the neighbor  $x$  of  $v'$  in the current tree  $T_b^\Delta(n)$ . If  $xv''$  spans a cube, then add the edge  $v''x$  to  $T_b^\Delta(n+1)$ . Otherwise add the edge  $v'v''$  to  $T_b^\Delta(n+1)$ . By induction on  $n$  we show that the path  $P_{(b,v'')}$  of  $T_b^\Delta(n+1)$  connecting  $b$  and  $v''$  is a normal cube-path in  $G(n+1)$  (and therefore in  $G$ ). Let  $C, C'$  be the cubes spanned by the first two edges of  $P_{(b,v'')}$ . We must show that  $C \cap \text{star}(C', \mathcal{X})$  consists of one vertex, namely the neighbor of  $v''$  in




FIG. 5. Combing trees  $T_b^+$  and  $T_b^-$  of a median graph.

$T_b^\Delta(n+1)$ . If  $v''$  and  $v'$  have a common neighbor  $x$  in  $T_b^\Delta(n+1)$ , then a cube  $Q$  from  $\text{star}(C', \mathcal{X})$  intersects the facet of  $C$  from  $H''$  if and only if it intersects the complementary facet spanned by the edge  $v'x$ , contrary to the induction assumption. Now, let  $v'$  be the neighbor of  $v''$  in  $T_b^\Delta(n+1)$ . Then  $C$  consists only of  $v', v''$ . If  $v'' \in Q$ , then  $v''$  and the neighbor  $x$  of  $v'$  both belong to  $Q$ , implying that actually  $x$  must be the predecessor of  $v''$ . The 1-fellow traveler property easily follows by induction. Since every normal cube path is a geodesic of  $G^\Delta$ , we recover the same combing as that of Niblo and Reeves [44]. Concluding, we have proven the following result.

**PROPOSITION 9.5.** *If  $G$  is a locally finite median graph then  $G$  and  $G^\Delta$  have geodesic 2-combings and 1-combings defined by the trees  $T_b^+$ ,  $T_b^-$ , and  $T_b^\Delta$ .*

## 9.2. Bicombings

Finding a bicombing of a graph seems to be more difficult than finding a combing. For example, Gersten and Short [29, 30] use a special structure of minimal disk diagrams and Noskov [45] presented a bicombing of triangle buildings based on geometric properties of the tessellation of the plane into equilateral triangles. We continue with algorithms for bicombing bridged graphs and F-graphs. Using some results from [24] we can extend them to all hereditary weakly modular graphs and graphs with convex disks, but the correction proofs become more technical and will be presented elsewhere. Notice that one cannot extend these results to all graphs in which the isometric cycles have length three or four. For this take the square grid  $n \times n$ , delete each of four corners, and make its neighbors in the grid adjacent. We will obtain an octagon with four sides of length 1 and four sides of length  $n-1$  subdivided into  $n^2-4$  squares and 4 triangles. The resulting graph  $\Gamma$  does not contain other isometric cycles. Let  $vw$  and  $v'w'$  be two opposite short sides of the octagon. Then  $d_\Gamma(v, v') = d_\Gamma(w, w') = 2n-3$ . The pairs  $v, v'$  and  $w, w'$  are connected by unique shortest paths which are not  $k$ -fellow travelers for any  $k < n$ . One can obtain the same effect if one triangulates the previous graph  $\Gamma$  by adding in each square the diagonal parallel to  $vw$ . In the resulting graph all isometric cycles have length three, except two cycles which have length five.

We analyze a few simple ways of generating a geodesic bicombing in a graph. First, we can fix a base point  $b$  and an ordering  $<_b$  refining  $\leq_b$  (in our cases this will be an BFS ordering). Suppose that every  $G_k$  is an isometric subgraph of  $G$  (this is the case of graphs occurring in this section; see [24]). Assume that we have defined bicombing paths between each pair of vertices of  $G_{k-1}$  and we wish to extend this bicombing further. Let  $\alpha(v) = k$ . For any vertex  $v'$  with  $\alpha(v') < k$ , let  $P_{(v', v)}$  be the union of the path  $P_{(v', v^*)}$  and the edge  $v^*v$ , where  $v^*$  is the neighbor of  $v$  in the interval  $I(v, v')$

with the smallest  $\alpha(v^*)$ . Clearly, if the father  $v_b$  of  $v$  belongs to  $I(v', v)$  then  $v^*$  coincides with  $v_b$ . If the neighbor  $x$  of  $v_b$  in  $P_{(v', v_b)}$  is adjacent to  $v$ , then  $v^* = x$ . The resulting path  $P_{(v', v)}$  is unimodal with respect to the function  $\alpha$ . Namely, when moving from  $v$  to  $v'$  along  $P_{(v', v)}$ ,  $\alpha$  decreases until we arrive at the vertex where it achieves the minimum value and then it strictly increases until we reach  $v'$ . As a result, we obtain a symmetric geodesic bicombing. (In Fig. 4 we present certain paths in a bicombing of a bridged graph obtained by this procedure. We do not list the complete BFS ordering, only the labels of the neighbors of the root-vertex  $b$ .)

If we let  $P'_{(v', v)}$  be the union of the path  $P'_{(v', v_b)}$  and the edge  $v_b v$ , we obtain a symmetric bicombing, but the bicombing paths are not necessarily geodesics. If  $v_b$  dominates the vertex  $v$  in  $G_k$  (this means that every neighbor of  $v$  in  $G_k$  is a neighbor of  $v_b$ , too), then one can easily show by induction that  $\{P'_{(v', v)} : v', v \in V(G)\}$  constitutes a 1-bicombing. Such graphs are known in the literature under the name *dismantlable graphs*. It is known and it can be easily shown that Helly graphs are dismantlable [12]. That bridged graphs are dismantlable has been established in [3]. In [23] it is shown that this can be done by BFS. In this case a vertex  $v$  with  $\alpha(v) = k$  is dominated in  $G_k$  by its father in the BFS-tree.

**PROPOSITION 9.6.** *Let  $G$  be a locally finite bridged graph. Then the symmetric geodesic bicombing  $\{P_{(v', v)} : v', v \in V(G)\}$  satisfies the 2-fellow traveler property.*

*Proof.* Let  $\alpha(v) = k$ . By induction on  $n$  we show that if  $d_G(v, w) \leq 1$ ,  $d_G(v', w') \leq 1$  and all  $v', w', w, v'$  belong to  $G_k$ , then  $P_{(v', v)} = (v' = v_0, v_1, \dots, v_{n-1}, v_n = v)$  and  $P_{(w, w')} = (w' = w_0, w_1, \dots, w_{m-1}, w_m = w)$  lie in a 1-Hausdorff neighborhood of each other.

If  $v$  is adjacent to some vertex  $w_i$  or  $w$  is adjacent to a vertex  $v_i$ , then we can apply the induction hypothesis. The same applies if the vertices  $v_{n-1}$  and  $w_{m-1}$  are adjacent. Thus we may assume the converse. Let  $v_b$  be the father of  $v$  and denote by  $x$  the neighbor of  $v_b$  in the path  $P_{(v', v_b)}$ . Since  $v_b$  dominates  $v$ ,  $v_b$  is adjacent to  $w$  and  $v_{n-1}$  (from our assumption, clearly  $v_b \neq v_{n-1}$ ).

*Case 1.*  $x = v_{n-1}$ .

By the induction assumption, the paths  $P_{(w', w)}$  and  $P_{(v', v_b)}$  lie in a 1-Hausdorff neighborhood of each other. Then we obtain the desired property, unless some vertex  $w_i$  is adjacent in  $P_{(v', v_b)}$  only to  $v_b$ . Clearly  $i \geq m - 2$ . To avoid induced 4-cycles,  $v_b$  and  $w_{m-1}$  must be adjacent. If  $w_{m-1}$  is adjacent to some  $v_j$ , then  $j \geq n - 3$ . To avoid induced 5-cycles or 4-cycles, the vertices  $w_{m-1}$  and  $v_{n-1}$  must be adjacent, which is impossible. Therefore  $i = n - 1$ . By the induction assumption,  $v_{n-1}$  is adjacent to a vertex  $w_l$ .




FIG. 6.

Since  $P_{(w,w')}$  is a shortest path, necessarily  $l = m - 2$  or  $m - 3$ . If  $v_{n-1}$  is adjacent to  $w_{m-3}$  but not to  $w_{m-2}$ , then the vertices  $v_b, v_{n-1}, w_{m-3}, w_{m-2}, w_m$  induce a 5-cycle, or  $v_b$  and  $w_{m-3}$  are adjacent, and we get a 4-cycle formed by  $v_b, v_{n-1}, w_{m-3}, w_{m-2}$ . Thus  $v_{n-1}$  and  $w_{m-2}$  are adjacent. But then we have obtained an induced 5-cycle  $(w, v, v_{n-1}, w_{m-2}, w_{m-1}, w)$ . This settles the case  $x = v_{n-1}$ ; for an illustration see Fig. 6.

Case 2.  $x \neq v_{n-1}$ .

This implies that  $x, v_{n-1} \in I(v_b, v')$ , therefore  $x$  and  $v_{n-1}$  are adjacent. On the other hand,  $x$  is not adjacent to  $v$ . Applying the induction assumption to the paths  $P_{(w',w)}$  and  $P_{(v',v_b)}$ , we conclude that  $x$  is adjacent to  $w$  or to a vertex  $w_i, i \geq m - 3$ . If  $w$  and  $x$  are adjacent, then the vertices  $w, v, v_{n-1}, x$  form an induced 4-cycle. A forbidden 5-cycle  $(w, v, v_{n-1}, x, w_{m-1}, w)$  occurs if  $x$  and  $w_{m-1}$  are adjacent. If  $x$  is adjacent to  $w_{m-3}$  then  $w_{m-2}, x \in I(w_{m-3}, w)$ ; therefore  $x$  and  $w_{m-2}$  must be adjacent. Hence  $i = m - 2$ . To avoid forbidden cycles induced by  $w, v_b, x, w_{m-2}, w_{m-1}$ , the vertex  $v_b$  must be adjacent to both  $w_{m-1}, w_{m-2}$ ; see Fig. 6. This implies  $v_b \in I(w, w')$ , whence  $P = (v_b, w_{m-2}, w_{m-3}, \dots, w_1, w')$  is a shortest path. On the other hand, we know that  $Q = (v_b, v_{n-1}, v_{n-2}, \dots, v_1, v')$  is also a shortest path. Consider the cycle formed by the paths  $P, Q$  and the edge  $v'w'$ . It cannot be well bridged, because the neighbors  $v_{n-1}$  and  $w_{m-2}$  of  $v_b$  in this cycle are not adjacent, thus settling Case 2.

This concludes the proof that  $P_{(v',v)}$  and  $P_{(w',w)}$  lie in a 1-Hausdorff neighborhood of each other. To prove that these paths are 2-fellow travelers we must show that a vertex  $v_i$  is adjacent to at least one of the vertices  $w_{i-1}, w_i, w_{i+1}$ . Suppose that  $v_i$  is adjacent to a vertex  $w_j$ . Since

$P_{(v',v)}$  and  $P_{(w',w)}$  are shortest paths, we have  $|i - j| \leq 2$ . If  $j = i + 2$ , then  $v_i, w_{i+1} \in I(w_{i+2}, w')$ ; thus  $v_i$  and  $w_{i+1}$  are adjacent, as desired. Otherwise, if  $j = i - 2$ , then  $d_G(w_{i-2}, w) \geq d_G(v_i, v)$  and  $d_G(w_{i-2}, v) > d_G(v_i, v)$ . Consider the cycle  $C$  formed by the edges  $vw, v_iw_{i-2}$  and the portions of the paths  $P_{(v',v)}$  and  $P_{(w',w)}$  comprised between  $v, v_i$  and  $w, w_{i-2}$ , respectively. Since  $C$  is well bridged, from the previous inequalities we deduce that the neighbors  $v_i$  and  $w_{i-1}$  of  $w_{i-2}$  in  $C$  are adjacent, concluding the proof. ■

*Remark 9.7.* From the proof one can conclude that if  $|d_G(v, v') - d_G(w, w')| \leq 1$ , then the paths  $P_{(v',v)}, P_{(w',w)}$  are 1-fellow travelers (this is also true for bicomblings of F-graphs). We do not have an example of a bridged graph without a geodesic 1-bicombling. However, one can construct bridged graphs and BFS-bicomblings which do not satisfy the 1-fellow traveler property.

To find a bicombling of an F-graph  $G$  we use a slightly modified version of the previous procedure. It concerns the case when  $v$  belongs to a pendant folder  $F$  of  $G_k$ , where  $\alpha(v) = k$ . For a vertex  $v' \prec_b v$ , let  $v^*$  be the neighbor of  $v$  in  $I(v, v')$  with the value of  $\alpha$ . If  $v^b \notin I(v, v')$ , then let  $P_{(v',v)}$  be the union of the path  $P_{(v',v^*)}$  and the edge  $v^*v$ , otherwise; if  $v^b \in I(v, v')$ , then the combling path  $P_{(v',v)}$  is the union of the paths  $P_{(v',v^b)}$  and  $(v^b, v^*, v)$ . Before presenting its analysis, we formulate the following property of BFS orderings of  $G$ .

**LEMMA 9.8.** *Let  $G$  be a locally finite F-graph. If  $x, y$  are two opposite vertices of a folder  $F$  and  $x \prec_b y$ , then the father of  $y$  in the BFS ordering belongs to  $F$ .*

*Proof.* Suppose  $y_b \notin F$ . Then  $x_b \notin F$ ; otherwise  $y_b = x_b$  by BFS. Pick a vertex  $z \in F$  adjacent to both  $x$  and  $y$ , such that  $z \prec_b y$ . By Proposition 9.1 its father  $z_b$  is adjacent to  $y$ . If  $z_b$  is distinct from  $x$ , then by Proposition 9.1  $z_b$  is adjacent to  $x_b$  or  $z$  is adjacent to the father of  $x_b$ . In the first case we obtain a forbidden configuration of three folders. In the second case we obtain  $z \prec_b y_b$ , which is impossible. So suppose that  $z_b = x$ . By Proposition 9.1 either  $x$  and  $y_b$  are adjacent, implying  $y_b \in F$ , or  $z$  is adjacent to the father  $s$  of  $y_b$ . Then  $s$  must be adjacent to  $x_b$ , yielding an induced 6-cycle. ■

**PROPOSITION 9.9.** *Let  $G$  be a locally finite F-graph. Then the symmetric geodesic bicombling  $\{P_{(v',v)} : v', v \in V(G)\}$  satisfies the 2-fellow traveler property.*

*Proof.* Let  $\alpha(v) = k$ ,  $d_G(v, w) \leq 1, d_G(v', w') \leq 1$ , and suppose that  $v', w', w, v'$  belong to  $G_k$ . Denote  $P_{(v',v)} = (v' = v_0, v_1, \dots, v_{n-1}, v_n = v)$  and  $P_{(w',w)} = (w' = w_0, w_1, \dots, w_{m-1}, w_m = w)$ . By induction on  $n + m$  we

show that every  $v_i$  either is adjacent to a vertex of  $P_{(w',w)}$  or both its neighbors  $v_{i-1}, v_{i+1}$  have a common neighbor in  $P_{(w',w)}$  and vice versa. From this we immediately obtain the 2-fellow traveler property. We know that in  $G_k$  the vertex  $v$  either is pendant or belongs to a pendant folder  $F$ . The either case is trivial, because then  $v_{n-1} = w$ . Thus let  $v, w$  belong to a common folder  $F$ . Without loss of generality we may assume that  $v', w' \notin F$  and  $v \neq w$ . Recall that by  $v^b \in F$  we denote the vertex of  $F$  opposite to  $v$ . From Lemma 9.8 we conclude that  $v^b$  is the father of at least one of the vertices  $v_{n-1}$  and  $w$ . If  $v^b = v_{n-2}$  then  $\alpha(w) > \alpha(v_{n-1})$  and  $v^b$  is the father of  $w$ . Then either  $w_{m-1} = v^b$  or  $d_G(v^b, v') = d_G(w, w')$ . In both cases we can apply the induction hypothesis. The same holds when  $v^b = w_{m-1}$ . So assume  $v^b \neq v_{n-2}, w_{m-1}$ . By Lemma 7.6 we conclude that if  $v'$  and  $w'$  have distinct gates in  $F$ , then  $v^b$  is the gate of at least one of them. From the algorithm and our agreement we deduce that  $v^b$  is the gate of  $w'$  and the father of  $v_{n-1}$ . Then  $w_{m-1} \prec_b v^b$  and  $v^b \notin I(v_{n-1}, v')$ . Since  $G$  is bipartite, this implies that  $d_G(v_{n-1}, v') = d_G(v^b, w')$ , and therefore  $d_G(v, v') = d_G(w, w')$ . Let  $u$  be the neighbor of  $v^b$  in the combing path  $P_{(w',v^b)}$ . By the induction assumption we infer that  $v_{n-2}$  and  $u$  are adjacent. If  $u$  is adjacent to  $w_{m-1}$ , we obtain three folders with a forbidden intersection. Thus  $v^b$  is adjacent to  $w_{m-2}$ . Since  $v^b \in I(w, w')$ , from the choice of  $w_{m-1}$  we deduce that  $w_{m-1} \prec_b v^b$ . By Lemma 9.8  $w_{m-2}$  must be the father of  $v^b$ . Since  $w_{m-2} \in I(v^b, w')$ , we obtain a contradiction with the choice of the vertex  $u$ . Finally, assume that  $v'$  and  $w'$  have the same gate in  $F$  different from  $v^b$ . Evidently, this gate is the vertex  $v_{n-1}$ . Then  $v, v^b, w_{m-1} \in I(w, w')$ , and by Lemma 7.4  $v_{n-1}$  and  $w_{m-1}$  are adjacent. Since  $\alpha(w_{m-1})$  and  $\alpha(v^b)$  are smaller than  $\alpha(w)$ , by the algorithm we have  $w_{m-2} = w^b$ . To avoid forbidden subgraphs induced by the vertices  $v, v_{n-1}, w, v^b, w_{m-1}, w^b$ , necessarily  $w^b = v_{n-1}$ . ■

Although there exist geodesic 1-bicomblings of  $G^\Delta$  derived by a procedure similar to those for bridged graphs or F-graphs, we present the analysis of the bicombing generated by normal folder-paths.

**PROPOSITION 9.10.** *Let  $G$  be a locally finite F-graph. Then the bicombing of  $G^\Delta$  generated by normal folder-paths satisfies the 1-fellow traveler property.*

*Proof.* As before, suppose that  $d_{G^\Delta}(v, w) \leq 1, d_{G^\Delta}(v', w') \leq 1$  and let  $P_{(v,v')} = (v = v_0, v_1, \dots, v_n = v')$ ,  $P_{(w,w')} = (w = w_0, w_1, \dots, w_m = w')$  be the normal folder-paths connecting  $v, v'$  and  $w, w'$ , respectively. Let  $F'$  and  $F''$  be the cells (folders or edges) spanned by  $v, w$  and  $v', w'$ , respectively. It is sufficient to show that  $v_1$  and  $w_2$  are adjacent in  $G^\Delta$ , and the general result follows by induction on  $n + m$  and the fact that a subpath of a normal folder-path is itself normal. This can be checked directly, using the properties of F-graphs. We will outline a shorter proof, using the result from [44] in Proposition 9.12 below.

Every folder-path lifts to a path of  $G$  obtained by replacing each pair  $v_i v_{i+1}$  spanning a folder  $F_i$  by a path consisting of  $v_i, v_{i+1}$  and their common neighbor in  $F_i$ . If we perform this operation with all folders in  $P_{(v,v')}$  and  $P_{(w,w')}$ , we obtain two genuine paths  $P', P''$  of  $G$ . These paths together with some paths connecting  $v, w$  and  $v', w'$  in  $F', F''$  form a cycle  $C$  of  $G$ . Assume that  $C$  is selected so that the area of a minimal singular disk quadrangulation  $(Q, f)$  of  $C$  is as small as possible. Pick a path of length two  $(v_i, a, v_{i+1}) \subset \partial Q$ . Its image is contained in the folder  $F_i$ . Add a new face  $q_i$  to  $Q$  by taking a vertex  $a'$  adjacent to  $v_i$  and  $v_{i+1}$ . Performing this operation with all folders from both normal folder-paths and with  $F', F''$  as well, we surround  $Q$  by a ring of new quadrangles, thus obtaining a new quadrangulation  $Q'$  (some care is needed while performing this operation for  $F'$  and  $F''$ ). Since two consecutive folders from one normal folder-path intersect exactly in one vertex, this implies that  $f$  can be extended to a cellular map from  $Q'$  to  $G$  (namely, it will map  $q_i$  to a 4-cycle of the folder  $F_i$ ). From Theorem 6.1 and the claim in the proof of Theorem 7.1 we infer that  $Q'$  is a cube-free median graph. The vertices  $v$  and  $v'$  are connected in  $Q'$  by a normal cube-path consisting of new faces and boundary edges mapped to the folders and edges of  $P_{(v,v')}$  (a normal cube-path for  $w, w'$  is defined similarly). By Proposition 9.12, the vertices  $v_1$  and  $w_1$  belong to a common face of  $Q'$ ; therefore they belong to a common folder in  $G$ . ■

Geodesic bicomblings of median graphs  $G$  and associated Helly graphs  $G^\Delta$  can be found by modifying the combing procedure from Section 9.2. Assume that we have defined the combing paths between any two vertices of the graph  $G(n)$  and we wish to extend this bicombing to the graph  $G(n+1)$ . We denote the union of the combing paths of  $G(n)$  pointing from a vertex  $u$  by  $T_u(n)$ . To extend this tree, if  $u \in G(n)$ , then we let a vertex  $v'' \in H''$  be adjacent in  $T_u(n+1)$  with its father  $v'$ . If  $u \in H''$ , say  $u = u''$ , then define  $T_{u''}(n+1)$  be the image in  $H''$  of the tree  $T_{u''}(n) \cap H'$  (that this indeed is a tree follows from the convexity of  $H'$ ). Put  $T_u = \bigcup_{n \geq 1} T_u(n)$ . Clearly, the obtained bicombing consists of shortest paths only. By induction on  $n$  it easily follows that if  $v, w \in T_u$  are adjacent in  $G$ , then the paths of  $T_u$  between  $u$  and  $v, w$  are “1.5-fellow travelers.” Using this fact, a simple case analysis, and the induction on  $n$ , one can prove the following result.

**PROPOSITION 9.11.** *If  $G$  is a locally finite median graph, then the geodesic bicombing defined by the collection of trees  $T_v, v \in V(G)$  satisfies the 3-fellow traveler property.*

We continue with the following result of Niblo and Reeves.

**PROPOSITION 9.12 [44].** *If  $G$  is a locally finite median graph, then the bicombing of  $G^\Delta$  generated by normal cube-paths satisfies the 1-fellow traveler property.*

Using the preprocessing of a median graph  $G$  defined before, we outline an algorithmic way to find the normal cube-paths between two vertices  $v, w$  of  $G$ . Suppose that this task is accomplished for  $G(n)$ . If both vertices are in  $H''$ , say they are  $v'', w''$ , then define the paths  $P_{(v'', w'')}$  and  $P_{(w'', v'')}$  as the images in  $H''$  of the paths  $P_{(v', w')}$  and  $P_{(w', v')}$ , respectively. Now, assume that one vertex  $v$  is in  $G(n)$  and another  $w = w''$  is in  $H''$ . Let  $x$  be the neighbor of  $w'$  in  $P_{(v, w')}$ . Again, if the pair  $xw''$  spans a cube, then set  $P_{(v, w'')} = P_{(v, x)} \cup \{xw''\}$ ; otherwise let  $P_{(v, w'')} = P_{(v, w')} \cup \{w'w''\}$ . The proof that the paths constructed this way are normal is the same as in Section 9.1. To find the path  $P_{(w'', v)}$  we use the fact that  $H'$  and  $H''$  are convex and gated. If  $v \in H'$  and  $y$  is the neighbor of  $w'$  in  $P_{(w', v)}$ , then one can show that  $y$  and  $w''$  span a cube. Then  $P_{(w'', v)}$  consists of the pair  $w''y$  and the path  $P_{(y, v)}$ . Otherwise, if  $v \notin H'$ , let  $v_{H'}$  be its gate in  $H'$ . Consider the neighbor  $z$  of  $w''$  in the path  $P_{(w'', v_{H'})}$ . Then set  $P_{(w'', v)}$  be the union of  $w''z$  and the path  $P_{(z, v)}$ .

Proposition 9.12 can be used to establish that Helly graphs have a geodesic 1-bicombing. Indeed, the basic example of a cubing is the Euclidean space tiled into regular cubes. The underlying graph is the rectilinear grid  $\Gamma$ , which can be viewed as a Cartesian product of paths. The associated Helly graph  $\Gamma^\Delta$  is the direct product of paths. By Theorem 4.7, Helly graphs are exactly the retracts of such products (maybe with an infinite number of factors). Therefore, at least for Helly graphs  $G$  isometrically embeddable into the direct product of a finite number of paths, the following holds.

**COROLLARY 9.13.** *Helly graphs have geodesic 1-bicombings.*

We were not successful in developing an algorithm for finding such a bicombing, and leave this as a question.

*Note.* After the submission of the manuscript we learned about two papers closely related with the subject of our paper. First, G. Noskov informed us about the preprint of M. Roller (1998, “Pocsets, Median Algebras and Group Actions. An Extended Study of Dunwoody’s Construction and Sageev’s Theorem,” University of Southampton, Pure Mathematics Preprints and Reports, No. 8, 105 pp.). Based on the results of M. Sageev, M. Roller also established the relationship between cubings and median algebras, and investigated the properties of groups acting on discrete median algebras. He also mentioned V. Gerasimov for a similar result. Second, the paper of J. M. Corson and B. Trace (1998, “The 6-Property for Simplicial Complexes and a Combinatorial Cartan–Hadamard Theorem for Manifolds,” Proc. Amer. Math. Soc. Vol. 126) establishes that every locally finite, simply connected, two-dimensional simplicial complex  $K$  in which the links of vertices do not contain  $n$ -cycles for  $n \leq 5$  can be represented as a monotone union of a sequence of collapsible subcomplexes  $K_i$  (in particular, if

$K$  is finite, then it is collapsible). Clearly, the complexes considered by Corson and Trace constitute a subclass of bridged complexes. Dismantlability of the underlying graphs of bridged complexes yields their collapsibility in a somewhat stronger form. The result of Corson and Trace can be refined by requiring that for every  $i > 1$  the subcomplex  $K_{i+1}$  contains one vertex more than  $K_i$  does.

## ACKNOWLEDGMENTS

I express my appreciation to Walter Deuber for his support. I gratefully acknowledge the stimulating mathematical environment in Bielefeld, where this work was done. Finally, I acknowledge Alexander Karzanov for a careful reading of the first version and critical remarks improving the presentation.

## REFERENCES

1. A. D. Alexandrov and V. A. Zalgaller, "Intrinsic Geometry of Surfaces," Transl. Math. Monographs 15, Am. Math. Soc., Providence, 1967.
2. A. D. Alexandrov, V. N. Berestovskii, and I. G. Nikolaev, Generalized Riemannian spaces, *Russian Math. Surveys* **41** (1986), 1–54.
3. R. P. Anstee and M. Farber, On bridged graphs and cop-win graphs, *J. Combin. Theory Ser. B* **44** (1988), 22–28.
4. N. Aronszajn and P. Panitchpakdi, Extensions of uniformly continuous transformations and hyperconvex metric spaces, *Pacific J. Math.* **6** (1956), 405–439.
5. S. P. Avann, Metric ternary distributive semi-lattices, *Proc. Amer. Math. Soc.* **12** (1961), 407–414.
6. W. Ballman, Singular spaces of non-positive curvature, in "Sur les groupes hyperboliques d'après Mikhael Gromov" (E. Ghys and P. de la Harpe, Eds.), Progress in Math. 81, Chap. 10, pp. 189–201, Birkhäuser, Boston/Basel/Berlin, 1990.
7. H.-J. Bandelt, Retracts of hypercubes, *J. Graph Theory* **8** (1984), 501–510.
8. H.-J. Bandelt, Networks with Condorcet solutions, *European J. Oper. Res.* **20** (1985), 314–326.
9. H.-J. Bandelt, Hereditary modular graphs, *Combinatorica* **8** (1988), 149–157.
10. H.-J. Bandelt and V. Chepoi, A Helly theorem in weakly modular space, *Discrete Math.* **126** (1996) 25–39.
11. H.-J. Bandelt and J. Hedlíková, Median algebras, *Discrete Math.* **45** (1983), 1–30.
12. H.-J. Bandelt and E. Pesch, Dismantling absolute retracts of reflexive graphs, *European J. Combin.* **10** (1989), 211–220.
13. H.-J. Bandelt and M. van de Vel, A fixed cube theorem for median graphs, *Discrete Math.* **62** (1987), 129–137.
14. H.-J. Bandelt and M. van de Vel, Embedding topological median algebras in product of dendrons, *Proc. London Math. Soc.* **58** (1989), 439–453.
15. H.-J. Bandelt and M. Van de Vel, Superextensions and the depth of median graphs, *J. Combin. Theory Ser. A* **57** (1991), 187–202.
16. M. R. Bridson, Geodesics and curvature in metric simplicial complexes, in "Group Theory from a Geometrical Point of View (E. Ghys, A. Haefliger, and A. Verjovsky, Eds.), pp. 373–463, World Scientific, Singapore, 1991.

17. M. Bridson and A. Haefliger, "Metric Spaces of Non-Positive Curvature," forthcoming book.
18. H. Busemann, "The Geometry of Geodesics," Academic Press, New York, 1955.
19. R. Charney, Artin groups of finite type are biautomatic, *Math. Ann.* **292** (1992), 671–683.
20. R. Charney and M. W. Davis, Singular metrics of nonpositive curvature on branched covers of Riemannian manifolds, *Amer. J. Math.* **115** (1993), 929–1009.
21. R. Charney and M. W. Davis, The  $K(\pi, 1)$ -problem for hyperplane complements associated to infinite reflection groups, *J. Amer. Math. Soc.* **8** (1995), 597–627.
22. V. Chepoi, Classifying graphs by metric triangles (in Russian), *Metody Diskret. Anal.* **49** (1989), 75–93.
23. V. Chepoi, Bridged graphs are cop-win graphs: an algorithmic proof, *J. Combin. Theory Ser. B* **69** (1997), 97–100.
24. V. Chepoi, On distance-preserving and domination elimination orderings, *SIAM J. Discrete Math.* **11** (1998), 414–436.
25. D. Ž. Djoković, Distance-preserving subgraphs of hypercubes, *J. Combin. Theory Ser. B* **14** (1973), 263–267.
26. A. W. M. Dress and R. Scharlau, Gated sets in metric spaces, *Aequationes Math.* **34** (1987), 112–120.
27. A. Dress, K. Huber, and V. Moulton, Some variations on a theme by Peter Buneman, *Ann. Combin.* **1** (1997), 339–352.
28. D. B. A. Epstein, J. W. Cannon, D. F. Holt, S. V. F. Levy, M. S. Paterson, and W. P. Thurston, "Word Processing in Groups," Jones and Bartlett, Boston, 1992.
29. S. M. Gersten and H. B. Short, Small cancellation theory and automatic groups, *Invent. Math.* **102** (1990), 305–334.
30. S. M. Gersten and H. M. Short, Small cancellation theory and automatic groups: Part II, *Invent. Math.* **105** (1991), 641–662.
31. M. Gromov, Hyperbolic groups, in "Essays in Group Theory" (S. M. Gersten, Ed.), Mathematical Sciences Research Institute Publications 8, pp. 75–263, Springer, Berlin, 1987.
32. M. C. Golumbic, "Algorithmic Graph Theory and Perfect Graphs," Academic Press, New York, 1980.
33. M. Farber and R. E. Jamison, On local convexity in graphs, *Discrete Math.* **66** (1987), 231–247.
34. J. R. Isbell, Six theorems about injective metric spaces, *Comment. Math. Helv.* **39** (1964), 65–74.
35. J. R. Isbell, Median algebra, *Trans. Amer. Math. Soc.* **260** (1980), 319–362.
36. E. Jawhari, D. Misane, and M. Pouzet, Retracts: graphs and ordered sets from the metric point of view, *Contemp. Math.* **56** (1986), 175–225.
37. A. V. Karzanov, Minimum 0-extensions of graph metrics, *European J. Combin.* **19** (1998), 71–101.
38. A. V. Karzanov, "Metrics with Finite Sets of Primitive Extensions," Universität Bielefeld SFB343, preprint 97–110, 1997.
39. W. A. Kirk, An abstract fixed point theorem for nonexpansive mappings, *Proc. Amer. Math. Soc.* **82** (1981), 640–642.
40. R. C. Lyndon and P. E. Schupp, "Combinatorial Group Theory," Springer-Verlag, Berlin/Heidelberg/New York, 1977.
41. J.-H. Mai and Y. Tang, An injective metrization for collapsible polyhedra, *Proc. Amer. Math. Soc.* **88** (1983), 333–337.
42. H. M. Mulder, "The Interval Function of a Graph," Math. Centre Tracts 132, Amsterdam, 1980.
43. H. M. Mulder and A. Schrijver, Median graphs and Helly hypergraphs, *Discrete Math.* **25** (1979), 41–50.

44. G. A. Niblo and L. D. Reeves, The geometry of cube complexes and the complexity of their fundamental groups, *Topology* **37** (1998), 621–633.
45. G. A. Noskov, “Bicombing of Triangle Buildings,” Universität Bielefeld SFB343, preprint 95–038, 1995.
46. R. Nowakowski and I. Rival, The smallest graph variety containing all paths, *Discrete Math.* **43** (1983), 223–234.
47. F. Paulin, Constructions of hyperbolic groups via hyperbolizations of polyhedra, in “Group Theory from a Geometrical Point of View” (E. Ghys, A. Haefliger, and A. Verjovsky, Eds.), pp. 313–372, World Scientific, Singapore, 1991.
48. J. P. Penot, Fixed point theorem without convexity, *Bull. Soc. Math. France* **60** (1979), 129–152.
49. L. Reeves, Rational subgroups of cubed 3-manifold groups, *Michigan Math. J.* **42** (1994), 109–126.
50. A. Quilliot, On the Helly property working as a compactness criterion on graphs, *J. Combin. Theory Ser. A* **40** (1985), 186–193.
51. M. Sageev, Ends of group pairs and non-positively curved cube complexes, *Proc. London Math. Soc.* **71** (1995), 585–617.
52. H. Seifert and W. Threlfall, “A Textbook of Topology,” Academic Press, New York, 1980.
53. V. Soltan and V. Chepoi, Conditions for invariance of set diameters under  $d$ -convexification in a graph, *Cybernetics* **19** (1983), 750–756.
54. M. van de Vel, Matching binary convexities, *Topology Appl.* **16** (1983), 207–235.
55. M. van de Vel, “Theory of Convex Structures,” Elsevier, Amsterdam, 1993.
56. M. van de Vel, Collapsible polyhedra and median spaces, *Proc. Amer. Math. Soc.* **126** (1998), 2811–2818.