

# Dismantlability of weakly systolic complexes and applications

VICTOR CHEPOI<sup>1</sup> and DAMIAN OSAJDA<sup>2</sup>

<sup>1</sup>Laboratoire d'Informatique Fondamentale,  
Université d'Aix-Marseille,  
Faculté des Sciences de Luminy,  
F-13288 Marseille Cedex 9, France  
chepoi@lif.univ-mrs.fr

<sup>2</sup> Instytut Matematyczny, Uniwersytet Wrocławski  
pl. Grunwaldzki 2/4, 50-384 Wrocław, Poland  
and  
Université de Lille I, Laboratoire Paul Painlevé  
F-59655 Villeneuve d'Ascq, France  
dosaj@math.uni.wroc.pl

**Abstract.** In this paper, we investigate the structural properties of weakly systolic complexes introduced recently by the second author and of their 1-skeletons, the weakly bridged graphs. We present several characterizations of weakly systolic complexes and weakly bridged graphs. Then we prove that weakly bridged graphs are dismantlable. Using this, we establish the fixed point theorem for weakly systolic complexes. As a consequence, we get results about conjugacy classes of finite subgroups and classifying spaces for finite subgroups of weakly systolic groups. As immediate corollaries, we obtain new results on systolic complexes and systolic groups.

*MSC:* 20F65; 20F67; 05C12; 05C63.

*Keywords:* weakly systolic complex, weakly bridged graph, weakly systolic group, dismantlability, fixed point theorem.

## 1. INTRODUCTION

In his seminal paper [Gro87], among many other results, Gromov gave a pretty combinatorial characterization of  $\text{CAT}(0)$  cubical complexes as simply connected cubical complexes in which the links of vertices are simplicial flag complexes. Based on this result, [Che00, Rol98] established a bijection between the 1-skeletons of  $\text{CAT}(0)$  cubical complexes and the median graphs, well-known in metric graph theory [BC08]. A similar combinatorial characterization of  $\text{CAT}(0)$  simplicial complexes having regular Euclidean simplices as cells seems to be out of reach. Nevertheless, [Che00] characterized the bridged complexes (i.e., the simplicial complexes having bridged graphs as 1-skeletons) as the simply connected simplicial complexes in which the links of vertices are flag complexes without embedded 4- and 5-cycles; the bridged graphs are exactly the graphs which satisfy one of the basic feature of  $\text{CAT}(0)$  spaces: the

balls around convex sets are convex. Bridged graphs have been introduced and characterized in [FJ87, SC83] as graphs without embedded isometric cycles of length greater than 3 and have been further investigated in several graph-theoretical and algebraic papers; cf. [AF88, BC96, Che97, Pol02, Pol00] and the survey [BC08]. Januszkiewicz-Swiatkowski [JS06] and Haglund [Hag03] rediscovered this class of simplicial complexes (they call them *systolic complexes*) using them (and groups acting on them geometrically—*systolic groups*) fruitfully in the context of geometric group theory. Systolic complexes and groups turned out to be good combinatorial analogs of  $CAT(0)$  (nonpositively curved) metric spaces and groups; cf. [Hag03, JS06, Osa07, OP09, Prz08, Prz09].

One of the characteristic features of systolic complexes, related with convexity of balls around convex sets, is the following  $SD_n(\sigma^*)$  property introduced in [Osa09]: *if a simplex  $\sigma$  of a simplicial complex  $\mathbf{X}$  is located in the sphere of radius  $n + 1$  centered at some simplex  $\sigma^*$  of  $\mathbf{X}$ , then the set of all vertices  $x$  such that  $\sigma \cup \{x\}$  is a simplex and  $x$  has distance  $n$  to  $\sigma^*$  is a nonempty simplex  $\sigma_0$  of  $\mathbf{X}$ .* Relaxing this condition, Osajda [Osa09] called a simplicial complex  $\mathbf{X}$  *weakly systolic* if the property  $SD_n(\sigma^*)$  holds whenever  $\sigma^*$  is a vertex (i.e., a 0-dimensional simplex) of  $\mathbf{X}$ . He further showed that this  $SD_n$  property is equivalent with the  $SD_n(\sigma^*)$  property in which  $\sigma^*$  is a vertex and  $\sigma$  is a vertex or an edge (i.e., a 1-dimensional simplex) of  $\mathbf{X}$ . Finally, it is showed in [Osa09] that weakly systolic complexes can be characterized as simply connected simplicial complexes satisfying some local combinatorial conditions, cf. also Theorem A below. This is analogous to the cases of  $CAT(0)$  cubical complexes and systolic complexes. In graph-theoretical terms, the 1-skeletons of weakly systolic complexes (which we call *weakly bridged graphs*) satisfy the so-called triangle and quadrangle conditions [BC96], i.e., like median and bridged graphs, the weakly bridged graphs are weakly modular graphs. As is shown in [Osa09] and in this paper, the properties of weakly systolic complexes resemble very much properties of spaces of non-positive curvature.

The initial motivation of [Osa09] for introducing weakly systolic complexes was to exhibit a class of simplicial complexes with some kind of simplicial nonpositive curvature that will include the systolic complexes and some other classes of complexes appearing in the context of geometric group theory. As we noticed already, systolic complexes are weakly systolic. Moreover, for every simply connected locally 5-large cubical complex (i.e.  $CAT(-1)$  cubical complex [Gro87]) there exists a canonically associated simplicial complex, which is weakly systolic [Osa09]. In particular, the class of *weakly systolic groups*, i.e., groups acting geometrically by automorphisms on weakly systolic complexes, contains the class of  $CAT(-1)$  cubical groups and is therefore essentially bigger than the class of systolic groups; cf. [Osa07]. Other classes of weakly systolic groups are presented in [Osa09]. The ideas and results from [Osa09] allowed to construct in [Osa09b] new examples of Gromov hyperbolic groups of arbitrarily large (virtual) cohomological dimension. Furthermore, Osajda [Osa09] and Osajda-Świątkowski [OŚ09] provide new examples of high dimensional groups with interesting asphericity properties. On the other hand, as we will show below, the class of weakly systolic complexes seems also to

appear naturally in the context of graph theory and have not been studied before from this point of view.

In this paper, we present further characterizations and properties of weakly systolic complexes and their 1-skeletons, weakly bridged graphs. Relying on techniques from graph theory we establish dismantlability of locally-finite weakly bridged graphs. This result is used to show some interesting nonpositive-curvature-like properties of weakly systolic complexes and groups (see [Osa09] for other properties of this kind). As corollaries, we get also new results about systolic complexes and groups. We conclude this introductory section with the formulation of our main results (see respective sections for all missing definitions and notations as well as for other related results).

We start with a characterization of weakly systolic complexes proved in Section 3:

**Theorem A.** *For a flag simplicial complex  $\mathbf{X}$  the following conditions are equivalent:*

- (a)  $\mathbf{X}$  is weakly systolic;
- (b) the 1-skeleton of  $\mathbf{X}$  is a weakly modular graph without induced  $C_4$ ;
- (c) the 1-skeleton of  $\mathbf{X}$  is a weakly modular graph with convex balls;
- (d) the 1-skeleton of  $\mathbf{X}$  is a graph with convex balls in which any  $C_5$  is included in a 5-wheel  $W_5$ ;
- (e)  $\mathbf{X}$  is simply connected, satisfies the  $\widehat{W}_5$ -condition, and does not contain induced  $C_4$ .

In Section 5 we prove the following result:

**Theorem B.** *Any LexBFS ordering of vertices of a locally finite weakly systolic complex  $\mathbf{X}$  is a dismantling ordering of its 1-skeleton.*

This dismantlability result has several consequences presented in Section 5. This result also allows us to prove in Section 6 the following fixed point theorem concerning group actions:

**Theorem C.** *Let  $G$  be a finite group acting by simplicial automorphisms on a locally finite weakly systolic complex  $\mathbf{X}$ . Then there exists a simplex  $\sigma \in \mathbf{X}$  which is invariant under the action of  $G$ .*

The barycenter of an invariant simplex is a point fixed by  $G$ . An analogous theorem holds in the case of  $CAT(0)$  spaces; cf. [BH99, Corollary 2.8]. As a direct corollary of Theorem C, we get the fixed point theorem for systolic complexes. This was conjectured by Januszkiewicz-Świątkowski (personal communication) and Wise [Wis03], and later was formulated in the collection of open questions [08, Conjecture 40.1 on page 115]. A partial result in the systolic case was proved by Przytycki [Prz08]. In fact, in Section 8, based on a result of Polat [Pol02] for bridged graphs, we prove even a stronger version of the fixed point theorem in this case.

There are several important group theoretical consequences of Theorem C. The first one follows directly from this theorem and [Prz08, Remarks 7.7&7.8].

**Theorem D.** *Let  $k \geq 6$ . Free products of  $k$ -systolic groups amalgamated over finite subgroups are  $k$ -systolic. HNN extensions of  $k$ -systolic groups over finite subgroups are  $k$ -systolic.*

The following result (Corollary 6.4 below) has also its  $CAT(0)$  counterpart; cf. [BH99, Corollary 2.8]:

**Corollary.** *Let  $G$  be a weakly systolic group. Then  $G$  contains only finitely many conjugacy classes of finite subgroups.*

The next important consequence of the fixed point theorem concerns classifying spaces for proper group actions. Recall that if a group  $G$  acts properly on a space  $\mathbf{X}$  such that the fixed point set for any finite subgroup of  $G$  is contractible (and therefore non-empty), then we say that  $\mathbf{X}$  is a *model for  $\underline{EG}$* —the classifying space for finite groups. If additionally the action is cocompact, then  $\mathbf{X}$  is a *finite model for  $\underline{EG}$* . A (finite) model for  $\underline{EG}$  is in a sense a “universal”  $G$ -space (see [Lüc05] for details). The following theorem is a direct consequence of Theorem C and Proposition 7.6 below.

**Theorem E.** *Let  $G$  act properly by simplicial automorphisms on a finite dimensional weakly systolic complex  $\mathbf{X}$ . Then  $\mathbf{X}$  is a finite dimensional model for  $\underline{EG}$ . If, moreover, the action of  $G$  on  $\mathbf{X}$  is cocompact, then  $\mathbf{X}$  is a finite model for  $\underline{EG}$ .*

As an immediate consequence we get an analogous result about  $\underline{EG}$  for systolic groups. This was conjectured in [08, Chapter 40]. Przytycki [Prz09] showed that the Rips complex (with the constant at least 5) of a systolic complex is an  $\underline{EG}$  space. Our result gives a systolic—and thus much nicer—model of  $\underline{EG}$  in that case.

In the final Section 8 we present some further results about systolic complexes and groups. Besides a stronger version of the fixed point theorem mentioned above, we remark on another approach to this theorem initiated by Zawiślak [Zaw04] and Przytycki [Prz08]. In particular, our Proposition 8.5 proves their conjecture about round complexes; cf. [Zaw04, Conjecture 3.3.1] and [Prz08, Remark 8.1]. Finally, we show (cf. Claim 8.7) how our results about  $\underline{EG}$  apply to the questions of existence of particular boundaries of systolic groups (and thus to the Novikov conjecture for systolic groups with torsion). This relies on earlier results of Osajda-Przytycki [OP09].

## 2. PRELIMINARIES

**2.1. Graphs and simplicial complexes.** We continue with basic definitions used in this paper concerning graphs and simplicial complexes. All graphs  $G = (V, E)$  occurring here are undirected, connected, and without loops or multiple edges. The *distance*  $d(u, v)$  between two vertices  $u$  and  $v$  is the length of a shortest  $(u, v)$ -path, and the *interval*  $I(u, v)$  between  $u$  and  $v$  consists of all vertices on shortest  $(u, v)$ -paths, that is, of all vertices (metrically) *between*  $u$  and  $v$ :

$$I(u, v) = \{x \in V : d(u, x) + d(x, v) = d(u, v)\}.$$

An induced subgraph of  $G$  (or the corresponding vertex set  $A$ ) is called *convex* if it includes the interval of  $G$  between any of its vertices. By the *convex hull*  $\text{conv}(W)$  of  $W$  in  $G$  we mean the smallest convex subset of  $V$  (or induced subgraph of  $G$ ) that contains  $W$ . An *isometric*

*subgraph* of  $G$  is an induced subgraph in which the distances between any two vertices are the same as in  $G$ . In particular, convex subgraphs are isometric. The *ball* (or disk)  $B_r(x)$  of center  $x$  and radius  $r \geq 0$  consists of all vertices of  $G$  at distance at most  $r$  from  $x$ . In particular, the unit ball  $B_1(x)$  comprises  $x$  and the neighborhood  $N(x)$  of  $x$ . The *sphere*  $S_r(x)$  of center  $x$  and radius  $r \geq 0$  consists of all vertices of  $G$  at distance exactly  $r$  from  $x$ . The ball  $B_r(S)$  centered at a convex set  $S$  is the union of all balls  $B_r(x)$  with centers  $x$  from  $S$ . The *sphere*  $S_r(S)$  of center  $S$  and radius  $r \geq 0$  consists of all vertices of  $G$  at distance exactly  $r$  from  $S$ .

A graph  $G$  is called *thin* if for any two nonadjacent vertices  $u, v$  of  $G$  any two neighbors of  $v$  in the interval  $I(u, v)$  are adjacent. A graph  $G$  is *weakly modular* [BC96, BC08] if its distance function  $d$  satisfies the following conditions:

*Triangle condition* (T): for any three vertices  $u, v, w$  with  $1 = d(v, w) < d(u, v) = d(u, w)$  there exists a common neighbor  $x$  of  $v$  and  $w$  such that  $d(u, x) = d(u, v) - 1$ .

*Quadrangle condition* (Q): for any four vertices  $u, v, w, z$  with  $d(v, z) = d(w, z) = 1$  and  $2 = d(v, w) \leq d(u, v) = d(u, w) = d(u, z) - 1$ , there exists a common neighbor  $x$  of  $v$  and  $w$  such that  $d(u, x) = d(u, v) - 1$ .

An abstract *simplicial complex*  $\mathbf{X}$  is a collection of sets (called *simplices*) such that  $\sigma \in \mathbf{X}$  and  $\sigma' \subseteq \sigma$  implies  $\sigma' \in \mathbf{X}$ . The *geometric realization*  $|\mathbf{X}|$  of a simplicial complex is the polyhedral complex obtained by replacing every face  $\sigma$  of  $\mathbf{X}$  by a “solid” regular simplex  $|\sigma|$  of the same dimension such that realization commutes with intersection, that is,  $|\sigma' \cap \sigma''| = |\sigma' \cap \sigma''|$  for any two simplices  $\sigma'$  and  $\sigma''$ . Then  $|\mathbf{X}| = \bigcup\{|\sigma| : \sigma \in \mathbf{X}\}$ .  $\mathbf{X}$  is called *simply connected* if it is connected and if every continuous mapping of the 1-dimensional sphere  $S^1$  into  $|\mathbf{X}|$  can be extended to a continuous mapping of the disk  $D^2$  with boundary  $S^1$  into  $|\mathbf{X}|$ .

For a simplicial complex  $\mathbf{X}$ , denote by  $V(\mathbf{X})$  and  $E(\mathbf{X})$  the *vertex set* and the *edge set* of  $\mathbf{X}$ , namely, the set of all 0-dimensional and 1-dimensional simplices of  $\mathbf{X}$ . The pair  $(V(\mathbf{X}), E(\mathbf{X}))$  is called the (*underlying*) *graph* or the *1-skeleton* of  $\mathbf{X}$  and is denoted by  $G(\mathbf{X})$ . Conversely, for a graph  $G$  one can derive a simplicial complex  $\mathbf{X}(G)$  (the *clique complex* of  $G$ ) by taking all complete subgraphs (cliques) as simplices of the complex. A simplicial complex  $\mathbf{X}$  is a *flag complex* (or a *clique complex*) if any set of vertices is included in a face of  $\mathbf{X}$  whenever each pair of its vertices is contained in a face of  $\mathbf{X}$  (in the theory of hypergraphs this condition is called conformality). A flag complex can therefore be recovered by its underlying graph  $G(\mathbf{X})$ : the complete subgraphs of  $G(\mathbf{X})$  are exactly the simplices of  $\mathbf{X}$ . The *link* of a simplex  $\sigma$  in  $\mathbf{X}$ , denoted  $\text{lk}(\sigma, \mathbf{X})$  is the simplicial complex consisting of all simplexes  $\sigma'$  such that  $\sigma \cap \sigma' = \emptyset$  and  $\sigma \cup \sigma' \in \mathbf{X}$ . For a simplicial complex  $\mathbf{X}$  and a vertex  $v$  not belonging to  $\mathbf{X}$ , the *cone* with apex  $v$  and base  $\mathbf{X}$  is the simplicial complex  $v * \mathbf{X} = \mathbf{X} \cup \{\sigma \cup \{v\} : \sigma \in \mathbf{X}\}$ .

For a simplicial complex  $\mathbf{X}$  and any  $k \geq 1$ , the *Rips complex*  $\mathbf{X}_k$  is a simplicial complex with the same set of vertices as  $\mathbf{X}$  and with a simplex spanned by any subset  $S \subset V(\mathbf{X})$  such that  $d(u, v) \leq k$  in  $G(\mathbf{X})$  for each pair of vertices  $u, v \in S$  (i.e.,  $S$  has diameter  $\leq k$  in the graph  $G(\mathbf{X})$ ). From the definition immediately follows that the Rips complex of any complex is a flag complex. Alternatively, the Rips complex  $\mathbf{X}_k$  can be viewed as the clique complex

$\mathbf{X}(G^k(\mathbf{X}))$  of the  $k$ th power of the graph of  $\mathbf{X}$  (the  $k$ th power  $G^k$  of a graph  $G$  has the same set of vertices as  $G$  and two vertices  $u, v$  are adjacent in  $G^k$  if and only if  $d(u, v) \leq k$  in  $G$ ).

**2.2.  $SD_n$  property and weakly systolic complexes.** The following generalization of systolic complexes has been presented by Osajda [Osa09]. A flag simplicial complex  $\mathbf{X}$  satisfies the property of *simple descent on balls* of radii at most  $n$  centered at a simplex  $\sigma^*$  (*property  $SD_n(\sigma^*)$* ) [Osa09] if for each  $i = 0, 1, 2, \dots, n$  and each simplex  $\sigma$  located in the sphere  $S_{i+1}(\sigma^*)$  the set  $\sigma_0 := V(\text{lk}(\sigma^*, \mathbf{X})) \cap B_i(\sigma^*)$  spans a non-empty simplex of  $\mathbf{X}$ . Systolic complexes are exactly the flag complexes which satisfy the  $SD_n(\sigma^*)$  property for all simplices  $\sigma^*$  and all natural numbers  $n$ . On the other hand, the 5-wheel is an example of a (2-dimensional) simplicial complex which satisfies the  $SD_2$  property for vertices and triangles but not for edges. In view of this analogy and of subsequent results, we call *weakly systolic* a flag simplicial complex  $\mathbf{X}$  which satisfies the  $SD_n(v)$  property for all vertices  $v \in V(\mathbf{X})$  and for all natural numbers  $n$ . We also call *weakly bridged* the underlying graphs of weakly systolic complexes. It can be shown (cf. Theorem 3.1) that  $\mathbf{X}$  is a weakly systolic complex if for each vertex  $v$  and every  $i$  it satisfies the following two conditions:

*Vertex condition* (V): for every vertex  $w \in S_{i+1}(v)$ , the intersection  $V(\text{lk}(v, \mathbf{X})) \cap B_i(v)$  is a single simplex;

*Edge condition* (E): for every edge  $e \in S_{i+1}(v)$ , the intersection  $V(\text{lk}(e, \mathbf{X})) \cap B_i(v)$  is nonempty.

In fact, this is the original definition of a weakly systolic complex given in [Osa09]. Notice that these two conditions imply that weakly systolic complexes are exactly the flag complexes whose underlying graphs are thin and satisfy the triangle condition.

**2.3. Dismantlability of graphs and LC-contractibility of complexes.** Let  $G = (V, E)$  be a graph and  $u, v$  two vertices of  $G$  such that any neighbor of  $v$  (including  $v$  itself) is also a neighbor of  $u$ . Then there is a retraction of  $G$  to  $G - v$  taking  $v$  to  $u$ . Following [HN04], we call this retraction a *fold* and we say that  $v$  is *dominated* by  $u$ . A finite graph  $G$  is *dismantlable* if it can be reduced, by a sequence of folds, to a single vertex. In other words, an  $n$ -vertex graph  $G = (V, E)$  is dismantlable if its vertices can be ordered  $v_1, \dots, v_n$  so that for each vertex  $v_i, 1 \leq i < n$ , there exists another vertex  $v_j$  with  $j > i$ , such that  $N_1(v_i) \cap V_i \subseteq N_1(v_j) \cap X_i$ , where  $V_i := \{v_i, v_{i+1}, \dots, v_n\}$ . This order is called a *dismantling order*. Consider now for simplicial complexes  $\mathbf{X}$  the analogy of dismantlability investigated in the papers [CY07, Mat08]. A vertex  $v$  of  $\mathbf{X}$  is *LC-removable* if  $\text{lk}(v, \mathbf{X})$  is a cone. If  $v$  is an LC-removable vertex of  $\mathbf{X}$ , then  $\mathbf{X} - v := \{\sigma \in \mathbf{X} : v \notin \sigma\}$  is obtained from  $\mathbf{X}$  by an *elementary LC-reduction* (link-cone reduction) [Mat08]. Then  $\mathbf{X}$  is called *LC-contractible* [CY07] if there is a sequence of elementary LC-reductions transforming  $\mathbf{X}$  to one vertex. For flag simplicial complexes, the LC-contractibility of  $\mathbf{X}$  is equivalent to dismantlability of its graph  $G(\mathbf{X})$  because an LC-removable vertex  $v$  is dominated by the apex of the cone  $\text{lk}(v, \mathbf{X})$  and vice versa the link of any dominated vertex  $v$  is a cone having the vertex dominating

$v$  as its apex. It is clear that LC-contractible simplicial complexes are collapsible (see also [CY07, Corollary 6.5]).

Dismantlable graphs are closed under retracts and direct products, i.e., they constitute a variety [NW83]. Winkler and Nowakowski [NW83] and Quilliot [Qui83] characterized the dismantlable graphs as cop-win graphs, i.e., graphs in which a single cop captures the robber after a finite number of moves for all possible initial positions of two players. Cops and robber game is a pursuit-evasion game played on finite (or infinite) undirected graphs in which the two players move alternatively starting from their initial positions, where a move is to slide along an edge or to stay at the same vertex. The objective of the cop is to capture the robber, i.e., to be at some moment of time at the same vertex as the robber. The objective of the robber is to continue evading the cops.

The simplest algorithmic way to order the vertices of a graph is to apply the *Breadth-First Search* (BFS) starting from the root vertex (base point)  $b$ . We number with 1 the vertex  $u$  and put it on the initially empty queue. We repeatedly remove the vertex  $v$  at the head of the queue and consequently number and place onto the queue all still unnumbered neighbors of  $v$ . BFS constructs a spanning tree  $T_u$  of  $G$  with the vertex  $u$  as a root. Then a vertex  $v$  is the *father* in  $T_v$  of any of its neighbors  $w$  in  $G$  included in the queue when  $v$  is removed (notation  $f(w) = v$ ). The procedure is called once for each vertex  $v$  and proceeds  $v$  in  $O(|deg(v)|)$  time, so the total complexity of its implementation is linear. Notice that the distance from any vertex  $v$  to the root  $u$  is the same in  $G$  and in  $T_u$ . Another method to order the vertices of a graph in linear time is the *Lexicographic Breadth-First Search* (LexBFS) proposed by Rose, Tarjan, and Lueker [RTL76]. According to LexBFS, the vertices of a graph  $G$  are numbered from  $n$  to 1 in decreasing order. The *label*  $L(w)$  of an unnumbered vertex  $w$  is the list of its numbered neighbors. As the next vertex to be numbered, select the vertex with the lexicographic largest label, breaking ties arbitrarily. As in case of BFS, we remove the vertex  $v$  at the head of the queue and consequently number according to the lexicographic order and place onto the queue all still unnumbered neighbors of  $v$ . LexBFS is a particular instance of BFS, i.e., every ordering produced by LexBFS can also be generated by BFS.

Anstee and Farber [AF88] established that bridged graphs are cop-win graphs. Chepoi [Che97] noticed that any order of a bridged graph returned by BFS is a dismantling order. Namely, he showed a stronger result: *for any two adjacent vertices  $v_i, v_j$  with  $i < j$ , their fathers  $f(v_i), f(v_j)$  either coincide or are adjacent and moreover  $f(v_j)$  is adjacent to  $v_i$* . This property implies that bridged graphs admits a geodesic 1-combing and that the shortest paths participating in this combing are the paths to the root  $u$  of the BFS tree  $T_u$  [Che00]. Similar results have been established in [Che98] for larger classes of weakly modular graphs by using LexBFS instead of BFS.

Notice that the notions of dismantlable graph, BFS, and LexBFS can be defined in a straightforward way to all locally finite graphs. Polat [Pol02, Pol00] defined dismantlability and BFS for arbitrary (not necessarily locally finite) graphs and extended the results of [AF88, Che97] to all bridged graphs.




FIGURE 1. The  $\widehat{W}_5$ -condition

**2.4. Group actions on simplicial complexes.** Let  $G$  be a group acting by automorphisms on a simplicial complex  $\mathbf{X}$ . By  $\text{Fix}_G \mathbf{X}$  we denote the *fixed point set* of the action of  $G$  on  $\mathbf{X}$ , i.e.  $\text{Fix}_G \mathbf{X} = \{x \in \mathbf{X} \mid Gx = \{x\}\}$ . Recall that the action is *cocompact* if the orbit space  $G \backslash \mathbf{X}$  is compact. The action of  $G$  on a locally finite simplicial complex  $\mathbf{X}$  is *properly discontinuous* if stabilizers of simplices are finite. Finally, the action is *geometric* (or  $G$  *acts geometrically* on  $\mathbf{X}$ ) if it is cocompact and properly discontinuous.

### 3. CHARACTERIZATIONS OF WEAKLY SYSTOLIC COMPLEXES

We continue with the characterizations of weakly systolic complexes and their underlying graphs; some of those characterizations have been presented also in [Osa09]. We denote by  $C_k$  an induced  $k$ -cycle and by  $W_k$  an induced  $k$ -wheel, i.e., an induced  $k$ -cycle  $x_1, \dots, x_k$  plus a central vertex  $c$  adjacent to all vertices of  $C_k$ .  $W_k$  can also be viewed as a 2-dimensional simplicial complex consisting of  $k$  triangles  $\sigma_1, \dots, \sigma_k$  sharing a common vertex  $c$  and such that  $\sigma_i$  and  $\sigma_j$  intersect in an edge  $x_i c$  exactly when  $|j - i| = 1 \pmod{k}$ . In other words,  $\text{lk}(c, \mathbf{X}) = C_k$ . By  $\widehat{W}_k$  we denote a  $k$ -wheel  $W_k$  plus a triangle  $ax_i x_{i+1}$  for some  $i < k$  (we suppose that  $a \neq c$  and that  $a$  is not adjacent to any other vertex of  $W_k$ ). We continue with a condition which basically characterizes weakly bridged complexes among simply connected flag simplicial complexes:

$\widehat{W}_5$ -condition: for any  $\widehat{W}_5$ , there exists a vertex  $v \notin \widehat{W}_5$  such that  $\widehat{W}_5$  is included in  $\text{lk}(v, \mathbf{X})$ , i.e.,  $v$  is adjacent in  $G(\mathbf{X})$  to all vertices of  $\widehat{W}_5$  (see Fig. 1).

**Theorem 3.1** (Characterizations). *For a flag simplicial complex  $\mathbf{X}$  the following conditions are equivalent:*

- (i)  $\mathbf{X}$  is weakly systolic;
- (ii)  $\mathbf{X}$  satisfies the the vertex condition (V) and the edge condition (E);
- (iii)  $G(\mathbf{X})$  is a weakly modular thin graph;
- (iv)  $G(\mathbf{X})$  is a weakly modular graph without induced  $C_4$ ;
- (v)  $G(\mathbf{X})$  is a weakly modular graph with convex balls;

- (vi)  $G(\mathbf{X})$  is a graph with convex balls in which any  $C_5$  is included in a 5-wheel  $W_5$ ;
- (vii)  $\mathbf{X}$  is simply connected, satisfies the  $\widehat{W}_5$ -condition, and does not contain induced  $C_4$ .

*Proof.* The implications (i) $\Rightarrow$ (ii) and (iii) $\Rightarrow$ (iv) are obvious.

(ii) $\Rightarrow$ (iii): The condition (V) implies that all vertices of  $I(u, v)$  adjacent to  $v$  are pairwise adjacent, i.e., that  $G(\mathbf{X})$  is thin. On the other hand, from the condition (E) we conclude that if  $1 = d(v, w) < d(u, v) = d(u, w) = i + 1$ , then  $v$  and  $w$  have a common neighbor  $x$  in the sphere  $S_i(u)$ , implying the triangle condition. Finally, in thin graphs the quadrangle condition is automatically satisfied. This shows that  $G(\mathbf{X})$  is a weakly modular thin graph.

(iv) $\Rightarrow$ (v): To show that any ball  $B_i(u)$  is convex in  $G(\mathbf{X})$ , since  $G(\mathbf{X})$  is weakly modular and  $B_i(u)$  induces a connected subgraph, according to [Che89] it suffices to show that the ball  $B_i(u)$  is locally convex, i.e., if  $x, y \in B_i(u)$  and  $d(x, y) = 2$ , then  $I(x, y) \subseteq B_i(u)$ . Suppose by way of contradiction that  $z \in I(x, y) \setminus B_i(u)$ . Then necessarily  $d(x, u) = d(y, u) = i$  and  $d(z, u) = i + 1$ . Applying the quadrangle condition, we infer that there exists a vertex  $z'$  adjacent to  $x$  and  $y$  at distance  $i - 1$  from  $u$ . As a result, the vertices  $x, z, y, z'$  induce a forbidden 4-cycle, a contradiction.

(v) $\Rightarrow$ (vi): Pick a 5-cycle induced by the vertices  $x_1, x_2, x_3, x_4, x_5$ . Since  $d(x_4, x_1) = d(x_4, x_2) = 2$ , by the triangle condition there exists a vertex  $y$  adjacent to  $x_1, x_2$ , and  $x_4$ . Since  $G(\mathbf{X})$  does not contain induced 4-cycles, necessarily  $y$  must be also adjacent to  $x_3$  and  $x_5$ , yielding a 5-wheel.

(v) $\Rightarrow$ (i): Pick a simplex  $\sigma$  in the sphere  $S_{i+1}(u)$ . Denote by  $\sigma_0$  the set of all vertices  $x \in S_i(u)$  such that  $\sigma \cup \{x\}$  is a simplex of  $\mathbf{X}$ . Since the balls of  $G$  are convex, necessarily either  $\sigma_0$  is empty or the vertices of  $\sigma_0$  are pairwise adjacent, thus  $\sigma_0$  and  $\sigma \cup \sigma_0$  induce complete subgraphs of  $G(\mathbf{X})$ . Since  $\mathbf{X}$  is a flag complex,  $\sigma_0$  and  $\sigma \cup \sigma_0$  are simplices. Notice that obviously  $\sigma' \subseteq \sigma_0$  holds for any other simplex  $\sigma' \subseteq S_i(u)$  such that  $\sigma \cup \sigma' \in \mathbf{X}$ . Therefore, it remains to show that  $\sigma_0$  is non-empty. Let  $x$  be a vertex of  $S_i(u)$  which is adjacent to the maximum number of vertices of  $\sigma$ . Since  $G(\mathbf{X})$  is weakly modular and  $\sigma$  is contained in  $S_{i+1}(u)$ , the vertex  $x$  must be adjacent to at least two vertices of  $\sigma$ . Suppose by way of contradiction that  $x$  is not adjacent to a vertex  $v \in \sigma$ . Pick any neighbor  $w$  of  $x$  in  $\sigma$ . By triangle condition, there exists a vertex  $y \in S_i(u)$  adjacent to  $v$  and  $w$ . Since  $w$  is adjacent to  $x, y \in S_i(u)$  and  $w \in S_{i+1}(u)$ , the convexity of  $B_i(u)$  implies that  $x$  and  $y$  are adjacent. Pick any other vertex  $w'$  of  $\sigma$  adjacent to  $x$ . Since  $x$  is not adjacent to  $v$  and  $G(\mathbf{X})$  does not contain induced 4-cycles, the vertices  $y$  and  $w'$  must be adjacent. Hence,  $y$  is adjacent to  $v \in \sigma$  and to all neighbors of  $x$  in  $\sigma$ , contrary to the choice of  $x$ . Thus  $x$  is adjacent to all vertices of  $\sigma$ , i.e.,  $\sigma_0 \neq \emptyset$ . This shows that  $\mathbf{X}$  satisfies the  $SD_n(u)$  property.

(vi) $\Rightarrow$ (vii): To show that a flag complex  $\mathbf{X}$  is simply connected, it suffices to prove that every simple cycle in the underlying graph of  $\mathbf{X}$  is a modulo 2 sum of its triangular faces. Notice that the isometric cycles of an arbitrary graph  $G$  constitute a basis of cycles of  $G$ . Since  $G(\mathbf{X})$  is a graph with convex balls, the isometric cycles of  $G(\mathbf{X})$  have length 3 or 5

[FJ87, SC83]. By (vi), any 5-cycle  $C$  of  $G(\mathbf{X})$  extends to a 5-wheel, thus  $C$  is a modulo 2 sum of triangles. Hence  $\mathbf{X}$  is indeed simply connected. That  $\mathbf{X}$  does not contain induced 4-cycles and 4-wheels follows from the convexity of balls. Finally, pick an extended 5-wheel  $\widehat{W}_5$ : let  $x_1, x_2, x_3, x_4, x_5$  be the vertices of the 5-cycle,  $y$  be the center of the 5-wheel, and  $x_1, x_2, z$  be the vertices of the pendant triangle. Since  $x_3$  and  $x_5$  are not adjacent and the balls of  $G(\mathbf{X})$  are convex, necessarily  $d(z, x_4) = 2$ . Let  $u$  be a common neighbor of  $z$  and  $x_4$ . If  $u$  is adjacent to one of the vertices  $x_2$  and  $x_3$ , then by convexity of balls it will be also adjacent to the second vertex and to  $y$ . But if  $u$  is adjacent to  $y$ , then it will be adjacent to  $x_1$  and therefore to  $x_5$  as well. Hence, in this case  $u$  will be adjacent to all vertices  $x_1, x_2, x_3, x_4, x_5$ , and  $y$ , and we are done. So, we can suppose that  $u$  is not adjacent to anyone of the vertices  $x_1, x_2, x_3, x_5$ , and  $y$ . As a result, we obtain two 5-cycles induced by the vertices  $z, x_2, x_3, x_4, u$  and  $z, x_1, x_5, x_4, u$ . Each of these cycles extends to a 5-wheel. Let  $v$  be the center of the 5-wheel extending the first cycle. To avoid a 4-cycle induced by the vertices  $x_2, v, x_4, y$ , the vertices  $v$  and  $y$  must be adjacent. Subsequently, to avoid a 4-cycle induced by the vertices  $y, v, z, x_1$ , the vertices  $v$  and  $x_1$  must be adjacent. Finally, to avoid a 4-cycle induced by  $x_1, v, x_4, x_5$ , the vertices  $v$  and  $x_5$  must be adjacent. This way, we obtained that  $v$  is adjacent to all six vertices of  $\widehat{W}_5$ , establishing the  $\widehat{W}_5$ -condition.

(vii) $\Rightarrow$ (iv): To prove this implication, as in [Che00], we will use the minimal disk diagrams. Let  $\mathcal{D}$  and  $\mathbf{X}$  be two simplicial complexes. A map  $\varphi : V(\mathcal{D}) \rightarrow V(\mathbf{X})$  is called *simplicial* if  $\varphi(\sigma) \in \mathbf{X}$  for all  $\sigma \in \mathcal{D}$ . If  $\mathcal{D}$  is a planar triangulation (i.e. the 1-skeleton of  $\mathcal{D}$  is an embedded planar graph whose all interior 2-faces are triangles) and  $C = \varphi(\partial\mathcal{D})$ , then  $(\mathcal{D}, \varphi)$  is called a *singular disk diagram* (or Van Kampen diagram) for  $C$  (for more details see [LS77, Chapter V]). According to Van Kampen's lemma ([LS77], pp.150–151), for every cycle  $C$  of a simply connected simplicial complex one can construct a singular disk diagram. A singular disk diagram with no cut vertices (i.e., its 1-skeleton is 2-connected) is called a *disk diagram*. A *minimal (singular) disk* of  $C$  is a (singular) disk diagram  $\mathcal{D}$  of  $C$  with a minimum number of 2-faces. This number is called the (*combinatorial*) *area* of  $C$  and is denoted  $\text{Area}(C)$ . The minimal disks diagrams  $(\mathcal{D}, \varphi)$  of simple cycles  $C$  in 1-skeletons of simply connected simplicial complexes have the following properties [Che00]: (1)  $\varphi$  bijectively maps  $\partial\mathcal{D}$  to  $C$  and (2) the image of a 2-simplex of  $\mathcal{D}$  under  $\varphi$  is a 2-simplex, and two adjacent 2-simplices of  $\mathcal{D}$  have distinct images under  $\varphi$ .

Let  $C$  be a simple cycle in the underlying graph  $G(\mathbf{X})$  of a flag simplicial complex  $\mathbf{X}$  satisfying the condition (vii).

**Claim 1:** *If  $C$  has length 5, then the minimal disk diagram of  $C$  is a 5-wheel. Otherwise,  $C$  admits a minimal disk diagram  $\mathcal{D}$  which is a systolic complex, i.e., a plane triangulation whose all inner vertices have degrees  $\geq 6$ .*

**Proof of Claim 1:** First we show that any minimal disk diagram  $\mathcal{D}$  of  $C$  does not contain interior vertices of degrees 3 and 4. Let  $x$  be any interior vertex of  $\mathcal{D}$ . Let  $x_1, \dots, x_k$  be the neighbors of  $x$ , where  $\sigma_i = xx_i x_{i+1(\text{mod } k)}$  ( $i = 1, \dots, k$ ) are the faces incident to  $x$ .

Trivially,  $k \geq 3$ . Suppose by way of contradiction that  $k \leq 4$ . By properties of minimal disk diagrams,  $\varphi(\sigma_1), \dots, \varphi(\sigma_k)$  are distinct 2-simplices of  $\mathbf{X}$ . If  $k = 3$  then the 2-simplices  $\varphi(\sigma_1), \varphi(\sigma_2), \varphi(\sigma_3)$  of  $\mathbf{X}$  intersect in  $\varphi(x)$  and pairwise share an edge of  $\mathbf{X}$ . Therefore they are contained in a 3-simplex of  $\mathbf{X}$ . This implies that  $\delta = \varphi(x_1)\varphi(x_2)\varphi(x_3)$  is a 2-face of  $\mathbf{X}$ . Let  $\mathcal{D}'$  be a disk triangulation obtained from  $\mathcal{D}$  by deleting the vertex  $x$  and the triangles  $\sigma_1, \sigma_2, \sigma_3$ , and adding the 2-simplex  $x_1x_2x_3$ . The map  $\varphi : V(\mathcal{D}') \rightarrow V(\mathbf{X})$  is simplicial, because it maps  $x_1x_2x_3$  to  $\delta$ . Therefore  $(\mathcal{D}', \varphi)$  is a disk diagram for  $C$ , contrary to the minimality choice of  $\mathcal{D}$ . Now, let  $x$  has four neighbors. The cycle  $(x_1, x_2, x_3, x_4, x_1)$  is sent to a 4-cycle of  $\text{lk}(\varphi(x), \mathbf{X})$ , in which two opposite vertices, say  $\varphi(x_1)$  and  $\varphi(x_3)$ , are adjacent. Consequently,  $\delta' = \varphi(x_1)\varphi(x_3)\varphi(x_2)$  and  $\delta'' = \varphi(x_1)\varphi(x_3)\varphi(x_4)$  are 2-faces of  $\mathbf{X}$ . Let  $\mathcal{D}'$  be a disk triangulation obtained from  $\mathcal{D}$  by deleting the vertex  $x$  and the triangles  $\sigma_i (i = 1, \dots, 4)$ , and adding the 2-simplices  $\sigma' = x_1x_3x_2$  and  $\sigma'' = x_1x_3x_4$ . The map  $\varphi$  remains simplicial, since it sends  $\sigma', \sigma''$  to  $\delta', \delta''$ , respectively, contrary to the minimality choice of  $\mathcal{D}$ . This establishes that the degree of each interior vertex  $x$  of any minimal disk diagram is  $\geq 5$ .

Suppose now additionally that  $\mathcal{D}$  is a minimal disk diagram of  $C$  having a minimum number of inner vertices of degree 5. With some abuse of notation, we will denote the vertices of  $\mathcal{D}$  and their images in  $\mathbf{X}$  under  $\varphi$  by the same symbols. Let  $x$  be any interior vertex of  $\mathcal{D}$  of degree 5 and let  $x_1, \dots, x_5$  be the neighbors of  $x$ . If  $C = (x_1, x_2, x_3, x_4, x_5, x_1)$ , then we are done because  $\mathcal{D}$  is a 5-wheel. Now suppose that one of the edges of the 5-cycle  $(x_1, x_2, x_3, x_4, x_5, x_1)$ , say  $x_1x_2$ , belongs in  $\mathcal{D}$  to the second triangle  $x_1x_2x_6$ . The minimality of  $\mathcal{D}$  implies that  $x, x_1, x_2, x_3, x_4, x_5, x_6$  induce in  $\mathbf{X}$  a  $\widehat{W}_5$  or that  $x$  and  $x_6$  are adjacent in  $\mathbf{X}$ . In the first case, by the  $\widehat{W}_5$ -condition, there exists a vertex  $y$  of  $\mathbf{X}$  which is adjacent to all vertices of this  $\widehat{W}_5$ . Let  $\mathcal{D}'$  be a disk triangulation obtained from  $\mathcal{D}$  by deleting the vertex  $x$  and the five triangles incident to  $x$  as well as the triangle  $x_1x_2x_6$  and replacing them by the six triangles of the resulting 6-wheel centered at  $y$  (we call this operation a *flip*). In the second case, let  $\mathcal{D}'$  be a disk triangulation obtained from  $\mathcal{D}$  by deleting the the five triangles incident to  $x$  as well as the triangle  $x_1x_2x_6$  and replacing them by the six triangles of the resulting 6-wheel centered at  $x$ . In both cases, the resulting map  $\varphi$  remains simplicial.  $\mathcal{D}'$  has the same number of triangles as  $\mathcal{D}$ , therefore  $\mathcal{D}'$  is also a minimal disk diagram for  $C$ . The flip replaces in the first case the vertex  $x$  of degree 5 by the vertex  $y$  of degree 6. In the second case, it increases the degree of  $x$  from 5 to 6. In both cases, it preserves the degrees of all other vertices except the vertices  $x_1$  and  $x_2$ , whose degrees decrease by 1. Since, by the minimality choice of  $\mathcal{D}$ , the disk diagram  $\mathcal{D}'$  has at least as many inner vertices of degree 5 as  $\mathcal{D}$ , necessarily at least one of the vertices  $x_1, x_2$ , say  $x_1$ , is an inner vertex of degree at most 6 of  $\mathcal{D}$ . If the degree of  $x_1$  in  $\mathcal{D}$  is 5, then in  $\mathcal{D}'$  the degree of  $x_1$  will be 4, which is impossible by what has been shown above because  $\mathcal{D}'$  is also a minimal disk diagram and  $x_1$  is an interior vertex of  $\mathcal{D}'$ . Hence the degree of  $x_1$  in  $\mathcal{D}$  is 6 and its neighbors constitute an induced (in  $\mathcal{D}$ ) 6-cycle  $(x_6, x_2, x, x_5, u, v, x_6)$ .

*Case 1:*  $x$  and  $x_6$  are not adjacent in  $\mathbf{X}$ . Since  $\mathbf{X}$  does not contain induced  $C_4$  and the minimal disk diagrams of  $C$  do not contain interior vertices of degree 3 and 4, it can be easily

shown that the images in  $\mathbf{X}$  of the vertices  $x_5, y, x_6, v, u, x_1, x_4$  induce a  $\widehat{W}_5$  constituted by the 5-wheel centered at  $x_1$  and the triangle  $x_4yx_5$ . By the  $\widehat{W}_5$ -condition, there exists a vertex  $z$  of  $\mathbf{X}$  which is adjacent to all vertices of  $\widehat{W}_5$ . If  $z$  is adjacent in  $\mathbf{X}$  with all vertices of the 7-cycle  $(u, v, x_6, x_2, x_3, x_4, x_5, u)$ , then replacing in  $\mathcal{D}$  the 9 triangles incident to  $x$  and  $x_1$  by the 7 triangles of  $\mathbf{X}$  incident to  $z$ , we will obtain a disk diagram  $\mathcal{D}''$  for  $C$  having less triangles than  $\mathcal{D}$ , contrary to the minimality of  $\mathcal{D}$ . Therefore  $z$  is different from  $x$  and is not adjacent to one of the vertices  $x_2, x_3$ . Since  $x_1$  and  $x_4$  are not adjacent and both  $x$  and  $z$  are adjacent to  $x_1, x_4$ , to avoid an induced  $C_4$  we conclude that  $z$  is adjacent in  $\mathbf{X}$  to  $x$ . If  $z$  is not adjacent to  $x_2$ , then, since  $x$  and  $x_6$  are not adjacent, we will obtain a  $C_4$  induced by  $x, z, x_6, x_2$ . Thus  $z$  is adjacent to  $x_2$ , and therefore  $z$  is not adjacent to  $x_3$ . Since both  $z$  and  $x_3$  are adjacent to nonadjacent vertices  $x_2$  and  $x_4$ , we will obtain a  $C_4$  induced by  $z, x_2, x_3, x_4$ . This contradiction shows that the degree of  $x_1$  in  $\mathcal{D}$  is at least 7.

*Case 2:  $x$  and  $x_6$  are adjacent in  $\mathbf{X}$ .* Again, using the fact that the minimal disk diagrams for  $C$  do not contain interior vertices of degree 3 and 4, the fact that  $\mathbf{X}$  does not contain induced  $C_4$ , it can be easily shown that  $d(x, u) = 2$ . Therefore the vertices  $x_1, x_2, x_3, x_4, x_5, x, u$  induce a  $\widehat{W}_5$  constituted by the 5-wheel centered at  $x$  and the triangle  $x_1ux_5$ . Thus, by the  $\widehat{W}_5$ -condition, there exists a vertex  $y' \neq x$  containing  $\widehat{W}_5$  in its link. Then considering the minimal disk diagram obtained by the flip exchanging  $x$  and  $y'$  we conclude that the vertices  $u, v, x_6, x_2, y', x_1$  induce a 5-wheel. Together with the vertex  $x_3$  they induce a  $\widehat{W}_5$ , so that, by  $\widehat{W}_5$ -condition, there exists a vertex  $z$  adjacent to all the vertices  $u, v, x_6, x_2, y', x_1, x_3$ . If  $z$  is adjacent to  $x_4$  and  $x_5$  then we get a disk diagram for  $C$  having less triangles than  $\mathcal{D}$ , which contradicts the minimality of  $\mathcal{D}$ . If  $z$  is not adjacent to one of the vertices  $x_4, x_5$  then we also get a contradiction arguing as in Case 1. Therefore, in our case the degree of  $x_1$  in  $\mathcal{D}$  is also at least 7. This final contradiction shows that all interior vertices of  $\mathcal{D}$  have degrees  $\geq 6$ , establishing Claim 1.

From Claim 1 we deduce that any simple cycle  $C$  of the underlying graph of  $\mathbf{X}$  admits a minimal disk diagram  $\mathcal{D}$  which is either a 5-wheel or a systolic plane triangulation. We will call a *corner* of  $\mathcal{D}$  any vertex  $v$  of  $\partial\mathcal{D}$  which belongs in  $\mathcal{D}$  either to a unique triangle (first type) or to two triangles (second type). The corners of first type are the boundary vertices of degree two. The corners of second type are boundary vertices of degree three. In the first case, the two neighbors of  $v$  are adjacent. In the second case,  $v$  and its neighbors in  $\partial\mathcal{D}$  are adjacent to the third neighbor of  $v$ . From Gauss–Bonnet formula and Claim 1 we infer that  $\mathcal{D}$  contains at least three corners, and if  $\mathcal{D}$  has exactly three corners then they are all of first type. Furthermore, if  $\mathcal{D}$  contains four corners, then at least two of them are corners of first type.

Next we show that  $G(\mathbf{X})$  is weakly modular. To verify the triangle condition, pick three vertices  $u, v, w$  with  $1 = d(v, w) < d(u, v) = d(u, w) = k$ . We claim that if  $I(u, v) \cap I(u, w) = \{u\}$ , then  $k = 1$ . Suppose not. Pick two shortest paths  $P'$  and  $P''$  joining the pairs  $u, v$  and  $u, w$ , respectively, such that the cycle  $C$  composed of  $P', P''$  and the edge  $vw$  has minimal

$\text{Area}(C)$  (the choice of  $v, w$  implies that  $C$  is a simple cycle). Let  $\mathcal{D}$  be a minimal disk diagram of  $C$  satisfying Claim 1. Then either  $\mathcal{D}$  has a corner  $x$  different from  $u, v, w$  or the vertices  $u, v, w$  are the only corners of  $\mathcal{D}$ . In the second case,  $u, v, w$  are all three corners of first type, therefore the two neighbors of  $v$  in  $C$  will be adjacent. This means that  $w$  will be adjacent to the neighbor of  $v$  in  $P'$ , contrary to  $I(u, v) \cap I(u, w) = \{u\}$ . So, suppose that the corner  $x$  exists. Let  $x \in P'$ . Notice that  $x$  is a corner of second type, otherwise its neighbors  $y, z$  in  $P'$  are adjacent, contrary to the assumption that  $P'$  is a shortest path. Let  $p$  be the vertex of  $\mathcal{D}$  adjacent to  $x, y, z$ . If we replace in  $P'$  the vertex  $x$  by  $p$ , we will obtain a new shortest path between  $u$  and  $v$ . Together with  $P''$  and the edge  $vw$  this path forms a cycle  $C'$  whose area is strictly smaller than  $\text{Area}(C)$ , contrary to the choice of  $C$ . This establishes the triangle condition. As to the quadrangle condition, suppose by way of contradiction that we can find distinct vertices  $u, v, w, z$  such that  $v, w \in I(u, z)$  are neighbors of  $z$  and  $I(u, v) \cap I(u, w) = \{u\}$ , however  $u$  is not adjacent to  $v$  and  $w$ . Again, select two shortest paths  $P'$  and  $P''$  between  $u, v$  and  $u, w$ , respectively, so that the cycle  $C$  composed of  $P', P''$  and the edges  $vz$  and  $zw$  has minimum area. Choose a minimal disk  $\mathcal{D}$  of  $C$  as in Claim 1. From the initial hypothesis concerning the vertices  $u, v, w, z$  we deduce that  $\mathcal{D}$  has at most one corner of first type located at  $u$ . Hence  $\mathcal{D}$  contains at least four corners of second type. Since one corner  $x$  is distinct from  $u, v, w, z$ , then proceeding in the same way as before, we will obtain a contradiction with the choice of the paths  $P', P''$ . This shows that  $u$  is adjacent to  $v, w$ , establishing the quadrangle condition. Hence  $G(\mathbf{X})$  is a weakly modular graph without induced  $C_4$ , concluding the proof of the implication (vii) $\Rightarrow$ (iv) and of the theorem.  $\square$

In the analysis of his construction of locally homogeneous graphs  $H$  having a given regular graph of girth  $\geq 6$  (i.e., 6-large) as a link of each vertex of  $H$ , Weetman [Wee94] introduced *quasitrees* as the graphs  $G = (V, E)$  satisfying the following two conditions for each vertex  $v$ : (F1) each vertex  $x \in S_{i+1}(v)$  has one or two adjacent neighbors in  $S_i(v)$ ; (F2) any two adjacent vertices  $x, y \in S_{i+1}(v)$  have a common neighbor  $z \in S_i(v)$ . It can be easily seen that (F2) is a reformulation of the edge condition (E) (alias the triangle condition). On the other hand, (F1) is a particular case of the vertex condition (V). From Theorem 3.1(ii) we immediately obtain the following observation:

**Corollary 3.2.** *The simplicial complexes derived from quasitrees are weakly systolic. In particular, quasitrees are weakly bridged graphs.*

The 5-wheel is an example of a quasitree which is not a bridged graph, thus not all simplicial complexes derived from quasitrees are systolic.

#### 4. PROPERTIES OF WEAKLY SYSTOLIC COMPLEXES

In this section, we establish some further combinatorial and geometrical properties of weakly systolic complexes, which are well known for systolic complexes [Hag03, JS06]. In particular, we show that weakly systolic complexes  $\mathbf{X}$  satisfy the  $SD_n(\sigma^*)$  property for facets  $\sigma^*$  (maximal by inclusion simplices of  $\mathbf{X}$ ).

**Lemma 4.1** (Edges descend on balls). *Let  $\sigma$  be a simplex of a weakly systolic complex  $\mathbf{X}$ . Let  $e = zz'$  be an edge contained in the sphere  $S_i(\sigma)$ . Then there exists a vertex  $w \in \sigma$  and a vertex  $v \in B_{i-1}(\sigma)$  such that  $v$  is adjacent to  $z, z'$  and  $d(v, w) = i - 1$ .*

*Proof.* If there exists a vertex  $w \in \sigma$  such that  $z, z' \in S_i(w)$  then the assertion follows from the  $SD_n$  property. Therefore suppose that such a vertex of  $\mathbf{X}$  does not exist. Let  $w, w'$  be two vertices of  $\sigma$  with  $d(w, z) = d(w', z') = i$ . Since  $d(w', z) = d(w, z') = i + 1$ , we conclude that  $z \in I(w, z')$ . Since  $w, z' \in B_i(w')$  and  $z \notin B_i(w')$ , this contradicts the convexity of  $B_i(w')$ .  $\square$

**Lemma 4.2** (Big balls are convex). *Let  $\sigma$  be a simplex of a weakly systolic complex  $\mathbf{X}$  and let  $i \geq 2$ . Then the ball  $B_i(\sigma)$  is convex. In particular,  $B_i(\sigma) \cap N(v)$  is a simplex for any vertex  $z \in S_{i+1}(\sigma)$ .*

*Proof.* Since  $G(\mathbf{X})$  is weakly modular and  $B_i(\sigma)$  induces a connected subgraph, according to [Che89] it suffices to show that  $B_i(\sigma)$  is locally convex, i.e., if  $x, y \in B_i(\sigma)$ ,  $d(x, y) = 2$ , and  $z$  is a common neighbor of  $x$  and  $y$ , then  $z \in B_i(\sigma)$ . Suppose by way of contradiction that  $z \in S_{i+1}(\sigma)$ . Let  $u$  and  $v$  be vertices of  $\sigma$  located at distance  $i$  from  $x$  and  $y$ , respectively. If  $u = v$ , then  $x, y \in I(u, z)$ , thus  $x$  and  $y$  must be adjacent because  $G(\mathbf{X})$  is thin. So, suppose that  $u \neq v$ , i.e.,  $d(y, u) = d(z, u) = i + 1$  holds. By triangle condition there exists a common neighbor  $w$  of  $z$  and  $y$  having distance  $i$  to  $u$ . Since  $x, w \in I(z, u)$  and  $G(\mathbf{X})$  is thin, the vertices  $x$  and  $w$  are adjacent; moreover by triangle condition, there exists a common neighbor  $u'$  of  $w$  and  $x$  having distance  $i - 1$  to  $u$ . If  $d(w, v) = i + 1$ , then  $y, u' \in I(w, v)$ , thus  $y$  and  $u'$  must be adjacent because  $G(\mathbf{X})$  is thin. As a result, we obtain a 4-cycle defined by  $x, z, y, u'$ . Since  $d(z, u) = i + 1$  and  $d(u', u) = i - 1$ ,  $z$  and  $u'$  cannot be adjacent, thus this 4-cycle must be induced, which is impossible. Hence  $d(w, v) = i$ . Let  $u''$  be a neighbor of  $u$  in the interval  $I(u, u')$  (it is possible that  $u'' = u'$ ). Since  $d(y, u) = i + 1$  and  $d(u', u) = i - 1$ ,  $d(u', y) = 2$ , we conclude that  $u' \in I(y, u)$ , yielding  $u'' \in I(u, u') \subset I(u, y)$ . Since  $v$  also belongs to  $I(u, y)$  and  $G(\mathbf{X})$  is thin, the vertices  $u''$  and  $v$  must be adjacent. But in this case  $d(x, v) = 1 + d(u', u'') + 1 = i$ , contrary to the assumption that  $d(x, v) = i + 1$ . This contradiction shows that  $B_i(\sigma)$  is convex for any  $i \geq 2$ .  $\square$

**Proposition 4.3** ( $SD_n$  property for maximal simplices). *A weakly systolic complex satisfies the property  $SD_n(\sigma^*)$  for any maximal simplex  $\sigma^*$  of  $\mathbf{X}$ .*

*Proof.* Let  $\sigma$  be a simplex of  $\mathbf{X}$  located in the sphere  $S_{i+1}(\sigma^*)$ . For each vertex  $v \in \sigma$  denote by  $\sigma^*(v)$  the metric projection of  $v$  in  $\sigma^*$ , i.e., the set of all vertices of  $\sigma^*$  located at distance  $i + 1$  from  $v$ . Notice that the sets  $\sigma^*(v)$  ( $v \in \sigma$ ) can be linearly ordered by inclusion. Indeed, if we suppose the contrary, then there exist two vertices  $v', v'' \in \sigma$  and the vertices  $u' \in \sigma^*(v') \setminus \sigma^*(v'')$  and  $u'' \in \sigma^*(v'') \setminus \sigma^*(v')$ . This is impossible because in this case  $v'' \in I(v', u'') \setminus B_{i+1}(u')$  and  $v', u'' \in B_{i+1}(u')$ , contrary to the convexity of  $B_{i+1}(u')$ . Therefore the simplices  $\sigma^*(v)$  ( $v \in \sigma$ ) can be linearly ordered by inclusion. This means that  $\sigma \subset S_{i+1}(u)$  holds for any vertex  $u$  belonging to all metric projections  $\sigma_0^* = \bigcap \{\sigma^*(v) : v \in \sigma\}$ .

Applying the  $SD_n(u)$  property to  $\sigma$  we conclude that the set of all vertices  $x \in S_i(u) \subseteq S_i(\sigma^*)$  adjacent to all vertices of  $\sigma$  is a non-empty simplex. Pick two vertices  $x, y \in S_i(\sigma^*)$  adjacent to all vertices of  $\sigma$ . Let  $x \in S_i(u)$  and  $y \in S_i(w)$  for  $u, w \in \sigma_0^*$ . We assert that  $x$  and  $y$  are adjacent. Let  $v$  be a vertex of  $\sigma$  whose projection  $\sigma^*(v)$  is maximal by inclusion. If  $\sigma^*(v) = \sigma^*$ , then applying the  $SD_n(v)$  property we conclude that there exists a vertex  $v'$  at distance  $i$  to  $v$  and adjacent to all vertices of  $\sigma^*$ , contrary to maximality of  $\sigma^*$ . Hence  $\sigma^*(v)$  is a proper simplex of  $\sigma^*$ . Let  $s \in \sigma^* \setminus \sigma^*(v)$ . Then  $x, y \in I(v, s)$  and since  $G(\mathbf{X})$  is thin, the vertices  $x$  and  $y$  must be adjacent.  $\square$

We conclude this section by showing that the systolic complexes are exactly the flag complexes satisfying  $SD_n(\sigma^*)$  for all simplices  $\sigma^*$ .

**Proposition 4.4.** *A simplicial flag complex  $\mathbf{X}$  is systolic if and only if  $\mathbf{X}$  satisfies the property  $SD_n(\sigma^*)$  for all simplices  $\sigma^*$  of  $\mathbf{X}$  and all  $n \geq 0$ .*

*Proof.* If  $\mathbf{X}$  satisfies the property  $SD_n(v)$  for all vertices, then  $\mathbf{X}$  is weakly systolic by Theorem 3.1. Since  $\mathbf{X}$  satisfies the property  $SD_n(e)$  for all edges,  $\mathbf{X}$  does not contain 5-wheels. Hence  $\mathbf{X}$  is systolic. Conversely, let  $\sigma^*$  be an arbitrary simplex of a systolic complex  $\mathbf{X}$  and let  $\sigma$  be a simplex belonging to  $S_{i+1}(\sigma^*)$ . Since  $B_i(\sigma^*)$  is convex because  $G(\mathbf{X})$  is bridged, the set  $\sigma_0$  of all vertices  $x \in B_i(\sigma^*)$  such that  $\sigma \cup \{x\} \in \mathbf{X}$ , if nonempty, necessarily is a simplex. Thus it suffices to show that  $\sigma_0 \neq \emptyset$ . As in previous proof, for each vertex  $v \in \sigma$  denote by  $\sigma^*(v)$  the metric projection of  $v$  in  $\sigma^*$ . Then, as we showed in the proof of Proposition 4.3, the sets  $\sigma^*(v)$  can be linearly ordered by inclusion. Therefore there exists a vertex  $u \in \sigma^*$  belonging to all projections  $\sigma^*(v), v \in \sigma$ . Then  $\sigma \subset S_{i+1}(u)$ , whence  $\sigma_0$  is nonempty because of the  $SD_n(u)$  property.  $\square$

## 5. DISMANTLABILITY OF WEAKLY BRIDGED GRAPHS

In this section, we show that the underlying graphs of weakly systolic complexes are dismantlable and that a dismantling order can be obtained using LexBFS. Then we use this result to deduce several consequences about combings of weakly bridged graphs and about the collapsibility of weakly systolic complexes. Other consequences of dismantling are given in subsequent sections.

**Theorem 5.1** (LexBFS dismantlability). *Any LexBFS ordering of a locally finite weakly bridged graph  $G = G(\mathbf{X})$  is a dismantling ordering. In particular, locally finite weakly systolic complexes  $\mathbf{X}$  and their Rips complexes  $\mathbf{X}_k$  are LC-contractible and therefore collapsible.*

*Proof.* We will establish the result for finite weakly bridged graphs and finite weakly systolic complexes. The proof in the locally finite case is completely similar. Let  $v_n, \dots, v_1$  be the total order returned by the LexBFS starting from the basepoint  $u = v_n$ . Let  $G_i$  be the subgraph of  $G$  induced by the vertices  $v_n, \dots, v_i$ . For a vertex  $v \neq u$  of  $G$ , denote by  $f(v)$  its father in the LexBFS tree  $T_u$ , by  $L(v)$  the list of all neighbors of  $v$  labeled before  $v$ , and by  $\alpha(v)$  the label of  $v$  (i.e., if  $v = v_i$ , then  $\alpha(v) = i$ ). We decompose the label  $L(v)$  of each

vertex  $v$  into two parts  $L'(v)$  and  $L''(v)$  : if  $d(v, u) = i$ , then  $L'(v) = L(v) \cap S_{i-1}(u)$  and  $L''(v) = L(v) \cap S_i(u)$ . Notice that in the lexicographic order of  $L(v)$ , all vertices of  $L'(v)$  precede the vertices of  $L''(v)$ ; in particular, the father of  $v$  belongs to  $L'(v)$ . The proof of the theorem is a consequence of the following assertion, which we call the *fellow traveler property*:

**Fellow Traveler Property:** *If  $v, w$  are adjacent vertices of  $G$ , then their fathers  $v' = f(v)$  and  $w' = f(w)$  either coincide or are adjacent. If  $v'$  and  $w'$  are adjacent and  $\alpha(w) < \alpha(v)$ , then  $w'$  is adjacent to  $v$  and  $v'$  is not adjacent to  $w$ .*

Indeed, if this assertion holds, then we claim that  $v_n, \dots, v_1$  is a dismantling order. To see this, it suffices to show that any vertex  $v_i$  is dominated in  $G_i$  by its father  $f(v_i)$  in the LexBFS tree  $T_u$ . Pick any neighbor  $v_j$  of  $v_i$  in  $G_i$ . We assert that  $v_j$  coincides or is adjacent to  $f(v_i)$ . This is obviously true if  $f(v_j) = f(v_i)$ . Otherwise, if  $f(v_i) \neq f(v_j)$ , then the Fellow Traveler Property implies that  $f(v_i)$  and  $f(v_j)$  are adjacent and since  $i < j$  that  $v_j$  is adjacent to  $f(v_i)$ . This shows that indeed any LexBFS order is a dismantling order.

We will establish now the Fellow Traveler Property by induction on  $i + 1 := \max\{d(u, v), d(u, w)\}$ . First suppose that  $d(u, v) < d(u, w)$ . Then  $v, w' \in I(w, u)$  and since  $G$  is thin, we conclude that  $v$  and  $w'$  either coincide or are adjacent. In the first case we are done because  $v$  (and therefore  $w'$ ) is adjacent to its father  $v' = f(v)$ . If  $v$  and  $w'$  are adjacent, since  $i = d(u, v) = d(u, w')$ , the vertices  $v'$  and  $f(w')$  coincide or are adjacent by the induction assumption. Again, if  $v' = f(w')$ , the assertion is immediate. Now suppose that  $v'$  and  $f(w')$  are adjacent. Since  $w' = f(w)$  was labeled before  $v$  (otherwise the father of  $w$  is  $v$  and not  $w'$ ),  $f(w')$  must be labeled before  $v'$ , therefore by the induction hypothesis we deduce that  $v' = f(v)$  must be adjacent to  $w' = f(w)$ . This concludes the analysis of the case  $d(u, v) < d(u, w)$ .

From now on, suppose that  $d(u, v) = d(u, w) = i + 1$  and that  $\alpha(w) < \alpha(v)$ . If the vertices  $v' = f(v)$  and  $w' = f(w)$  coincide, then we are done. If the vertices  $v'$  and  $w'$  are adjacent, then the vertices  $v, w, w', v'$  define a 4-cycle, which cannot be induced by the  $SD_n$  property (see Theorem 3.1). Since  $v$  was labeled before  $w$ , the vertex  $v'$  must be labeled before  $w'$ . Therefore, if  $v'$  is adjacent to  $w$ , then LexBFS will label  $w$  from  $v'$  and not from  $w'$ , a contradiction. Thus  $v'$  and  $w$  are not adjacent, showing that  $w'$  must be adjacent to  $v$ , establishing the required assertion. So, assume by way of contradiction that the vertices  $v'$  and  $w'$  are not adjacent in  $G$ . Then  $w'$  is not adjacent to  $v$ , otherwise  $w', v' \in B_i(u)$  and  $v \in I(v', w') \cap S_{i+1}(u)$ , contrary to the convexity of the ball  $B_i(u)$  (similarly,  $v'$  is not adjacent to  $w$ ).

Since  $G$  is weakly modular by Theorem 3.1, by triangle condition applied to the vertices  $v, w$ , and  $u$ , there exists a common neighbor  $s$  of  $v$  and  $w$  located at distance  $i$  from  $u$ . Denote by  $S$  the set of all such vertices  $s$ . From the property  $SD_n$  we infer that  $S$  is a simplex of  $\mathbf{X}$  (i.e., its vertices are pairwise adjacent in  $G$ ). Since  $v'$  and  $w'$  do not belong to  $S$ , necessarily all vertices of  $S$  have been labeled later than  $v'$  and  $w'$  (but obviously before  $v$  and  $w$ ). Pick a vertex  $s$  in  $S$  with the largest label  $\alpha(s)$  and set  $z := f(s)$ . By induction assumption applied

to the pairs of adjacent vertices  $\{v', s\}$  and  $\{s, w'\}$ , we conclude that the vertices of each of the pairs  $\{f(v'), z\}$  and  $\{z, f(w')\}$  either coincide or are adjacent. Moreover, in all cases, the vertex  $z$  must be adjacent to the vertices  $v'$  and  $w'$ .

**Claim 1:**  $C := L'(v') = L'(s) = L'(w')$  and  $z$  is the father of  $v'$  and  $w'$ .

**Proof of Claim 1:** Since  $s$  was labeled later than  $v'$  and  $w'$ , it suffices to show that  $L'(v') = L'(s)$ . Indeed, if this is the case, then necessarily  $z$  is the father of  $v'$ . Then, as  $z$  is adjacent to  $w'$  and  $\alpha(w') < \alpha(v')$ , necessarily  $z$  is also the father of  $w'$ . Now, if  $L'(w')$  and  $L'(s) = L'(v')$  do not coincide, since  $L'(v')$  lexicographically precedes  $L''(v')$  and  $L'(w')$  precedes  $L''(w')$ , the fact that LexBFS labeled  $v'$  before  $w'$  means that  $L'(v')$  lexicographically precedes  $L'(w')$ . Since  $L'(s) = L'(v')$ , then necessarily LexBFS would label  $s$  before  $w'$ , a contradiction. This shows that  $L'(s) = L'(v')$  implies the equality of the three labels  $L'(v')$ ,  $L'(s)$ , and  $L'(w')$ .

To show that  $L'(v') = L'(s)$ , since  $\alpha(s) < \alpha(v')$ , it suffices to establish only the inclusion  $L'(v') \subseteq L'(s)$ . Suppose by way of contradiction that there exists a vertex in  $L'(v') \setminus L'(s)$  i.e., a vertex  $x \in S_{i-1}(u)$  which is adjacent to  $v'$  but is not adjacent to  $s$ . Let  $x$  be the vertex of  $L'(v') \setminus L'(s)$  having the largest label  $\alpha(x)$ . Since  $s$  was labeled by LexBFS later than  $v'$ , necessarily any vertex of  $L'(s) \setminus L'(v')$  must be labeled later than  $x$ . Notice that  $x$  cannot be adjacent to  $w'$ , otherwise we obtain an induced 4-cycle formed by the vertices  $v', s, w', x$ . Since  $x$  is not adjacent to  $v, w$ , and  $s$ , we conclude that the vertices  $v, w, w', z, v', s, x$  induce an extended 5-wheel. By the  $\widehat{W}_5$ -condition, there exists a vertex  $t$  adjacent to all vertices of this  $\widehat{W}_5$ . Hence  $t \in S$ . Further  $t$  must be adjacent to  $z$ , otherwise we obtain a forbidden 4-cycle induced by the vertices  $s, z, x$ , and  $t$ . For the same reason,  $t$  must be adjacent to any other vertex  $z'$  belonging to  $L'(v') \cap L'(s)$ . This means that LexBFS will label  $t$  before  $s$ . Since  $t$  belongs to  $S$  and  $\alpha(t) > \alpha(s)$ , we obtain a contradiction with the choice of the vertex  $s$ . This contradiction concludes the proof of the Claim 1.

Since  $v'$  and  $w'$  are not adjacent and  $G$  does not contain induced 4-cycles, any vertex  $s' \neq s$  adjacent to  $v'$  and  $w'$  is also adjacent to  $s$ . In particular, this shows that  $L''(v') \cap L''(w') \subseteq L''(s)$ . Therefore, if  $L''(w') \subseteq L''(v')$ , then  $L''(w') \subseteq L''(s)$ . Since  $v' \in L''(s) \setminus L''(w')$  and  $L'(s) = L'(w')$  by Claim 1, we conclude that the vertex  $s$  must be labeled before  $w'$ , contrary to the assumption that  $\alpha(s) < \alpha(w')$ . Therefore the set  $B := L''(w') \setminus L''(v')$  is nonempty. Since  $v'$  was labeled before  $w'$  and  $L'(v') = L'(w')$  by Claim 1, we conclude that the set  $A := L''(v') \setminus L''(w')$  is nonempty as well. Let  $p$  be the vertex of  $A$  with the largest label  $\alpha(p)$  and let  $q$  be the vertex of  $B$  with the largest label  $\alpha(q)$ . Since LexBFS labeled  $v'$  before  $w'$  and  $L'(v') = L'(w')$  holds, necessarily  $\alpha(q) < \alpha(p)$  holds. Since  $p \in L''(v')$ , we obtain that  $\alpha(w') < \alpha(v') < \alpha(p)$ . Since  $v' = f(v)$  and  $w' = f(w)$ , this shows that  $p$  cannot be adjacent to the vertices  $v$  and  $w$ . If  $s$  is adjacent to  $p$ , then  $p \in L''(s)$ . But then from Claim 1 and the inclusion  $L''(v') \cap L''(w') \subseteq L''(s)$  we will infer that LexBFS must label  $s$  before  $w'$ , contrary to the assumption that  $\alpha(s) < \alpha(w')$ . Therefore  $p$  is not adjacent to  $s$  either. On the other hand, since  $\alpha(v') < \alpha(p)$ , by the induction hypothesis applied to the adjacent vertices

$p$  and  $v'$ , we infer that  $z = f(v')$  must be adjacent to  $p$ . Hence the vertices  $v, w, w', z, v', s, p$  induce an extended 5-wheel. By the  $\widehat{W}_5$ -condition, there exists a vertex  $t$  adjacent to all these vertices. Since  $C := L'(v') = L'(w')$  and  $d(v', w') = 2$ , to avoid induced 4-cycles, the vertex  $t$  must be adjacent to any vertex of  $C$ . For the same reason,  $t$  must be adjacent to any vertex of  $L''(v') \cap L''(w')$ . Since additionally  $t$  is adjacent to the vertex  $p$  of  $A$  with the highest label, necessarily  $t$  will be labeled by LexBFS before  $w'$  and  $s$ . Since  $t$  is adjacent to  $v$  and  $w$ , this contradicts the assumption that  $w' = f(w)$ . This shows that the initial assumption that  $v'$  and  $w'$  are not adjacent lead to a final contradiction. Hence the order returned by LexBFS is indeed a dismantling order of the weakly bridged graph  $G = G(\mathbf{X})$ .

To show that any finite weakly systolic complex  $\mathbf{X}$  is LC-contractible it suffices to notice that, since  $\mathbf{X}$  is a flag complex, the LC-contractibility of  $\mathbf{X}$  is equivalent to the dismantlability of its graph  $G(\mathbf{X})$ , and hence the result follows from the first part of the theorem. To show that the Rips complex  $\mathbf{X}_k$  is LC-contractible, since  $\mathbf{X}_k$  is a flag complex, it suffices to show that its graph  $G(\mathbf{X}_k)$  is dismantlable. From the definition of  $\mathbf{X}_k$ , the graph  $G(\mathbf{X}_k)$  coincides with the  $k$ th power  $G^k$  of the underlying graph  $G$  of  $\mathbf{X}$ . Now notice that if a vertex  $v$  is dominated in  $G$  by a vertex  $u$ , then  $u$  also dominates  $v$  in the graph  $G^k$ . Indeed, pick any vertex  $x$  adjacent to  $v$  in  $G^k$ . Then  $d(v, x) \leq k$  in  $G$ . Let  $y$  be the neighbor of  $v$  on some shortest path  $P$  connecting the vertices  $v$  and  $x$  in  $G$ . Since  $u$  dominates  $v$ , necessarily  $u$  is adjacent to  $y$ . Hence  $d(u, x) \leq k$  in  $G$ , therefore  $u$  is adjacent to  $x$  in  $G^k$ . This shows that  $v$  is dominated by  $u$  in the graph  $G^k$  as well. Therefore the dismantling order of  $G$  returned by LexBFS is also a dismantling order of  $G^k$ , establishing that the Rips complex  $\mathbf{X}_k$  is LC-contractible. This completes the proof of the theorem.  $\square$

**Remark 5.2.** BFS orderings of weakly bridged graphs do not satisfy the property that each vertex is dominated by its father. For example, let  $G$  be a 5-wheel whose vertices are labeled as in Fig. 1. If BFS starts from the vertex  $x_1$  and orders the remaining vertices as  $x_2, x_5, c, x_3, x_4$ , then the father of the last vertex  $x_4$  is  $x_5$ , however  $x_5$  does not dominate  $x_4$  in the whole graph. On the other hand, LexBFS starting from  $x_1$  and continuing with  $x_2$ , necessarily will label the vertex  $c$  next. As a consequence,  $c$  will be the father of the last labeled vertex  $x_4$  and obviously  $c$  dominates  $x_4$ . Nevertheless, the order  $x_1, x_2, x_5, c, x_3, x_4$  returned by BFS is a domination order of the 5-wheel. Is this true for all weakly bridged graphs?

**Corollary 5.3.** *Graphs of Rips complexes  $\mathbf{X}_n$  of locally finite systolic and weakly systolic complexes are dismantlable.*

**Corollary 5.4.** *Finite weakly bridged graphs are cop-win.*

For a locally finite weakly bridged graph  $G$  and integer  $k$  denote by  $G_k$  the subgraph of  $G$  induced by the first  $k$  labeled vertices in a LexBFS order, i.e., by the vertices of  $G$  with  $k$  lexicographically largest labels.

**Corollary 5.5.** *Any  $G_k$  is an isometric weakly bridged subgraph of  $G$ .*

*Proof.* By Theorem 5.1, LexBFS returns a dismantling order of  $G$ , hence any  $G_k$  is an isometric subgraph of  $G$ . Therefore  $G_k$  is a thin graph, because any interval  $I(x, y)$  in  $G_k$  is contained in the interval of  $G$  between  $x$  and  $y$ . Moreover,  $G_k$ , as an isometric subgraph of a  $G$ , does not contain isometric cycles of length  $> 5$ . Hence, by a result of [SC83, FJ87],  $G_k$  is a graph with convex balls. By Theorem 3.1(vi) it remains to show that any induced 5-cycle  $C$  of  $G_k$  is included in a 5-wheel. Suppose by induction assumption that this is true for  $G_{k-1}$ . Therefore  $C$  must contain the last labeled vertex of  $G_k$ , denote this vertex by  $v$ . Let  $x$  and  $y$  be the neighbors of  $v$  in  $C$ . Let  $v' = f(v)$  be the vertex (of  $G_k$ ) dominating  $v$  in  $G_k$ . Since  $C$  is induced, necessarily  $v'$  is adjacent to  $x$  and  $y$  but different from these vertices. Denote by  $C'$  the 5-cycle obtained by replacing in  $C$  the vertex  $v$  by  $v'$ . If  $C'$  is not induced, then  $v'$  will be adjacent to a third vertex of  $C$ , and since  $G_k$  does not contain induced 4-cycles,  $v'$  will be adjacent to all vertices of  $C$ , showing that  $C$  extends to a 5-wheel. So, suppose that  $C'$  is induced. Applying the induction hypothesis to  $G_{k-1}$ , we conclude that  $C'$  extends to a 5-wheel in  $G_{k-1}$ . Let  $w$  be the central vertex of this wheel. To avoid a 4-cycle induced by the vertices  $x, y, v$ , and  $w$ , necessarily  $v$  and  $w$  must be adjacent. Hence  $C$  extends in  $G_k$  to a 5-wheel centered at  $w$ . This establishes that indeed  $G_k$  is weakly bridged.  $\square$

A *homomorphism* of a graph  $G = (V, E)$  to itself is a mapping  $\varphi : V \rightarrow V$  such that for any edge  $uv \in E$  we have  $\varphi(u)\varphi(v) \in E$  or  $\varphi(u) = \varphi(v)$ . A set  $S \subset V$  is fixed by  $\varphi$ , if  $\varphi(S) = S$ . A *simplicial map* on a simplicial complex  $\mathbf{X}$  is a map  $\varphi : V(\mathbf{X}) \rightarrow V(\mathbf{X})$  such that for all  $\sigma \in \mathbf{X}$  we have  $\varphi(\sigma) \in \mathbf{X}$ . A simplicial map fixes a simplex  $\sigma \in \mathbf{X}$  if  $\varphi(\sigma) = \sigma$ . Every simplicial map on  $\mathbf{X}$  is a homomorphism of its graph  $G(\mathbf{X})$ . Every homomorphism of a graph  $G$  is a simplicial map on its clique complex  $\mathbf{X}(G)$ . Therefore, if  $\mathbf{X}$  is a flag complex, then the set of simplicial maps of  $\mathbf{X}$  coincides with the set of homomorphisms of its graph  $G(\mathbf{X})$ . It is well known (see, for example, [HN04, Theorem 2.65]) that any homomorphism of a finite dismantlable graph to itself fixes some clique. From Theorem 5.1 we know that the graphs of weakly systolic complexes as well as the graphs of their Rips complexes are dismantlable. Therefore from preceding discussion we obtain:

**Corollary 5.6.** *Any homomorphism of a finite weakly bridged graph  $G = G(\mathbf{X})$  to itself fixes some clique. Any simplicial map of a weakly systolic complex  $\mathbf{X}$  to itself or of its Rips complex  $\mathbf{X}_k$  to itself fixes some simplex of respective complex.*

Let  $u$  be a base point of a graph  $G$ . A (*geodesic*) *k-combing* [ECH<sup>+</sup>92] is a choice of a shortest path  $P_{(u,x)}$  between  $u$  and each vertex  $x$  of  $G$ , such that  $P := P_{(u,v)}$  and  $Q := P_{(u,w)}$  are  $k$ -fellow travelers for any adjacent vertices  $v$  and  $w$  of  $G$ , i.e.,  $d(P(t), Q(t)) \leq k$  for all integers  $t \geq 0$ . One can imagine the union of combing paths as a spanning tree  $T_u$  of  $G$  rooted at  $u$  and preserving the distances from  $u$  to all vertices. A natural way to comb a graph  $G$  from  $u$  is to run BFS and to take as a shortest path  $P_{(u,x)}$  the unique path of the BFS-tree  $T_u$ . It is shown in [Che00] that for bridged graphs this geodesic combing satisfies the 1-fellow

traveler property. We will show now that in the case of weakly bridged graphs the same combing property is satisfied by the paths of any LexBFS tree  $T_u$  :

**Corollary 5.7.** *Locally finite weakly bridged graphs  $G$  admit a geodesic 1-combing defined by the paths of any LexBFS tree  $T_u$  of  $G$ .*

*Proof.* Pick two adjacent vertices  $v, w$  of  $G$  and suppose that  $w$  was labeled by LexBFS after  $v$ . Then  $d(u, v) \leq d(u, w) = n$ . We proceed by induction on  $n$ . Denote by  $v' = f(v)$  and  $w' = f(w)$  the fathers of  $v$  and  $w$ . By definition of the combing,  $v'$  and  $w'$  are the neighbors of  $v$  and  $w$  in the combings paths  $P_{(u,v)}$  and  $P_{(u,w)}$ , respectively. If  $d(u, v) = d(u, w)$ , then the fellow traveler property established in Theorem 5.1 shows that  $v'$  and  $w'$  either are adjacent or coincide. In the second case,  $P_{(u,v)}$  and  $P_{(u,w)}$  coincide everywhere except the last vertices  $v$  and  $w$ . In the first case, since  $d(u, v') = d(u, w') = n - 1$ , the paths  $P_{(u,v')}$  and  $P_{(u,w')}$  are 1-fellow travelers. Since  $P_{(u,v')} \subset P_{(u,v)}$  and  $P_{(u,w')} \subset P_{(u,w)}$  we conclude that  $P_{(u,v)}$  and  $P_{(u,w)}$  are 1-fellow travelers as well. Now suppose that  $d(u, v) < d(u, w)$ . If  $w' = v$ , then  $P_{(u,v)} \subset P_{(u,w)}$  and we are done. Otherwise,  $w'$  is adjacent to  $v$  and  $v'$ . Applying the induction hypothesis to the combing paths  $P_{(u,v')} \subset P_{(u,v)}$  and  $P_{(u,w')} \subset P_{(u,w)}$ , again we conclude that  $P_{(u,v)}$  and  $P_{(u,w)}$  are 1-fellow travelers.  $\square$

## 6. FIXED POINT THEOREM

In this section, we establish the fixed point theorem (Theorem C from Introduction). We start with two auxiliary results. The first is an easy corollary of Theorem 5.1:

**Lemma 6.1** (Strictly dominated vertex). *Let  $\mathbf{X}$  be a finite weakly systolic complex. Then either  $\mathbf{X}$  is a single simplex or it contains two vertices  $v, w$  such that  $B_1(v)$  is a proper subset of  $B_1(w)$ , i.e.  $B_1(v) \subsetneq B_1(w)$ .*

*Proof.* Let  $v$  be the last vertex of  $\mathbf{X}$  labeled by LexBFS which started at vertex  $u$  (see Theorem 5.1). If  $d(u, v) = 1$ , then the construction of our ordering implies that  $B_1(u) = V(\mathbf{X})$ . Hence, either there exists a vertex  $w$ , such that  $B_1(w) \subset V(\mathbf{X}) = B_1(u)$ , and we are done, or every two vertices of  $\mathbf{X}$  are adjacent, i.e.,  $\mathbf{X}$  is a simplex. Now suppose that  $d(u, v) \geq 2$ . From Theorem 5.1 we know that  $B_1(v) \subseteq B_1(w)$ , where  $w$  is the father of  $v$ . Since  $d(u, v) = d(u, w) + 1 \geq 2$ , we conclude that  $u \neq w$  and that  $z \in B_1(w) \setminus B_1(v)$ , where  $z$  is the father of  $w$ . Hence  $B_1(v)$  is a proper subset of  $B_1(w)$ .  $\square$

**Lemma 6.2** (Elementary LC-reduction). *Let  $\mathbf{X}$  be a finite weakly systolic complex. Let  $v, w$  be two vertices such that  $B_1(v)$  is a proper subset of  $B_1(w)$ . Then the full subcomplex  $\mathbf{X}_0$  of  $\mathbf{X}$  spanned by all vertices of  $\mathbf{X}$  except  $v$  is weakly systolic.*

*Proof.* It is easy to see that  $\mathbf{X}_0$  is simply connected (see also the discussion in Section 2.3). Thus, by Theorem 3.1, it suffices to show that  $\mathbf{X}_0$  does not contain induced 4-cycles and satisfies the  $\widehat{W}_5$ -condition. Since, by Theorem 3.1,  $\mathbf{X}$  does not contain induced  $C_4$ , the same is true for its full subcomplex  $\mathbf{X}_0$ . Let  $\widehat{W}_5 \subseteq \mathbf{X}_0$  be a given 5-wheel plus a triangle as defined

in Section 3. By Theorem 3.1 there exists a vertex  $v' \in \mathbf{X}$  adjacent in  $\mathbf{X}$  to all vertices of  $\widehat{W}_5$ . If  $v' \neq v$  then  $v' \in \mathbf{X}_0$  and if  $v' = v$  then  $\widehat{W}_5 \subseteq \text{lk}(v, \mathbf{X}_0)$ . In both cases all vertices of  $\widehat{W}_5$  are adjacent to a vertex of  $\mathbf{X}_0$ . Thus  $\mathbf{X}_0$  also satisfies the  $\widehat{W}_5$ -condition and hence the lemma follows.  $\square$

**Theorem 6.3** (The fixed point theorem). *Let  $G$  be a finite group acting by simplicial automorphisms on a locally finite weakly systolic complex  $\mathbf{X}$ . Then there exists a simplex  $\sigma \in \mathbf{X}$  which is invariant under the action of  $G$ .*

*Proof.* Let  $\mathbf{X}'$  be the subcomplex of  $\mathbf{X}$  spanned by the convex hull of the set  $\{gv : g \in G\}$ . Then it is clear that  $\mathbf{X}'$  is a bounded and  $G$ -invariant full subcomplex of  $\mathbf{X}$ . Moreover, as a convex subcomplex of a weakly systolic complex,  $\mathbf{X}'$  is itself weakly systolic. Thus there exists a minimal non-empty  $G$ -invariant subcomplex  $\mathbf{X}_0$  of  $\mathbf{X}$ , that is itself weakly systolic. Since  $\mathbf{X}$  is locally finite,  $\mathbf{X}_0$  is finite. We assert that  $\mathbf{X}_0$  must be a single simplex.

Assume by way of contradiction that  $\mathbf{X}_0$  is not a simplex. Then, by Lemma 6.1,  $\mathbf{X}_0$  contains two vertices  $v, w$  such that  $B_1(v) \subsetneq B_1(w)$  (i.e.,  $v$  is a strictly dominated vertex). Since the strict inclusion of 1-balls is a transitive relation and  $\mathbf{X}_0$  is finite, there exists a finite set  $S$  of strictly dominated vertices of  $\mathbf{X}$  with the following property: for a vertex  $x \in S$  there is no vertex  $y$  with  $B_1(y) \subsetneq B_1(x)$ . Let  $\mathbf{X}'_0$  be the full subcomplex of  $\mathbf{X}$  spanned by  $V(\mathbf{X}_0) \setminus S$ . It is clear that  $\mathbf{X}'_0$  is a non-empty  $G$ -invariant proper subcomplex of  $\mathbf{X}_0$ . By Lemma 6.2,  $\mathbf{X}'_0$  is weakly systolic. This contradicts the minimality of  $\mathbf{X}_0$  and thus shows that  $\mathbf{X}_0$  has to be a simplex.  $\square$

**Corollary 6.4** (Conj. classes of finite subgroups). *Let  $G$  be a group acting geometrically by automorphisms on a weakly systolic complex  $\mathbf{X}$  (i.e.,  $G$  is weakly systolic). Then  $G$  contains only finitely many conjugacy classes of finite subgroups.*

*Proof.* Suppose by way of contradiction that we have infinitely many conjugacy classes of finite subgroups represented by  $H_1, H_2, \dots \subset G$ . Since  $G$  acts geometrically on  $\mathbf{X}$ , there exists a compact subset  $K \subset V(\mathbf{X})$  with  $\bigcup_{g \in G} gK = \mathbf{X}$ . For  $i = 1, 2, \dots$ , let  $\sigma_i$  be an  $H_i$ -invariant simplex of  $\mathbf{X}$  (whose existence is assured by the fixed point Theorem 6.3) and let  $g_i \in G$  be such that  $g_i(\sigma_i) \cap K \neq \emptyset$ . Then  $g_i(\sigma_i)$  is  $g_i H_i G_i^{-1}$  invariant and  $\bigcup_i g_i H_i G_i^{-1}$  is infinite. But for every element  $g \in \bigcup_i g_i H_i G_i^{-1}$  we have  $g(B_1(K)) \cap B_1(K) \neq \emptyset$ , a contradiction with the properness of  $G$ -action on  $\mathbf{X}$ .  $\square$

## 7. CONTRACTIBILITY OF THE FIXED POINT SET

The aim of this section is to prove that for a group acting on a weakly systolic complex its fixed point set is contractible (Proposition 7.6). As explained in the Introduction, this result implies Theorem E asserting that weakly systolic complexes are models for  $\underline{E}G$  for groups acting on them properly.

Our proof closely follows Przytycki's proof of an analogous result for the case of systolic complexes [Prz09]. There are however minor technical difficulties. In particular, since balls

around simplices in weakly systolic complexes need not to be convex, we have to work with other convex objects that are defined as follows. For a simplex  $\sigma$  of a simplicial complex  $\mathbf{X}$ , set  $K_0(\sigma) = \sigma$  and  $K_i(\sigma) = \bigcap_{v \in \sigma} B_i(v)$  for  $i = 1, 2, \dots$

**Lemma 7.1** (Properties of  $K_i(\sigma)$ ). *Let  $\sigma$  be a simplex of a weakly systolic complex  $\mathbf{X}$ . Then, for  $i = 0, 1, 2, \dots$ ,  $K_i(\sigma)$  is convex and  $K_{i+1}(\sigma) \subseteq B_1(K_i(\sigma))$ .*

*Proof.* Trivially,  $K_0(\sigma) = \sigma$  is convex. For  $i > 0$ ,  $K_i(\sigma)$  is the intersection of the balls  $B_i(v), v \in \sigma$ . By Theorem 3.1, balls around vertices are convex, whence  $K_i(\sigma)$  is convex as well. To establish the inclusion  $K_{i+1}(\sigma) \subseteq B_1(K_i(\sigma))$ , pick any vertex  $w \in K_{i+1}(\sigma)$ . Let  $l + 1 = d(w, \sigma)$  and denote by  $\sigma_0$  the metric projection of  $w$  in  $\sigma$ . By the property  $SD_l(w)$ , there exists a vertex  $z \in S_l(w)$  adjacent to all vertices of the simplex  $\sigma_0$ . Let  $w'$  be a neighbor of  $w$  in the interval  $I(w, z)$ . Then obviously  $d(w', \sigma) = l$  and therefore  $\sigma_0$  is the metric projection of  $w'$  in  $\sigma$ . Since  $d(w', v) = d(w, v) - 1$  for any vertex  $v \in \sigma$  and  $w \in K_{i+1}(\sigma)$ , we conclude that  $w' \in K_i(\sigma)$ , whence  $w \in B_1(w') \subset B_1(K_i(\sigma))$ .  $\square$

We recall now two general results that were proved in [Prz09] and which will be important in the proof of Proposition 7.6.

**Proposition 7.2** ([Prz09, Proposition 4.1]). *If  $\mathcal{C}, \mathcal{D}$  are posets and  $F_0, F_1: \mathcal{C} \rightarrow \mathcal{D}$  are functors such that for each object  $c$  of  $\mathcal{C}$  we have  $F_0(c) \leq F_1(c)$ , then the maps induced by  $F_0, F_1$  on the geometric realizations of  $\mathcal{C}, \mathcal{D}$  are homotopic. Moreover this homotopy can be chosen to be constant on the geometric realization of the subposet of  $\mathcal{C}$  of objects on which  $F_0$  and  $F_1$  agree.*

**Proposition 7.3** ([Prz09, Proposition 4.2]). *Let  $F_0: \mathcal{C}' \rightarrow \mathcal{C}$  be the functor from the flag poset  $\mathcal{C}'$  of a poset  $\mathcal{C}$  into the poset  $\mathcal{C}$ , assigning to each object of  $\mathcal{C}'$ , which is a chain of objects of  $\mathcal{C}$ , its minimal element. Then the map induced by  $F_0$  on geometric realizations of  $\mathcal{C}', \mathcal{C}$  (that are homeomorphic in a canonical way) is homotopic to identity.*

The following property of flag complexes will be crucial in the definition of expansion by projection below. It says that in weakly systolic case we can define projections on convex subcomplexes the same way as projections on balls.

**Lemma 7.4** (Projections on convexes). *Let  $\mathbf{X}$  be a simplicial flag complex and let  $Y$  be its convex subset. If a simplex  $\sigma$  belongs to  $S_1(Y)$ , i.e.  $\sigma \subseteq B_1(Y)$  and  $\sigma \cap Y = \emptyset$ , then  $\tau := \text{lk}(\sigma, \mathbf{X}) \cap Y$  is a single simplex.*

*Proof.* By definition of links,  $\tau$  consists of all vertices  $v$  of  $Y$  adjacent in  $G(\mathbf{X})$  to all vertices of  $\sigma$ . Since the set  $Y$  is convex and  $\sigma$  is disjoint from  $Y$ , necessarily the vertices of  $\tau$  are pairwise adjacent. As  $\mathbf{X}$  is a flag complex,  $\tau$  is a simplex of  $\mathbf{X}$ .  $\square$

We will call the simplex  $\tau$  as in the lemma above the *projection of  $\sigma$  on  $Y$* . Now we are in position to define the following notion introduced (in a more general version) by Przytycki [Prz09, Definition 3.1] in the systolic case. Let  $Y$  be a convex subset of a weakly systolic

complex  $\mathbf{X}$  and let  $\sigma$  be a simplex in  $B_1(Y)$ . The *expansion by projection*  $e_Y(\sigma)$  of  $\sigma$  is a simplex in  $B_1(Y)$  defined in the following way: if  $\sigma \subseteq Y$ , then  $e_Y(\sigma) = \sigma$ , otherwise  $e_Y(\sigma)$  is the join of  $\sigma \cap S_1(Y)$  and its projection on  $Y$ . A version of the following simple lemma was proved in [Prz09] in the systolic case. Its proof given there is valid also in our case.

**Lemma 7.5** ([Prz09, Lemma 3.8]). *Let  $Y$  be a convex subset of a weakly systolic complex  $\mathbf{X}$  and let  $\sigma_1 \subseteq \sigma_2 \subseteq \dots \subseteq \sigma_n \subseteq B_1(Y)$  be an increasing sequence of simplices. Then the intersection  $(\bigcap_{i=1}^n e_Y(\sigma_i)) \cap Y$  is nonempty.*

Let  $\sigma$  be a simplex of a weakly systolic complex  $\mathbf{X}$ . As in [Prz09], we define an increasing sequence of full subcomplexes  $\mathbf{D}_{2i}(\sigma)$  and  $\mathbf{D}_{2i+1}(\sigma)$  of the barycentric subdivision  $\mathbf{X}'$  of  $\mathbf{X}$  in the following way. Let  $\mathbf{D}_{2i}(\sigma)$  be the subcomplex spanned by all vertices of  $\mathbf{X}'$  corresponding to simplices of  $\mathbf{X}$  which have all their vertices in  $K_i(\sigma)$ . Let  $\mathbf{D}_{2i+1}(\sigma)$  be the subcomplex spanned by all vertices of  $\mathbf{X}'$  which correspond to those simplices of  $\mathbf{X}$  that have all their vertices in  $K_{i+1}(\sigma)$  and at least one vertex in  $K_i(\sigma)$ . The proof of the main proposition in this section follows closely the proof of [Prz09, Proposition 1.4].

**Proposition 7.6** (Contractibility of the fixed point set). *Let  $H$  be a group acting by simplicial automorphisms on a weakly systolic complex  $\mathbf{X}$ . Then the complex  $\text{Fix}_H \mathbf{X}'$  is contractible or empty.*

*Proof.* Assume that  $\text{Fix}_H \mathbf{X}'$  is nonempty and let  $\sigma$  be a maximal  $H$ -invariant simplex. By  $\mathbf{D}_i$  we will denote here  $\mathbf{D}_i(\sigma)$ . We will prove the following three assertions.

- (i)  $\mathbf{D}_0 \cap \text{Fix}_H \mathbf{X}'$  is contractible;
- (ii) the inclusion  $\mathbf{D}_{2i} \cap \text{Fix}_H \mathbf{X}' \subseteq \mathbf{D}_{2i+1} \cap \text{Fix}_H \mathbf{X}'$  is a homotopy equivalence;
- (iii) the identity on  $\mathbf{D}_{2i+2} \cap \text{Fix}_H \mathbf{X}'$  is homotopic to a mapping with image in  $\mathbf{D}_{2i+1} \cap \text{Fix}_H \mathbf{X}' \subseteq \mathbf{D}_{2i+2} \cap \text{Fix}_H \mathbf{X}'$ .

As in the proof of [Prz09, Proposition 1.4], the three assertions imply that  $\mathbf{D}_k \cap \text{Fix}_H \mathbf{X}'$  is contractible for every  $k$ , thus the proposition holds. To show (i), note that  $\mathbf{D}_0 \cap \text{Fix}_H \mathbf{X}'$  is a cone over the barycenter of  $\sigma$  and hence it is contractible.

To prove (ii), let  $\mathcal{C}$  be the poset of  $H$ -invariant simplices in  $\mathbf{X}$  with vertices in  $K_{i+1}(\sigma)$  and at least one vertex in  $K_i(\sigma)$ . Its geometric realization is  $\mathbf{D}_{2i+1} \cap \text{Fix}_H \mathbf{X}'$ . Consider a functor  $F: \mathcal{C} \rightarrow \mathcal{C}$  assigning to each object of  $\mathcal{C}$  (i.e., each simplex of  $\mathbf{X}$ ), its subsimplex spanned by its vertices in  $K_i(A)$ . By Proposition 7.2, the geometric realization of  $F$  is homotopic to identity (which is the geometric realization of the identity functor). Moreover this homotopy is constant on  $\mathbf{D}_{2i} \cap \text{Fix}_H \mathbf{X}'$ . The image of the geometric realization of  $F$  is contained in  $\mathbf{D}_{2i} \cap \text{Fix}_H \mathbf{X}'$ . Hence  $\mathbf{D}_{2i} \cap \text{Fix}_H \mathbf{X}'$  is a deformation retract of  $\mathbf{D}_{2i+1} \cap \text{Fix}_H \mathbf{X}'$ , as desired.

To establish (iii), let  $\mathcal{C}$  be the poset of  $H$ -invariant simplices of  $\mathbf{X}'$  with vertices in  $K_{i+1}(\sigma)$  and let  $\mathcal{C}'$  be its flag poset. Let also  $F_0: \mathcal{C}' \rightarrow \mathcal{C}$  be the functor assigning to each object of  $\mathcal{C}'$  its minimal element; cf. Proposition 7.3. Now we define another functor  $F_1: \mathcal{C}' \rightarrow \mathcal{C}$ . For any object  $c'$  of  $\mathcal{C}'$ , which is a chain of objects  $c_1 < c_2 < \dots < c_k$  of  $\mathcal{C}$ , recall that  $c_j$

are some  $H$ -invariant simplices in  $K_{i+1}(\sigma)$ . Let  $c'_j = e_{K_i(\sigma)}(c_j)$ . Then by Lemma 7.5 the intersection  $\bigcap_{j=1}^k c'_j$  contains at least one vertex in  $K_i(\sigma)$ . Thus  $\bigcap_{j=1}^k c'_j$  is an  $H$ -invariant non-empty simplex and hence it is an object of  $\mathcal{C}$ . We define  $F_1(c')$  to be this object. In the geometric realization of  $\mathcal{C}$ , which is  $\mathbf{D}_{2i+2} \cap \text{Fix}_H \mathbf{X}'$ , the object  $F_1(c')$  corresponds to a vertex of  $\mathbf{D}_{2i+1} \cap \text{Fix}_H \mathbf{X}'$ . It is obvious that  $F_1$  preserves the partial order. Notice that for any object  $c'$  of  $\mathcal{C}'$  we have  $F_0(c') \subseteq F_1(c')$ , hence, by Proposition 7.3, the geometric realizations of  $F_0$  and  $F_1$  are homotopic. We have that  $F_0$  is homotopic to the identity and that  $F_1$  has image in  $\mathbf{D}_{2i+1} \cap \text{Fix}_H \mathbf{X}'$ , thus establishing (iii).  $\square$

## 8. FINAL REMARKS ON THE CASE OF SYSTOLIC COMPLEXES

In this final section, we restrict to the case of systolic complexes and we present some further results in that case. First, using Lemma 3.10 and Theorem 3.11 of Polat [Pol02] for bridged graphs, we prove a stronger version of the fixed point theorem for systolic complexes. Namely, Polat [Pol02] established that for any subset  $\bar{Y}$  of vertices of a graph with finite intervals, there exists a minimal isometric subgraph of this graph which contains  $\bar{Y}$ . Moreover, if  $\bar{Y}$  is finite and the graph is bridged, then [Pol02, Theorem 3.11(i)] shows that this minimal isometric (and hence bridged) subgraph is also finite. We continue with two lemmata which can be viewed as  $G$ -invariant versions of these two results of Polat [Pol02].

**Lemma 8.1** (Minimal subcomplex). *Let a group  $G$  act by simplicial automorphisms on a systolic complex  $\mathbf{X}$ . Let  $\bar{Y}$  be a  $G$ -invariant set of vertices of  $\mathbf{X}$ . Then there exists a minimal  $G$ -invariant subcomplex  $\mathbf{Y}$  of  $\mathbf{X}$  containing  $\bar{Y}$ , which is itself a systolic complex.*

*Proof.* Let  $\Sigma$  be a chain (with respect to the subcomplex relation) of  $G$ -invariant subcomplexes of  $\mathbf{X}$ , which contain  $\bar{Y}$  and induce isometric subgraphs of the underlying graph of  $\mathbf{X}$  (and thus are systolic complexes themselves). Then, as in the proof of [Pol02, Lemma 3.10], we conclude that the subcomplex  $\mathbf{Y} = \bigcap \Sigma$  is a minimal  $G$ -invariant subcomplex of  $\mathbf{X}$ , containing  $\bar{Y}$  and which is itself a systolic complex.  $\square$

**Lemma 8.2** (Minimal finite subcomplex). *Let a group  $G$  act by simplicial automorphisms on a systolic complex  $X$ . Let  $\bar{Y}$  be a finite  $G$ -invariant set of vertices of  $X$ . Then there exists a minimal (as a simplicial complex) finite  $G$ -invariant subcomplex  $Y$  of  $X$ , which is itself a systolic complex.*

*Proof.* Let  $co(\bar{Y})$  be the convex hull of  $\bar{Y}$  in  $\mathbf{X}$ . The full subcomplex  $\mathbf{Z}$  of  $\mathbf{X}$  spanned by  $co(\bar{Y})$  is a bounded systolic complex. By Lemma 8.1, there exists a minimal  $G$ -invariant subcomplex  $\mathbf{Y}$  of  $\mathbf{Z}$  containing the set  $\bar{Y}$  and which itself is a systolic complex. Then, applying the proof of [Pol02, Theorem 3.11] to the bounded bridged graphs which are the underlying graphs of the systolic complexes  $\mathbf{Y}$  of  $\mathbf{Z}$ , it follows that  $\mathbf{Y}$  is finite.  $\square$

**Theorem 8.3** (The fixed point theorem). *Let  $G$  be a finite group acting by simplicial automorphisms on a systolic complex  $\mathbf{X}$ . Then there exists a simplex  $\sigma \in \mathbf{X}$  which is invariant under the action of  $G$ .*

*Proof.* Let  $\bar{Y} = Gv = \{gv \mid g \in G\}$ , for some vertex  $v \in \mathbf{X}$ . Then  $\bar{Y}$  is a finite  $G$ -invariant set of vertices of  $\mathbf{X}$  and thus, by Lemma 8.2, there exists a minimal finite  $G$ -invariant subcomplex  $\mathbf{Y}$  of  $\mathbf{X}$ , which is itself a systolic complex. Then, the same way as in the proof of Theorem 6.3, we conclude that there exists a simplex in  $\mathbf{Y}$  that is  $G$ -invariant.  $\square$

**Remark 8.4.** We believe that, as in the systolic case, the stronger version of Theorem 6.3 holds also for weakly systolic complexes, i.e., one can drop the assumption on the local finiteness of  $\mathbf{X}$  in Theorem 6.3. This needs extensions of some results of Polat (in particular, Theorems 3.8 and 3.11 from [Pol02]) to the class of weakly bridged graphs.

Zawiślak [Zaw04] initiated another approach to the fixed point theorem in the systolic case based on the following notion of round subcomplexes. A systolic complex  $\mathbf{X}$  of finite diameter  $k$  is *round* (cf. [Prz08]) if  $\cap\{B_{k-1}(v) : v \in V(\mathbf{X})\} = \emptyset$ . Przytycki [Prz08] established that all round systolic complexes have diameter at most 5 and used this result to prove that for any finite group  $G$  acting by simplicial automorphisms on a systolic complex there exists a subcomplex of diameter at most 5 which is invariant under the action of  $G$ . Zawiślak [Zaw04, Conjecture 3.3.1] and Przytycki (Remark 8.1 of [Prz08]) conjectured that in fact the diameter of round systolic complexes must be at most 2. Zawiślak [Zaw04, Theorem 3.3.1] showed that if this is true, then it implies that  $G$  has an invariant simplex, thus paving another way to the proof of Theorem 8.3. We will show now that the positive answer to the question of Zawiślak and Przytycki directly follows from an earlier result of Farber [Far89] on diameters and radii of finite bridged graphs.

**Proposition 8.5** (Round systolic complexes). *Any round systolic complex  $\mathbf{X}$  has diameter at most 2.*

*Proof.* Let  $\text{diam}(\mathbf{X})$  and  $\text{rad}(\mathbf{X})$  denote the diameter and the radius of a systolic complex  $\mathbf{X}$ , i.e., the diameter and radius of its underlying bridged graph  $G = G(\mathbf{X})$ . Recall that  $\text{rad}(\mathbf{X})$  is the smallest integer  $r$  such that there exists a vertex  $c$  of  $\mathbf{X}$  (called a central vertex) so that the ball  $B_r(c)$  of radius  $r$  and centered at  $c$  covers all vertices of  $\mathbf{X}$ , i.e.,  $B_r(c) = V(\mathbf{X})$ .

Farber [Far89, Theorem 4] proved that if  $G$  is a finite bridged graph, then  $3\text{rad}(G) \leq 2\text{diam}(G) + 2$ . We will show first that this inequality holds for infinite bridged graphs  $G$  of finite diameter  $\text{diam}(G)$  and containing no infinite simplices. Set  $k := \text{rad}(G) \leq \text{diam}(G)$ . By definition of  $\text{rad}(G)$  the intersection of all balls of radius  $k - 1$  of  $G$  is empty. Then using an argument of Polat (personal communication) presented below, we can find a finite subset of vertices  $Y$  of  $G$  such that the intersection of the balls  $B_{k-1}(v)$ ,  $v$  running over all vertices of  $Y$ , is still empty. By [Pol02, Theorem 3.11], there exists a finite isometric bridged subgraph  $H$  of  $G$  containing  $Y$ . From the choice of  $Y$  we conclude that the radius of  $H$  is at least  $k$ , while the diameter of  $H$  is at most the diameter of  $G$ . As a result, applying Farber's inequality to  $H$ , we obtain  $3\text{rad}(G) \leq 3\text{rad}(H) \leq 2\text{diam}(H) + 2 \leq 2\text{diam}(G) + 2$ , whence  $3\text{rad}(G) \leq 2\text{diam}(G) + 2$ .

To show the existence of a finite set  $Y$  such that  $\cap\{B_{k-1}(v) : v \in Y\} = \emptyset$ , we use an argument of Polat (personal communication). According to Theorem 3.9 of [Pol98], any

graph without isometric rays (in particular, any bridged graph of finite diameter) can be endowed with a topology, called *geodesic topology*, so that the resulting topological space is compact. On the other hand, it is shown in [Pol04, Corollary 6.26] that any convex set of a bridged graph containing no infinite simplices is closed in the geodesic topology. As a result, the balls of a bridged graph  $G$  of finite diameter containing no infinite simplices are compact convex sets. Hence any family of balls with an empty intersection contains a finite subfamily with an empty intersection, showing that such a finite set  $Y$  indeed exists.

Now suppose that  $\mathbf{X}$  is a round systolic complex and let  $k := \text{diam}(\mathbf{X})$ . Since  $\mathbf{X}$  is round, one can easily deduce that  $\text{rad}(\mathbf{X}) = k$ : indeed, if  $\text{rad}(\mathbf{X}) \leq k - 1$  and  $c$  is a central vertex, then  $c$  will belong to the intersection  $\cap \{B_{k-1}(v) : v \in V(\mathbf{X})\}$ , which is impossible. Applying Farber's inequality to the (bridged) underlying graph of  $\mathbf{X}$ , we conclude that  $3k \leq 2k + 2$ , whence  $k \leq 2$ .  $\square$

**Remark 8.6.** It would be interesting to extend Proposition 8.5 and the relationship of [Far89] between radii and diameters to weakly systolic complexes.

Osajda-Przytycki [OP09] constructed a  $Z$ -set compactification  $\overline{\mathbf{X}} = \mathbf{X} \cup \partial\mathbf{X}$  of a systolic complex  $\mathbf{X}$ . The main result there ([OP09, Theorem 6.3]) together with Theorem E from the Introduction of our paper, allowed them to claim the following result (without proving it):

**Claim 8.7** ([OP09, Theorem 6.3 and Claim 14.2]). Let a group  $G$  act geometrically by simplicial automorphisms on a systolic complex  $\mathbf{X}$ . Then the compactification  $\overline{\mathbf{X}} = \mathbf{X} \cup \partial\mathbf{X}$  of  $\mathbf{X}$  satisfies the following properties:

1.  $\overline{\mathbf{X}}$  is a Euclidean retract (ER);
2.  $\partial\mathbf{X}$  is a  $Z$ -set in  $\overline{\mathbf{X}}$ ;
3. for every compact set  $K \subset \mathbf{X}$ ,  $(gK)_{g \in G}$  is a null sequence;
4. the action of  $G$  on  $\mathbf{X}$  extends to an action, by homeomorphisms, of  $G$  on  $\overline{\mathbf{X}}$ ;
5. for every finite subgroup  $F$  of  $G$ , the fixed point set  $\text{Fix}_F \overline{\mathbf{X}}$  is contractible;
6. for every finite subgroup  $F$  of  $G$ , the fixed point set  $\text{Fix}_F \mathbf{X}$  is dense in  $\text{Fix}_F \overline{\mathbf{X}}$ .

This result asserts that  $\overline{\mathbf{X}}$  is an  $EZ$ -structure, sensu Rosenthal [Ros03], for a systolic group  $G$ ; for details, see [OP09]. The existence of such a structure implies, by [Ros03], the Novikov conjecture for  $G$ .

#### ACKNOWLEDGEMENTS

Work of V. Chepoi was supported in part by the ANR grant BLAN06-1-138894 (projet OPTICOMB). Work of D. Osajda was supported in part by MNiSW grant N201 012 32/0718 and by the ANR grants Cannon and Théorie Géométrique des Groupes. We are grateful to Norbert Polat for his help in the proof of Proposition 8.5.

## REFERENCES

- [AF88] Richard P. Anstee and Martin Farber, *On bridged graphs and cop-win graphs*, J. Combin. Theory Ser. B **44** (1988), no. 1, 22–28. MR923263 (89h:05053)
- [BC96] Hans-Jürgen Bandelt and Victor Chepoi, *A Helly theorem in weakly modular space*, Discrete Math. **160** (1996), no. 1-3, 25–39. MR1417558 (97h:52006)
- [BC08] ———, *Metric graph theory and geometry: a survey*, Surveys on discrete and computational geometry, Contemp. Math., vol. 453, Amer. Math. Soc., Providence, RI, 2008, pp. 49–86. MR2405677 (2009h:05068)
- [BH99] Martin R. Bridson and André Haefliger, *Metric spaces of non-positive curvature*, Grundlehren der Mathematischen Wissenschaften [Fundamental Principles of Mathematical Sciences], vol. 319, Springer-Verlag, Berlin, 1999. MR1744486 (2000k:53038)
- [08] *Guido's book of conjectures*, Monographies de L'Enseignement Mathématique [Monographs of L'Enseignement Mathématique], vol. 40, L'Enseignement Mathématique, Geneva, 2008. A gift to Guido Mislin on the occasion of his retirement from ETHZ June 2006; Collected by Indira Chatterji. MR2499538
- [Che89] Victor Chepoi, *Classification of graphs by means of metric triangles*, Metody Diskret. Analiz. **49** (1989), 75–93, 96 (Russian). MR1114014 (92e:05041)
- [Che97] ———, *Bridged graphs are cop-win graphs: an algorithmic proof*, J. Combin. Theory Ser. B **69** (1997), no. 1, 97–100. MR1426753 (97g:05150)
- [Che98] ———, *On distance-preserving and domination elimination orderings*, SIAM J. Discrete Math. **11** (1998), no. 3, 414–436 (electronic). MR1628110 (99h:05032)
- [Che00] ———, *Graphs of some CAT(0) complexes*, Adv. in Appl. Math. **24** (2000), no. 2, 125–179. MR1748966 (2001a:57004)
- [CY07] Yusuf Civan and Ergün Yalçın, *Linear colorings of simplicial complexes and collapsing*, J. Combin. Theory Ser. A **114** (2007), no. 7, 1315–1331. MR2353125 (2009a:05067)
- [ECH<sup>+</sup>92] David B. A. Epstein, James W. Cannon, Derek F. Holt, Silvio V. F. Levy, Michael S. Paterson, and William P. Thurston, *Word processing in groups*, Jones and Bartlett Publishers, Boston, MA, 1992. MR1161694 (93i:20036)
- [Far89] Martin Farber, *On diameters and radii of bridged graphs*, Discrete Math. **73** (1989), no. 3, 249–260. MR983023 (90d:05192)
- [FJ87] Martin Farber and Robert E. Jamison, *On local convexity in graphs*, Discrete Math. **66** (1987), no. 3, 231–247. MR900046 (89e:05167)
- [Gro87] Mikhail Gromov, *Hyperbolic groups*, Essays in group theory, Math. Sci. Res. Inst. Publ., vol. 8, Springer, New York, 1987, pp. 75–263. MR919829 (89e:20070)
- [Hag03] Frédéric Haglund, *Complexes simpliciaux hyperboliques de grande dimension*, Prepublication Orsay **71** (2003), preprint, available at [http://www.math.u-psud.fr/~biblio/ppo/2003/fic/ppo\\_2003\\_71.pdf](http://www.math.u-psud.fr/~biblio/ppo/2003/fic/ppo_2003_71.pdf).
- [HN04] Pavol Hell and Jaroslav Nešetřil, *Graphs and homomorphisms*, Oxford Lecture Series in Mathematics and its Applications, vol. 28, Oxford University Press, Oxford, 2004. MR2089014 (2005k:05002)
- [JŚ06] Tadeusz Januszkiewicz and Jacek Świątkowski, *Simplicial nonpositive curvature*, Publ. Math. Inst. Hautes Études Sci. **104** (2006), 1–85, DOI 10.1007/s10240-006-0038-5. MR2264834 (2007j:53044)
- [Lüc05] Wolfgang Lück, *Survey on classifying spaces for families of subgroups*, Infinite groups: geometric, combinatorial and dynamical aspects, Progr. Math., vol. 248, Birkhäuser, Basel, 2005, pp. 269–322.
- [LS77] Roger C. Lyndon and Paul E. Schupp, *Combinatorial group theory*, Springer-Verlag, Berlin, 1977. Ergebnisse der Mathematik und ihrer Grenzgebiete, Band 89. MR0577064 (58 #28182)
- [Mat08] Jiří Matoušek, *LC reductions yield isomorphic simplicial complexes*, Contrib. Discrete Math. **3** (2008), no. 2, 37–39. MR2455228 (2009g:55031)

- [NW83] Richard Nowakowski and Peter Winkler, *Vertex-to-vertex pursuit in a graph*, Discrete Math. **43** (1983), no. 2-3, 235–239. MR685631 (84d:05138)
- [Osa07] Damian Osajda, *Connectedness at infinity of systolic complexes and groups*, Groups Geom. Dyn. **1** (2007), no. 2, 183–203. MR2319456 (2008e:20064)
- [Osa09a] ———, *A combinatorial non-positive curvature I:  $SD_n$  property* (2009), in preparation.
- [Osa09b] ———, *A construction of hyperbolic Coxeter groups* (2009), in preparation.
- [OP09] Damian Osajda and Piotr Przytycki, *Boundaries of systolic groups*, Geom. Topol. **13** (2009), no. 5, 2807–2880.
- [OŚ09] Damian Osajda and Jacek Świątkowski, *On asymptotically hereditarily aspherical groups* (2009), in preparation.
- [Pol98] Norbert Polat, *Graphs without isometric rays and invariant subgraph properties. I*, J. Graph Theory **27** (1998), no. 2, 99–109. MR1491562 (99a:05086)
- [Pol00] ———, *On infinite bridged graphs and strongly dismantlable graphs*, Discrete Math. **211** (2000), no. 1-3, 153–166. MR1735348 (2000k:05232)
- [Pol02] ———, *On isometric subgraphs of infinite bridged graphs and geodesic convexity*, Discrete Math. **244** (2002), no. 1-3, 399–416. Algebraic and topological methods in graph theory (Lake Bled, 1999). MR1844048 (2003c:05070)
- [Pol04] ———, *Fixed finite subgraph theorems in infinite weakly modular graphs*, Discrete Math. **285** (2004), no. 1-3, 239–256. MR2062847 (2005c:05174)
- [Prz08] Piotr Przytycki, *The fixed point theorem for simplicial nonpositive curvature*, Math. Proc. Cambridge Philos. Soc. **144** (2008), no. 3, 683–695.
- [Prz09] ———,  *$EG$  for systolic groups*, Comment. Math. Helv. **84** (2009), no. 1, 159–169.
- [Qui83] Alain Quilliot, *Problèmes de jeux, de point fixe, de connectivité et de représentation sur des graphes, des ensembles ordonnés et des hypergraphes*, Thèse de doctorat d'état, Université de Paris VI, 1983 (French).
- [Rol98] Martin A. Roller, *Poc sets, median algebras and group actions. An extended study of Dunwoody's construction and Sageev's theorem*, Univ. of Southampton Preprint Ser. (1998), preprint.
- [RTL76] Donald J. Rose, R. Endre Tarjan, and George S. Lueker, *Algorithmic aspects of vertex elimination on graphs*, SIAM J. Comput. **5** (1976), no. 2, 266–283. MR0408312 (53 #12077)
- [Ros03] David Rosenthal, *Split injectivity of the Baum–Connes assembly map* (2003), preprint, available at [arXiv:math/0312047](https://arxiv.org/abs/math/0312047).
- [SC83] Valeriu P. Soltan and Victor Chepoi, *Conditions for invariance of set diameters under  $d$ -convexification in a graph*, Kibernetika (Kiev) **6** (1983), 14–18 (Russian, with English summary); English transl., Cybernetics **19** (1983), no. 6, 750–756 (1984). MR765117 (86k:05102)
- [Wee94] Graham M. Weetman, *A construction of locally homogeneous graphs*, J. London Math. Soc. (2) **50** (1994), no. 1, 68–86. MR1277755 (95d:05113)
- [Wis03] Daniel T. Wise, *Sixtolic complexes and their fundamental groups* (2003), unpublished manuscript.
- [Zaw04] Paweł Zawiślak, *O pewnych własnościach 6-systolicznych kompleksów symplecjalnych*, M.Sc. thesis, Wrocław University, 2004 (Polish).