

The interval number of a planar graph is at most three

Guillaume Guégan¹, Kolja Knauer^{2,3}, Jonathan Rollin⁴, and
Torsten Ueckerdt⁵

¹Université de Montpellier, CNRS, LIRMM, France

²Aix Marseille Univ, Univ de Toulon, CNRS, LIS, Marseille, France

³Departament de Matemàtiques i Informàtica, Universitat de
Barcelona (UB), Barcelona, Spain

⁴FernUniversität in Hagen, Hagen, Germany

⁵Karlsruhe Institute of Technology, Karlsruhe, Germany

July 25, 2019

Abstract

The interval number of a graph G is the minimum k such that one can assign to each vertex of G a union of k intervals on the real line, such that G is the intersection graph of these sets, i.e., two vertices are adjacent in G if and only if the corresponding sets of intervals have non-empty intersection.

In 1983 Scheinerman and West [The interval number of a planar graph: Three intervals suffice. *J. Comb. Theory, Ser. B*, 35:224–239, 1983] proved that the interval number of any planar graph is at most 3. However the original proof has a flaw. We give a different and shorter proof of this result.

1 Introduction

For a positive integer k , a k -interval representation of a graph $G = (V, E)$ is a set $\{f(v) \mid v \in V\}$ where $f(v)$ is the union of at most k intervals on the real line representing vertex $v \in V$, such that uv ($u \neq v$) is an edge of G if and only if $f(u) \cap f(v) \neq \emptyset$. In 1979 Trotter and Harary [16] introduced the *interval number* of G , denoted by $i(G)$, as the smallest k such that G has a k -interval representation. In 1983 Scheinerman and West [13] constructed planar graphs with interval number at least 3 and proposed a proof for the following:

Theorem 1 (Scheinerman, West [13]). *If G is planar, then $i(G) \leq 3$.*

In this paper we point out a flaw in that proof and give an alternative proof of Theorem 1. The original proof is based on induction, maintaining a very comprehensive description of the (partial) representation and its properties. However in certain cases, one of the required properties can not be maintained

during the construction. A precise description of the problematic case is given in Section 4. Let us also remark that in his thesis [12], Scheinerman proposes a slightly different argumentation for the considered case which also leads to the same problem.

In the remaining part of the introduction we discuss some related work. In Section 2 we introduce the notions used in the new proof, which is then given in Section 3. We conclude with final remarks in Section 5.

Related work. A graph with $i(G) \leq 1$ is called an *interval graph*. A concept closely related to the interval number is the *track number* of G , denoted by $t(G)$, which is the smallest k such that G is the union of k interval graphs. Equivalently, $t(G)$ is the smallest k such that G admits a k -interval representation that is the union of k 1-interval representations, each on a different copy of the real line (called a track) and each containing one interval per vertex. More recently, Knauer and Ueckerdt [10] defined the *local track number* of G , denoted by $t_\ell(G)$, to be the smallest k such that G is the union of d interval graphs, for some d , where every vertex of G is contained in at most k of them. It is easy to see that for every graph G we have $i(G) \leq t_\ell(G) \leq t(G)$.

Gonçalves [7] proved that the track number of planar graphs is at most 4, which is best-possible [8]. It is an open problem whether there is a planar graph with local track number 4 [10]. Balogh *et al.* [2] show that any planar graph of maximum degree at most 4 has interval number at most 2. For further recent results about interval numbers of other classes of planar graphs and general graphs, let us refer to [3] and [6], respectively.

Finally, let us mention that West and Shmoys [17] proved that deciding $i(G) \leq k$ is NP-complete for each $k \geq 2$, while Jiang [9] proved the same for $t(G) \leq k$, and Stumpf [14] for $t_\ell(G) \leq k$.

2 Preliminaries

All graphs considered here are finite, simple, undirected, and non-empty. If $G = (V, E)$ is a graph and $\{f(v) \mid v \in V\}$ is a k -interval representation of G for some k , we say that a subset S of the real line is *intersected* by some $f(v)$ if $S \cap f(v) \neq \emptyset$. For a point p with $p \in f(v)$, we also say that p is *covered* by $f(v)$. A vertex v has a *broken end* b if b is an endpoint of an interval in $f(v)$ and b is not covered by any other $f(w)$ for $w \in V - \{v\}$. We say that a vertex v is *displayed* if $f(v)$ contains a *portion* (i.e., a non-empty open interval) not intersected by any other $f(w)$ for $w \in V - \{v\}$. An edge $uv \in E$ is *displayed* if $f(u) \cap f(v)$ contains a portion not intersected by any other $f(w)$ for $w \in V - \{u, v\}$.

The *depth* of the representation $\{f(v) \mid v \in V\}$ is the largest integer d such that some point of the real line is covered by at least d sets $f(v)$ with $v \in V$. For $d \geq 1$, the *depth d interval number* of G , denoted by $i_d(G)$, is the smallest k such that G admits a k -interval representation of depth d . Scheinerman and West [13] proved that there are planar graphs G with $i_2(G) \geq 4$, while Gonçalves [7] proved that $i_2(G) \leq 4$ for all planar graphs G . In Theorem 3 we obtain that for any 4-connected planar graph G it holds $i_2(G) \leq 3$. Finally, we shall prove here that $i_3(G) \leq 3$ for all planar graphs G , which is also the original claim of Scheinerman and West.

3 An alternative proof of Theorem 1

In this section we give a new proof for Theorem 1, i.e., that every planar graph G has interval number at most 3. A *triangulation* is a plane embedded graph in which every face is bounded by a triangle. In other words, a triangulation is a maximally planar graph on at least three vertices with a fixed plane embedding. As every planar graph is an induced subgraph of some triangulation (For example, iteratively adding a new vertex to a non-triangular face together with one edge to each of the face's incident vertices, eventually results in such a triangulation.) and the interval number is monotone under taking induced subgraphs, we may assume without loss of generality that G is a triangulation.

A triangle in G is *non-empty* if its interior contains at least one vertex of G . We shall construct a 3-interval representation of G by recursively splitting G along its non-empty triangles. This leaves us with the task to represent 4-connected triangulations, i.e., triangulations whose only non-empty triangle is the outer triangle, and to “glue” those representation along the non-empty triangles of G . More precisely, we shall roughly proceed as follows:

- (I) Consider a non-empty triangle Δ with inclusion-minimal interior and the set X of all vertices in its interior.
- (II) Call induction on the graph $G_1 = G - X$, obtaining a 3-interval representation f_1 of G_1 with additional properties on how inner faces are represented.
- (III) Since the subgraph G_2 of G induced by $V(\Delta) \cup X$ is 4-connected, we can utilize a recent result of the second and fourth author to decompose G_2 into a path and two forests.
- (IV) Using this decomposition, we define a 3-interval representation f_2 of G_2 that coincides with f_1 on $V(\Delta)$.

We remark that the essentials of the construction in step (IV) can be already found in [1]. Also, the decomposition of a triangulation along its non-empty triangles is a common method in the field of intersection graphs; see e.g., [4,5,15]. Hence the key to our new proof is the most recent [11] decomposition used in step (III), which we shall state next.

Let $G = (V, E)$ be a 4-connected triangulation with outer vertices x, y, z . We denote by u_x the unique inner vertex of G adjacent to y and z (if there were several, G would not be 4-connected), and call the vertex u_x the vertex *opposing* x . Similarly, the vertex u_y opposing y and the vertex u_z opposing z are defined. Note that since G is 4-connected we have that u_x, u_y, u_z all coincide if $|V| = 4$, and are pairwise distinct if $|V| \geq 5$. We use the following recent result of the second and fourth author (Lemma 3.1 in [11]), which we have adapted here to the case of 4-connected triangulations (see Figure 1 for an illustration):

Lemma 2 (Knauer, Ueckerdt [11]). *Let G be a plane 4-connected triangulation with outer triangle $\Delta_{\text{out}} = x, y, z$ and corresponding opposing vertices u_x, u_y, u_z . Then the inner edges of G can be partitioned into three forests F_x, F_y, F_z such that*

- F_x is a Hamiltonian path of $G \setminus \{y, z\}$ going from x to u_x ,




Figure 1: The decomposition in Lemma 2 for the case $G \not\cong K_4$ (left) and $G \cong K_4$ (right).

- F_y is a spanning tree of $G \setminus \{x, z\}$,
- F_z is a spanning forest of $G \setminus \{y\}$ consisting of two trees, one containing x and one containing z , unless $G \cong K_4$. In this case $F_z = zu_z$.

Note that the conditions on the decomposition in Lemma 2 imply that the edge xu_y is in F_x and the edge zu_y is in F_z (since x and z are in different components of F_z and z is not in F_x). We are now ready to give our new proof of Theorem 1.

Proof of Theorem 1. We have to show that every planar graph G admits a 3-interval representation of depth at most 3, in particular that $i(G) \leq 3$. As every planar graph is an induced subgraph of some planar triangulation and the interval number is monotone under taking induced subgraphs, we may assume without loss of generality that G is a triangulation.

We proceed by induction on the number n of vertices in G , showing that G admits a 3-interval representation of depth 3 with the additional invariant that **(I1)** every vertex is displayed, and **(I2)** every inner face contains at least one displayed edge.

The base case is $n = 3$, i.e., G is a triangle with vertices x, y, z . In this case, it is easy to define a 3-interval representation of G with invariants **(I1)** and **(I2)**. For example, take $f(x) := [0, 3]$, $f(y) := [1, 4] \cup [6, 7]$, and $f(z) := [2, 5]$, and note that edges xy and yz are displayed.

Now assume that $n \geq 4$, i.e., G contains at least one non-empty triangle. Let Δ be a non-empty triangle in G with inclusion-minimal interior among all non-empty triangles in G . Let $X \subset V$ be the (non-empty) set of vertices in the interior of Δ , $G_{\text{out}} = G - X$ be the triangulation induced by $V - X$, and G_{in} be the 4-connected triangulation induced by $V(\Delta) \cup X$. By induction hypothesis there exists a 3-interval representation $\{f(v) \mid v \in V - X\}$ of G_{out} of depth 3 satisfying the invariant **(I1)** and **(I2)**. In particular, the three vertices x, y, z of Δ , which now form an inner face of G_{out} , are already assigned to up to three intervals each. We shall now extend this representation to the vertices in X .

By invariant **(I2)** at least one edge of Δ , say xz , is a displayed edge. Consider the vertices u_x, u_y, u_z opposing x, y, z in G_{in} , respectively, and let F_x, F_y, F_z be the decomposition of the inner edges of G_{in} given by Lemma 2 (note that we can choose x, y, z arbitrarily in Lemma 2). For convenience, if $G_{\text{in}} \cong K_4$, let F_z additionally have vertex x as a one-vertex component. We define, based on this decomposition, three intervals for each vertex $v \in X$ as follows.

- Represent the path $F_x - \{x\}$ on an unused portion of the real line by using one interval per vertex, and in such a way that every vertex and every edge is displayed. Let this representation be denoted by $\{f_1(v) \mid v \in X\}$.
- Consider y to be the root of the tree F_y , and for each vertex v in $F_y - \{y\}$ consider the parent w of v in F_y , i.e., the neighbor of v on the v -to- y path in F_y . Create a new interval for v strictly inside the displayed portion of w . If $w = y$, such a portion exists in $f(y)$ as **(I1)** holds for f , and if $w \neq y$, such a portion exists in $f_1(w)$. Let these new intervals be denoted by $\{f_2(v) \mid v \in X\}$.
- Consider the three trees that are the components of $F_z - \{u_y z\}$, that is, of F_z after removing the edge $u_y z$. Say T_1 contains vertex u_y (and possibly no other vertex), T_2 contains z , and T_3 contains x . Consider u_y , respectively z and x , to be the root of T_1 , respectively T_2 and T_3 . For each vertex v in $T_1 - \{u_y\}$, respectively $T_2 - \{z\}$ and $T_3 - \{x\}$, consider the parent w of v in T_1 , respectively T_2 and T_3 , and create a new interval for v in the displayed portion of w . (Again, if $w \in \{x, z\}$, such a portion exists in $f(w)$ as **(I1)** holds for f , and if $w \notin \{x, z\}$, such a portion exists in $f_1(w)$.) Let these new intervals be denoted by $\{f_3(v) \mid v \in X - \{u_y\}\}$.
- Finally, create a new interval $f_3(u_y)$ in the displayed portion of edge xz (to represent xu_y and zu_y).

Defining $f(v) = f_1(v) \cup f_2(v) \cup f_3(v)$ for each $v \in X$, it is straightforward to check that $\{f(v) \mid v \in V\}$ is a 3-interval representation of $G = G_{\text{out}} \cup G_{\text{in}}$ of depth 3. Moreover, the edge xz , every inner edge of G_{in} except for $u_y x$ and $u_y z$, and every vertex of X is displayed. Hence this representation satisfies our invariants **(I1)** and **(I2)**, which concludes the proof. \square

4 The approach of Scheinerman and West

To explain the proof strategy of Scheinerman and West, we introduce some more terminology from their paper. For a plane embedded graph G let G^0 denote the (outerplanar) subgraph induced by its *external vertices*, i.e., those lying on the unbounded, outer face. The edges of G^0 that bound the outer face are the *external edges*, while those that bound two inner faces are the *chords*. The graph G^0 is considered together with its decomposition into its inclusion-maximal 2-connected subgraphs, called blocks. Furthermore, we fix for each component of G^0 a non-cut-vertex z as the root of that component and say that $z = z_H$ is the root of its corresponding block H . For each remaining block H of G^0 let z_H be the cut-vertex of G^0 in H that is closest to the root of the corresponding component. For a chord xy of G^0 contained in block H with root z_H and a third vertex $u \neq x, y$ of H , say that u is *on the z_H^* -side* of xy if u and z_H lie in different connected components of $H - \{x, y\}$.

The argument of Scheinerman and West proceeds by induction on the number of vertices in G with a stronger induction hypothesis on how G^0 , i.e., the subgraph of G induced by the external vertices, is represented. Every edge xy of G^0 shall be represented in such a way that for a possible future vertex v which




FIG. 3. Adding an outervertex joined to the endpoints of a non-displayed chord.

Figure 2: Figure 3 from [13] illustrating how to use a z_H^* -reusable endpoint b of $f(x)$, which is covered by some vertex u and assigned to a chord xy of H , in order to insert a new vertex v that is adjacent to x and y . Note that the interval for u is split into two.

is adjacent to x and y we can define only one interval I that has at least two of the following three properties:

P1: $I \cap f(x) \neq \emptyset$, **P2:** $I \cap f(y) \neq \emptyset$, **P3:** I is displayed.

If xy is displayed, it is easy to find an interval I for v having properties **P1** and **P2**. Moreover if x , respectively y , has a broken end, it is easy to satisfy **P1**, respectively **P2**, and **P3**. However if xy is not displayed and neither x nor y has a broken end, Scheinerman and West propose to alter the existing representation of another vertex u whose interval covers an endpoint of x . As their modification, which is depicted in Figure 2, splits an interval of u into two, it is necessary that u appears at most twice so far, meaning that $f(u)$ consists of at most two intervals. This shall be achieved by specifying the current representation of G quite precisely.

Given a rooted plane embedded graph $G = (V, E)$, i.e., with a fixed root z_H for each block H of G^0 , a representation $\{f(v) \mid v \in V\}$ is called *P-special* if every vertex is displayed and each of the following holds:

- (1) Each root is represented by one interval and every other external vertex is represented by at most two intervals.
- (2) For each block H of G^0 all edges incident to z_H are displayed.
- (3) Each non-displayed edge xy of any block H of G^0 is assigned to an endpoint b of $f(x)$ or $f(y)$, say $f(x)$, such that the following hold.
 - (3.1) If xy is an external edge, the endpoint b is a broken end.
 - (3.2) If xy is a chord, the endpoint b is a broken end, or b is covered by $f(u)$ for only one other vertex u , where additionally u is on the z_H^* -side of xy and ux is an external edge of G^0 . An endpoint b satisfying this condition is called z_H^* -reusable for edge xy .
 - (3.3) For each endpoint b there is at most one edge assigned to b .
 - (3.4) Each vertex covers at most one endpoint which has an edge assigned.




Figure 3: A plane embedded 9-vertex graph G with one block H and root $z_H = x_1$, and how it is reduced to an independent set according to the induction rules in [13]. In each of the six steps, the set X is highlighted and the smaller graph is obtained by removing all vertices in X .

Now Scheinerman and West propose to show by induction on the number of vertices in G that every rooted plane embedded graph G admits a P-special 3-interval representation. They remove a small set of carefully chosen external vertices from G , induct on the smaller instance to obtain a P-special representation, create intervals for the removed vertex/vertices, extend and/or alter the existing representation, and argue that the result is a P-special 3-interval representation of G .

The problem lies in the last step. Some removed vertex v may cover an edge ux that is external in $G - v$, but a chord in G . But it may be that some non-displayed chord xy of $G - v$ is assigned to an endpoint b of $f(x)$ that is z_H^* -reusable and covered by u . This assignment cannot be kept, since ux is no longer external, which however is required for z_H^* -reusability as it is defined in (3.2) above. Thus, the invariants of the induction cannot be maintained.

Let us explain in more detail below (discussing only the relevant cases) on basis of a small example graph that the above problem can indeed occur, and why some straightforward attempt to fix it does not work. To this end, consider the plane embedded 9-vertex graph G in the top-left of Figure 3. The thick edges highlight the subgraph G^0 , which has only one block H , and let us pick vertex x_1 to be the root of that block, i.e., $z_H = x_1$. The base case of the inductive construction is an independent set of vertices. Otherwise, we identify a set X of external vertices that we want to remove as follows. Consider a leaf-block H of G^0 , i.e., one that contains no root of another block, its root z_H , and distinguish the following cases.

Case I: Every chord of H is incident to z_H . In this case label the vertices of H as z_H, v_1, \dots, v_k , $k \geq 1$, in this cyclic order around the outer face of H . If $k \geq 2$, let $X = \{v_1, \dots, v_p\}$, where p is the smallest integer greater than or equal to 2 for which v_p has degree 2 in H .

Case II: Some chord of H is not incident to z_H . Consider an inner face C of H that does not contain z_H and is bounded by only one chord xy , and distinguish further.

Case IIa: C is a triangle. Let $X = \{v\}$, where $v \neq x, y$ is the third vertex in C .

Case IIb: C has at least four vertices. In this case label the vertices of C as x, v_1, \dots, v_k, y , $k \geq 2$, in this cyclic order around C , and let $X = \{v_1, v_2\}$.

Figure 3 shows how our example graph G is reduced to an independent set according to these rules. Next, we consider these steps in reverse order and construct an interval representation following the case distinction as proposed in [13]. Let us refer to Figure 4 for a step-by-step illustration of this construction. The problem arises in Step 5.

Step 1: base case. Represent x_1 with a single interval.

Step 2: Case I, $X = \{x_2, x_3\}$. Insert intervals for x_1, x_2 into the displayed interval for $x_1 = z_H$. Place overlapping, displayed intervals for x_1, x_2 in an unused portion of the real line.

Step 3: Case IIa, $X = \{x_4\}$. As the chord $xy = x_2x_3$ is displayed, called subcase (1), add a displayed interval for $v = x_4$ and place the second interval available for v in the displayed portion for xy . Assign one broken end of v to each of the external edges $xv = x_2x_4$ and $yv = x_3x_4$.

Step 4: Case IIa, $X = \{x_5\}$. As the chord $xy = x_3x_4$ is not displayed, but assigned to a broken end b of $x = x_4$, called subcase (2), add the displayed interval for $v = x_5$ so that it overlaps $f(x)$ at b . Add a second interval for v in the displayed portion of $f(y)$. Assign the chord xy to the z_H^* -reusable endpoint b , which is here covered by $u = x_5$ with ux being an external edge.

Step 5: Case IIa, $X = \{x_6\}$. As the chord $xy = x_4x_5$ is displayed, we are in subcase (1) again. We add a displayed interval for $v = x_6$ and place a second interval for v in the displayed portion for xy . Now the problem is that the chord x_3x_4 can no longer be assigned to the endpoint b , since the edge x_4x_5 is no longer external.

One might be tempted to assume that the problem can be fixed by relaxing the definition of z_H^* -reusability as follows. Instead of ux being external, maybe it suffices to require that u appears at most twice so far. Let us call this new concept *almost z_H^* -reusability*. However, the assumption that ux is external is crucial for Case IIb, as we will demonstrate in Steps 6 and 7 below.

Step 6: Case IIb, $X = \{x_7, x_8\}$. Assign $v_1 = x_8$ an interval in the displayed interval for $x = x_3$, and assign $v_2 = x_7$ an interval in the displayed interval for $v_3 = x_6$. As the chord $xy = x_3x_4$ is not displayed, but assigned to an almost z_H^* -reusable endpoint b of $y = x_4$, we are in subcase (2c) of [13]. Here it is concluded in [13] that the vertex u covering b must be $u = v_k = x_6$ due to the externality of edge ux . Hence it would suffice to place overlapping displayed intervals for v_1 and v_2 in an unused portion of the line and add intervals for the inner neighbors of v_1, v_2 in the displayed portion for v_1, v_2 , and v_1v_2 , according to the subset of these vertices that they are adjacent to.




Figure 4: The steps of the inductive construction in [13] applied to the graph in Figure 3. Note that after Step 5 the chord x_3x_4 is not displayed and the right endpoint of the longer interval for x_4 is not z_H^* -reusable since edge x_4x_5 is not external. Further note that after Step 7 vertex x_5 is represented by four intervals.

However, in our example we have $u = x_5$, which at least appears only twice so far, but which is an inner neighbor of v_1 and v_2 , and thus has to spend its third interval in the displayed portion of v_1v_2 . We end up with chord xy still assigned to the endpoint b that is covered by u , but neither is ux external, nor is u appearing only twice so far. In particular, we are not prepared to insert a future vertex adjacent to x and y , as we illustrate in Step 7.

Step 7: Case IIa, $X = \{x_9\}$. The chord $xy = x_3x_4$ is not displayed, but assigned to the endpoint b of $x = x_4$, which is covered by $u = x_5$ with u however being internal and appearing three times already. Thus, if we apply the proposed modification as illustrated in Figure 2, we split one interval of u into two, causing $f(u)$ to consist of four intervals.

This concludes our example. Let us remark that the fourth interval for x_5 in Step 7 of Figure 4 might seem superfluous, but is actually needed to give u a displayed portion. This in turn is necessary as there might be a neighbor w of x_5 of degree 1. For example, with w inside face $x_5x_6x_7$, it would have been inserted between Steps 5 and 6, according to Case I with $z_H = x_5$, spending one interval in the displayed portion of u .

Also note that in this particular example other ways of assigning endpoints in Step 5 would have allowed the process to continue. However, the first author together with Daniel Gonçalves tried some time to obtain new invariants to fix this induction but did not succeed.

5 Concluding remarks

As mentioned above, Axenovich *et al.* [1] used some steps of the construction from Section 3 to prove that every graph G whose edges decompose into $k - 1$ forests and another forest of maximum degree 2 admits a k -interval representation of depth at most 2. Note that if G is a 4-connected triangulation with outer triangle Δ_{out} , the decomposition of $G - E(\Delta_{\text{out}})$ given by Lemma 2 can be easily extended to a decomposition of all edges in G into two forests and a path, which immediately gives the following.

Theorem 3. *If G is planar and 4-connected, then $i_2(G) \leq 3$.*

Given that the largest interval number among all planar graphs is 3, while the largest track number is 4, it remains open to determine whether the largest local track number among all planar graphs is 3 or 4, c.f., [10, Question 19]. We feel that the gluing of several triangulations along separating triangles is likely to be possible along the lines discussed in [10] for planar 3-trees. However, finding a 3-local track representation of a 4-connected planar triangulation, strengthening Theorem 3, seems to be more difficult. What is the largest local track number among all planar graphs?

Acknowledgments. We would like to thank Maria Axenovich and Daniel Gonçalves for fruitful discussions. We also thank the anonymous reviewers for their comments and suggestions. Moreover, we thank Ed Scheinerman and Douglas West for their helpful comments on an earlier version of this manuscript and

for providing us a copy of Ed's PhD thesis. The second author was partially supported by ANR projects ANR-16-CE40-0009-01 and ANR-17-CE40-0015 and by the Spanish Ministerio de Economía, Industria y Competitividad, through grant RYC-2017-22701.

References

- [1] M. Axenovich, A. Beveridge, J. P. Hutchinson, and D. B. West. Visibility number of directed graphs. *SIAM J. Discrete Math.*, 27(3):1429–1449, 2013.
- [2] J. Balogh, P. Ochem, and A. Pluhár. On the interval number of special graphs. *J. Graph Theory*, 46(4):241–253, 2004.
- [3] J. Balogh and A. Pluhár. A sharp edge bound on the interval number of a graph. *J. Graph Theory*, 32(2):153–159, 1999.
- [4] J. Chalopin and D. Gonçalves. Every planar graph is the intersection graph of segments in the plane. In *Proceedings of the forty-first annual ACM symposium on Theory of computing*, pages 631–638. ACM, 2009.
- [5] S. Chaplick and T. Ueckerdt. Planar graphs as VPG-graphs. *Journal of Graph Algorithms and Applications*, 17(4):475–494, 2013.
- [6] A. B. De Queiroz, V. Garnero, and P. Ochem. On interval representations of graphs. *Discrete Appl. Math.*, 202:30–36, 2016.
- [7] D. Gonçalves. Caterpillar arboricity of planar graphs. *Discrete Math.*, 307(16):2112–2121, 2007.
- [8] D. Gonçalves and P. Ochem. On star and caterpillar arboricity. *Discrete Math.*, 309(11):3694–3702, 2009.
- [9] M. Jiang. Recognizing d -interval graphs and d -track interval graphs. *Algorithmica*, 66(3):541–563, 2013.
- [10] K. Knauer and T. Ueckerdt. Three ways to cover a graph. *Discrete Math.*, 339(2):745–758, 2016.
- [11] K. Knauer and T. Ueckerdt. Decomposing 4-connected planar triangulations into two trees and one path. *J. Comb. Theory, Ser. B*, 134:88–109, 2019.
- [12] E. R. Scheinerman. *Intersection Classes and Multiple Intersection Parameters of Graphs*. PhD thesis, Princeton University, 1984.
- [13] E. R. Scheinerman and D. B. West. The interval number of a planar graph: Three intervals suffice. *J. Comb. Theory, Ser. B*, 35:224–239, 1983.
- [14] P. Stumpf. On covering numbers of different kinds. Bachelor thesis, Karlsruhe Institute of Technology, August 2015.
- [15] C. Thomassen. Interval representations of planar graphs. *J. Comb. Theory, Ser. B*, 40(1):9–20, 1986.

- [16] W. T. Trotter and F. Harary. On double and multiple interval graphs. *J. Graph Theory*, 3:205–211, 1979.
- [17] D. B. West and D. B. Shmoys. Recognizing graphs with fixed interval number is NP-complete. *Discrete Appl. Math.*, 8:295–305, 1984.